

## Journal of Plant Production

Journal homepage: [www.jpp.mans.edu.eg](http://www.jpp.mans.edu.eg)  
Available online at: [www.jpp.journals.ekb.eg](http://www.jpp.journals.ekb.eg)

### Improving Growth, Productivity and Oil Yield of *Nigella sativa*, L. Plants by Foliar Spraying with some Stimulants

Ayyat, A. M.<sup>1\*</sup>; A. G. M. Kenawy<sup>2</sup>; M. A. Aboel-Ainin<sup>3</sup> and M. A. M. Abdel-Mola<sup>4</sup>




<sup>1</sup>Department of Medicinal and Aromatic plants, Faculty of Agriculture, Beni-Suef University, Egypt.

<sup>2</sup>Medicinal and Aromatic Plants Res. Dept., Hort. Res. Inst., ARC, Giza, Egypt.

<sup>3</sup>Department of Biochemistry, Faculty of Agriculture, Beni-Suef University, Egypt.

<sup>4</sup>Department of Horticulture, Faculty of Agriculture, Beni-Suef University, Egypt.



#### ABSTRACT

A field experiment was conducted during the two successive seasons of 2018/2019 and 2019/2020 at the experimental farm of Horticultural Research Station at Sides, Beni-Suef Governorate to improve the growth, productivity and seed oil yield of black cumin plants to foliar application with  $\alpha$ -tocopherol (vitamin E), amino acids (Aminoactal) and moringa leaf extract (MLE). Obtained results revealed that the best vegetative growth criteria (Plant height, stem diameter, number of main branches and plant weight at harvest), yield parameters (number of capsules per plant, and seeds yield per plant and per fed.), oil determinations characters (volatile and fixed oil percentages and oil yield /plant and /fed.) and chemical properties such as photosynthetic pigments, herb percentages of N, P and K, free fatty acid contents, iodine and saponification values were obtained due to foliar spray of the high dose of moringa extract at 5 ml/l then amino acids 2ml/l and  $\alpha$ -tocopherol (vitamin E) at 100 ppm and with no significant differences being detected between such three concentration of the stimulants treatments. There for, it could be recommended that spraying *Nigella sativa* L. plants with the high concentration of moringa leaf extract (5 ml/l) to enhance growth, seeds yield / plant and /feddan and fixed oil components under the same environmental conditions.

**Keywords:** Black cumin, Moringa leaf extract (MLE),  $\alpha$ -tocopherol (Vitamin E), Amino acids, Volatile and fixed oil

#### INTRODUCTION

The black cumin (*Nigella sativa* L.) which belongs to family Ranunculaceae is a native herbaceous plant in the Mediterranean region and is one of the most valuable plants containing both volatile and fixed oil (Leung and Foster, 1996 and Burits and Bucar, 2000). In folk medicine, its seeds have been commonly used to treat a range of diseases, including diarrhea, jaundice, amenorrhea, helminthiasis, ophthalmia, paralysis and asthma. Black cumin seeds have a fixed oil rich in linoleic, oleic and palmitic acids (Al-Jassir, 1992 and Takruri and Dameh, 1998).

*Moringa oleifera* is a tree that can be called a miracle tree and is very useful. It has a strong role in various purposes, such as food, medicine and even in the industry (Moyo *et al.*, 2011). Moringa leaf extract (MLE) is known as a natural plant stimulant that, when applied as a foliar spray, improves yield. The active ingredients for promoting growth in MLE are stated to be zeatin, dihydrozeatin and isopentyladenine (Andrews, 2006 and Price, 2007). Leaves of moringa which collected from all over the world and discovered to have high zeatin concentrations of 5 to 200  $\mu$ g/g of leaves (Fuglie, 2000). *Moringa oleifera* has also drawn immense attention because its leaves contain antioxidants such as ascorbic acid, flavonoids, phenolics, carotenoids, amino acids, macro and micro nutrients (Foidl *et al.*, 2001 and Yasmeen, 2011).

Amino acids are considered as precursors of proteins, which are vital for stimulating the growth of cells. They have a number of functions in plants, including regulatory and signaling molecules. Also, influence gene expression, redox homeostasis and the synthesis and activity of certain enzymes. Glycine and glutamic acids are essential for the development

of vegetative growth and chlorophyll contents. They also chelate micronutrients, making them easier for plants to absorb and transport (Rai, 2002).

Vitamins are bio-regulator compounds that have profound effects on plant growth controlling factors that control many physiological processes, such as enzyme synthesis, function as co-enzymes and affect plant growth at relatively low concentrations (Abdel Aziz *et al.*, 2009). Tocopherols are a class of compounds generated exclusively by photosynthetic organisms that are concerned in the quenching and scavenging of oxygen (Neely *et al.*, 1988) and serve as extremely effective recyclable chain reaction terminators for the removal of polyunsaturated fatty acid (PUFA) radical species produced during lipid peroxidation (Munne-Bosch and Falk, 2004).

The aim of this study to investigate the response of *Nigella sativa* L. plants to foliar application of MLE, amino acids and  $\alpha$ -tocopherol (vitamin E).

#### MATERIALS AND METHODS

The present study was conducted in the experimental farm of the Horticultural Research Station at Sides, Beni-Suef Governorate, over two successive seasons of 2018/2019 and 2019/2020. Prior to land preparation, several soil samples (0-40 cm depth) were taken for nutrient and trace element analysis (Table, 1) according to, Jackson, 1973.

The black cumin seeds were sown on the third week of October in both seasons in 3 X 3.60 m plots with 60 cm distance between the rows and 30 cm between hills within each row. So, each plot contained 6 rows and 60 hills. The plants were thinned to two plants per hill after one month of sowing. In addition, all experimental plots, including control, received 20 Kg/ plot of farm

\* Corresponding author.

E-mail address: [Ahmed.Ayyat@agr.bsu.edu.eg](mailto:Ahmed.Ayyat@agr.bsu.edu.eg)

DOI: 10.21608/jpp.2021.69108.1024

yard manure (FYM) before sowing date and all other agricultural practices were followed as usual. FYM was received from Sides Research Station belonging to Animal Production Research Institute (APRI), ARC, Egypt. Table (2), clearly show that some chemical properties of the used FYM at the both seasons, were determined according the method described by A.O.A.C., (1995).

Complete randomized block design (CRD) was also used in this investigation. The foliar sprayed for all treatments were done three times, the first sprayed was applied after 45 days from sowing date and repeated two times at 15 days intervals through the both growing season .The plants were harvested in the third week of April in both seasons.

**Preparation of MLE**

100 g of young fresh moringa leaves were collected and combined with 1 litre of distilled water to prepare an aqueous extract. A household blender was used to homogenies the suspension for 15 minutes. The solution was then filtered by squeezing with a mutton cloth. Finally, using No. 2 Whatman filter paper, the solution were re-filtered (Fuglie, 2000). Plants were sprayed directly with the extract. The extract was sprayed within 5 hours of the collection and extraction process (if not ready for use, the prepared extract was stored at 0.0 °C, it is only taken out when it is needed for use) Abdel-Rahman and Abdel-Kader (2020).

**Table 1. Chemical and physical properties of the experimental soil at 2018 and 2019.**

Seasons	Particle size distribution*			Textural Class	OM %	Chemical properties**							pH
	Clay %	Silt %	Sand %			EC, dSm <sup>-1</sup> (at 25°C)	N	P	Available (ppm)			Mn	
2018/2019	49.80	33.20	17.00	Clay	1.75	1.12	41.00	14.18	260.5	2.40	0.25	0.60	7.7
2019/2020	48.10	33.60	18.30	Clay	1.60	1.03	38.00	12.60	244.6	2.25	0.30	0.65	7.8

**Table 2. Chemical properties of the used FYM (2018 and 2019).**

Parameter	Total N (%)	Total P (%)	Total K (%)	Organic matter (%)	Organic carbon (%)	C/N ratio	pH (1: 10 soil water suspension)	EC, dSm <sup>-1</sup> (1:10 soil water extraction)
2018	1.42	0.31	1.05	45.22	26.23	1:18	7.82	4.75
2019	1.46	0.29	1.11	47.31	27.44	1.19	7.90	4.83

**Amino acids source**

The commercial product "Aminoactal" was used that contains the following amino acids mixture (Methionine, Cysteine and Treptophan) which was obtained from Shoura Chemical Company, Cairo-Alexandria Desert Road, Giza Governorate.

**Recorded data**

**A- Vegetative growth criteria**

Plant height, stem diameter, main branch number/ plant and plant weight at harvest.

**B- Yield components**

Number of capsules per plant, seeds yield per plant and per feddan (calculated as plot yield × plots number).

**C- Oil productivity**

Volatile oil percentage and yield/feddan, according to (Guenther, 1961), fixed oil percentage and yield/feddan, according to (Lo´pez-Basco´ and de Castro, 2020).

**D- Chemical properties**

The three photosynthetic pigments were determined according to Moran (1982), N, P and K percentages were determined according to Wilde *et al.* (1985), Chapman and Pratt (1975) and Cottenie *et al.* (1982), respectively, the free fatty acid (FFAs), iodine value (IVs) and saponification value (SVs) of black cumin seed oil were measured as general indicator of the oil's condition and edibility, according to A.O.A.C. (1990).

The data were statistically analyzed using the MSTAT-C (1985). the differences between means were tested using the least significant differences (L.S.D) test at 0.05.

**RESULTS AND DISCUSSION**

**A- Vegetative growth criteria**

Obtained results in Table (3) for plant height, stem diameter, main branch number/plant and plant weight at harvest of black cumin plants showed that all tested treatments, namely, α-tochopherol (50 and 100 ppm), amino acids (1 and 2 ml/l) and MLE (3 and 5 ml/l) caused a substantial and significant increase in the four tested vegetative characters in the two seasons, in comparison with untreated treatment. The highest overall values were given due to MLE at 5 ml/l followed by amino acids at 2 ml/l then α-tochopherol at 100 ppm, in the two growing seasons, in most cases. The increase in plant height, stem diameter, branch number/plant and plant weight at harvest due to MLE at 5 ml/l, in comparison with control treatment reached 36.65, 37.31, 39.29 and 51.38 % , respectively, in the first season and by 38.00, 37.68, 40.29 and 41.59 % , respectively, in the second one. The role of MLE in enhancing vegetative growth characters was demonstrated by Mostafa (2015) on fennel, Ali *et al.* (2018) on geranium, meanwhile, The importance of amino acids was highlighted by Zedan (2000) on Coriander and Caraway plants and Gamal El-Din *et al.* (2005) on chamomile plant. Vitamins also increased the height and diameter of coriander plants' stems, as defined by Abdou *et al.* (2015).

**Table 3. Effect of the three stimulants treatments on some vegetative growth criteria of black cumin plants during 2018/2019 and 2019/2020.**

characters	Plant height (cm)		Stem diameter (cm)		Branch number per plant		Plant weight at harvest (g)	
	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season
Seasons								
Treatments								
Control (water)	41.2	40.0	0.67	0.69	5.60	6.13	26.47	26.93
Vitamin E 50 ppm	46.5	45.6	0.78	0.81	6.20	7.20	31.93	31.07
Vitamin E 100 ppm	50.6	51.0	0.88	0.91	7.27	8.03	37.53	36.20
Amino acids 1 ml/l	48.6	48.9	0.84	0.87	6.87	7.63	35.47	33.80
Amino acids 2 ml/l	54.2	51.4	0.89	0.93	7.33	8.20	39.73	37.40
MLE 3 ml/l	54.5	52.3	0.86	0.88	6.80	7.73	37.00	36.07
MLE 5 ml/l	56.3	55.2	0.92	0.95	7.80	8.60	40.07	38.13
L.S.D 5%	3.7	4.0	0.07	0.04	0.75	0.94	2.39	1.93

These results may be explained by a high level of endogenous cytokinin (zeatin, dihydrozeatin, and isopentyladenine) in MLE (Fuglie, 2000, Andrews, 2006 and

Price, 2007). Zeatin is a type of cytokinin naturally found in plants, and its role is not only necessary in cell division and elongation which stimulates plant growth, but it also has

defensive properties on plants (Nagar *et al.*, 2006 and Anwar *et al.*, 2007). Moringa leaf extract is also high in Ca, P, Mg, Fe, Zn, Mn. It also contains other growth-promoting compounds, making it a natural plant growth promoter (Makkar *et al.*, 2007, Jacob and Shenbagaraman, 2011 and Abd El-Hamied and El-Amari, 2015). Tao *et al.* (2008) observed an increase in plant growth criteria after amino acid application as a consequence of synthesis of protein. Amino acids are precursors of phytohormones and growth factors (El-Awadi and Hassan, 2010). The beneficial effect of  $\alpha$ -tocopherol may be due to induced improvements in growth characteristics such as better water capacity, high antioxidant accumulation, less oxidative harm, and improved cross-talk among different growth regulators (Soltani *et al.*, 2012).

**B- Yield components**

Table (4) showed that there were statistically significant differences, in both seasons, between the control treatment on the one hand and the other treatments examined ( $\alpha$ -tocopherol, amino acids and MLE) except for the  $\alpha$ -tocopherol (50 ppm) treatment, for seeds yield per plant. Among the six tested treatments, the three treatments of MLE at 5 ml/l, amino acids at 2 ml/l and  $\alpha$ -tocopherol at 100 ppm resulted in the best values for number of capsules / plant, seeds yield / plant and / feddan without significant differences being obtained between them, except for seeds yield/fed. for second season. They increased the number of capsules per plant of black cumin plants, in comparison with check treatment, by 36.76%, 24.66% and 23.65%, respectively, in the first season and by 39.65%, 30.80% and 29.75%, respectively, in the second season. Numerically, the rise in seeds yield / plant and / fed. reached 38.15, 31.11 and 25.07 % for seeds yield / plant and by 37.90, 32.59 and 30.98 % for seeds yield / fed. in the first season, respectively, over that check treatment. The corresponding values in the second season were obtained for the same treatments, for seeds yield / plant 44.27, 33.36 and 29.73 % and by 44.74, 37.59 and 36.33 % for seeds yield / fed., respectively.

**C- Oil productivity**

Oil productivity of black cumin plants were greatly and significantly increased due to the use of six tested treatments, the only exceptional for vitamin E (50 ppm) for volatile and fixed oil

percentages compared to control treatments in both experimental seasons as shown in Table (5). The highest volatile and fixed oil percentages among these five treatments was obtained from MLE at 5 ml/l treatment, followed by amino acids at 2 ml/l treatment, then,  $\alpha$ -tocopherol (100 ppm) treatment which increased volatile oil percentage over check treatment by 31.71, 25.61 and 23.17 %, respectively, in the first season and by 25.88, 24.71 and 21.18 %, respectively, in the second one, while, the increments of fixed oil percentage of seeds due to these three treatments over control treatment reached 18.44, 18.01 and 12.32 %, respectively, in the first season and by 17.09, 14.43 and 11.76 % respectively, in the second one. Numerically, the rise in volatile and fixed oil yield / feddan for these three treatments was 73.31, 59.55 and 58.71 % for volatile oil and by 71.76, 59.88 and 55.52 % for fixed oil in the first season, respectively, over the control treatment. The corresponding values, for the same treatments in the second season came to 61.86, 50.52 and 48.45 % for volatile oil and by 70.73, 58.99 and 53.87 %, for fixed oil, respectively, as shown in Table (5). In accordance with the results which explored the capability of MLE, amino acids and vitamin E in promoting yield components and oil productivity of different plant species were the findings by Abdou *et al.* (2015) on coriander, Abdel-Rahman and Abdel-Kader (2020) on fennel and Poorghadir *et al.* (2020) on savory.

**Table 4. Effect of the three stimulants treatments on yield components of black cumin plants during 2018/2019 and 2019/2020.**

characters	Number of capsules per plant		Seeds yield per plant (g)		Seeds yield per feddan (Kg)	
	1st season	2nd season	1st season	2nd season	1st season	2nd season
Control (water)	26.93	24.87	10.93	11.00	442.97	438.50
Vitamin E 50 ppm	32.40	29.40	12.17	12.63	520.13	520.67
Vitamin E 100 ppm	33.30	32.27	13.67	14.27	580.20	597.83
Amino acids 1 ml/l	31.47	28.93	12.87	13.40	542.73	548.10
Amino acids 2 ml/l	33.57	32.53	14.33	14.67	587.33	603.33
MLE 3 ml/l	32.70	30.33	13.73	14.00	541.53	560.13
MLE 5 ml/l	36.83	34.73	15.10	15.87	610.87	634.67
L.S.D 5%	4.14	2.84	2.03	1.68	65.36	40.64

**Table 5. Effect of the three stimulants treatments on oil productivity of black cumin plants during 2018/2019 and 2019/2020.**

characters	Volatile oil % of seeds		Volatile oil yield L. per feddan		Fixed oil % of seeds.		fixed oil yield L. per feddan	
	1st season	2nd season	1st season	2nd season	1st season	2nd season	1st season	2nd season
Control (water)	0.82	0.85	3.56	3.88	28.09	28.49	98.70	103.17
Vitamin E 50 ppm	0.89	0.91	4.32	4.68	29.34	29.86	133.13	136.57
Vitamin E 100 ppm	1.01	1.03	5.65	5.76	31.55	31.84	153.50	158.75
Amino acids 1 ml/l	0.97	0.94	4.79	4.87	31.08	31.29	134.65	136.87
Amino acids 2 ml/l	1.03	1.06	5.68	5.84	33.15	32.60	157.80	164.03
MLE 3 ml/l	0.99	1.01	5.33	5.48	31.09	31.72	150.92	162.13
MLE 5 ml/l	1.08	1.07	6.17	6.28	33.27	33.36	169.53	182.33
L.S.D 5%	0.08	0.05	0.54	0.60	2.51	1.97	24.67	28.29

MLE improves crop yield mainly due to the presence of zeatin and other hormones and also a variety of active compounds (Duke, 1992). When zeatin-related hormone is applied as an extract from fresh ML, it enhances crop yields, according to Fuglie (2000). Furthermore, the simulative impact of such treatments on enhancing characteristics of seeds, essential and fixed oil percentages and oil yield may be due to their improving impact on vegetative growth criteria and plant chemical properties for example, NPK in black cumin plants, which can affect seeds and oil yield. The influence of amino acids may be attributed to their functions in improving a variety of physiological processes, such as nutrient absorption by roots and

metabolism in treated plants (Hanafy *et al.*, 2010). Also, their role in changes in levels and/or enzymes involved. Furthermore, micronutrients, especially iron and Zinc, serve as metal components of a variety of enzymes as well as functional, structural, and regulatory cofactors. Sadak *et al.* (2010) on sunflower and Soltani *et al.* (2012) on calendula plants stated that foliar spray of  $\alpha$ -tocopherol (vitamin E) led to increase seed yield. The impact of vitamins on protein synthesis and delaying senescence is primarily responsible for the increment in black cumin seed yield as a result to foliar application of vitamin E, but it could also be linked to an improve in photosynthetic products.

**D- Chemical properties**  
**Photosynthetic pigments**

The response of photosynthetic pigment content to various concentrations of MLE, amino acids, and  $\alpha$ -tocopherol, as shown in Table (6), indicates that these treatments were found to be positively affected compared to the check treatment in both growing seasons. The exceptional were the treatments of  $\alpha$ -tocopherol at 50 ppm for the three photosynthetic pigments and amino acids at 1 ml/l for carotenoids content in the two growing seasons. The highest contents of photosynthetic pigments were obtained from the following treatments in descending order, MLE at 5 ml/l followed by the amino acids at 2 ml/l then the  $\alpha$ -tocopherol at 100 ppm. These three treatments increased the photosynthetic pigments over those of control plants by 14.87, 10.94 and 9.40 % in the first season and by 14.31, 10.52 and 9.86 % in the second one, respectively, for chlorophyll a, and by 19.42, 17.63 and 16.29 % in the first season and 19.16, 18.28 and 17.40 % in the second one, successively, for chlorophyll b, while for carotenoids, by 10.31, 7.80 and 6.48 % in the first season and 9.27, 7.84 and 6.70 % in the second one, consecutively. These results of this study back up the findings of the other studies such as Aslam *et al.* (2016) on spinach and Abdou *et al.* (2018) on gladiolus.

**Table 6. Effect of the three stimulants treatments on photosynthetic pigments of black cumin plants during 2018/2019 and 2019/2020.**

characters	Chlorophyll (a) content (mg / g F.W)		Chlorophyll (b) content (mg / g F.W)		Carotenoids content (mg / g F.W)	
	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season
Control (water)	1.042	1.055	0.448	0.454	0.679	0.701
Vitamin E 50 ppm	1.101	1.112	0.485	0.499	0.685	0.715
Vitamin E 100 ppm	1.140	1.159	0.521	0.533	0.723	0.748
Amino acids 1 ml/l	1.126	1.138	0.495	0.525	0.705	0.725
Amino acids 2 ml/l	1.156	1.166	0.527	0.537	0.732	0.756
MLE 3 ml/l	1.135	1.150	0.518	0.522	0.715	0.744
MLE 5 ml/l	1.197	1.206	0.535	0.541	0.749	0.766
L.S.D 5%	0.082	0.059	0.044	0.051	0.032	0.027

The existence of significant quantities of chlorophyll and carotenoids in moringa leaf extract could explain its positive impact on photosynthetic pigments (Ashfaq *et al.*, 2012). Moringa leaves also contain a variety of components, including Mg (Yameogo *et al.*, 2011), a chlorophyll component, as well as cytokinins, which play an important role in biosynthesis of chlorophyll (Taiz and Zeiger, 2010) therefore, the quantity of photosynthetic pigments in black cumin herb were increased.

The beneficial effect of amino acids on chlorophylls content of black cumin plants can be explained simply because of their crucial function in chlorophyll biosynthesis. Saburi *et al.* (2014) found that foliar tryptophan application resulted in an increase in total chlorophyll and carotenoids in basil plants. This stimulatory influence of  $\alpha$ -tocopherol could be attributed to their antioxidant scrounging impact, which preserved chloroplasts and inhibited chlorophyll deterioration by toxic reactive oxidative damage (Bowler *et al.*, 1992 and Aono *et al.*, 1993).

**N, P and K percentages**

The percentages of N, P and K in the dry herb of black cumin plants was significantly affected by different treatments (MLE, amino acids, and  $\alpha$ -tocopherol at 100 ppm) in both seasons, according to data provided in Table (7). Plants treated with the two concentrations of MLE yielded the highest percentages of the three elements over those of the control treatment by 27.90 and 21.45 % in the first season and

by 29.92 and 25.34 % in the second one, respectively for N % and by 20.62 and 17.90 % in the first season and 16.73 and 14.87 % in the second one, successively for P %, as well as, by 6.63 and 5.72 % in the first season and by 12.68 and 10.13 % in the second one, respectively for K %. The above-mentioned findings on the benefits of vitamin E, amino acids and MLE on N, P and K % in the black cumin herb were on line with those of Mady (2009) on tomato, Abd-Elkader *et al.* (2020) on gerbera and Abdel-Rahman and Abdel-Kader (2020) on fennel, respectively.


The presence of amino acids, P, Ca, Fe, vitamin E, ascorbates, phenolic compounds and growth regulating hormones like zeatin in MLE can explain the increase in the three photosynthetic pigments and the NPK (Siddhuraju and Becker, 2003 and Nagar *et al.*, 2006). Moringa leaf extract contains Ca and P, which may play a vital role in crop growth and development through osmoregulation, enzyme activation, photosynthesis and different physiological processes (Hasegawa *et al.*, 2000 and Epstein and Bloom, 2005). The amino acids may have had a beneficial effect because of their essential function in the chlorophyll molecules biosynthesis, which reflected NPK percentages. Furthermore, antioxidants like vitamin E protected cell membranes and their binding transporter proteins (H<sup>+</sup>-ATPase membrane pumps), preserving their structure and function against the toxic and destructive effects of reactive oxygen species during stress, resulting in increased mineral absorption and translocation (Dickson *et al.*, 1991).

**Free fatty acid, iodine value and saponification value**

The results for free fatty acid contents, iodine value and saponification value as shown in Figure (1), were ranked in order of transactions (amino acids,  $\alpha$ -tocopherol, MLE and control) in descending order of transactions. FFAs were 14.25, 13.75, 13.25 and 12.25, respectively., IVs were (102, 109, 110 and 113 g I2/100 g oil), successively, and for SVs were (206.25, 205.5, 205.25 and 197), consecutively.

**Table 7. Effect of the three stimulants treatments on NPK percentages of black cumin plants during 2018/2019 and 2019/2020.**

characters	Nitrogen%		Phosphorus%		Potassium%	
	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season	1 <sup>st</sup> season	2 <sup>nd</sup> season
Control (water)	1.147	1.093	0.257	0.269	1.977	2.003
Vitamin E 50 ppm	1.220	1.207	0.284	0.292	2.010	2.077
Vitamin E 100 ppm	1.283	1.297	0.293	0.304	2.070	2.186
Amino acids 1 ml/l	1.237	1.280	0.279	0.293	1.996	2.067
Amino acids 2 ml/l	1.371	1.349	0.297	0.308	2.073	2.198
Moringa ex. 3 ml/l	1.393	1.370	0.303	0.309	2.090	2.206
Moringa ex. 5 ml/l	1.467	1.420	0.310	0.314	2.108	2.257
L.S.D 5%	0.119	0.131	0.019	0.017	0.055	0.087



**Figure 1. Effects of treatments on free fatty acid, iodine value and saponification value of Nigella sativa seed oil**

\* (AA) Amino Acid, ( $\alpha$ -T)  $\alpha$ -Tocopherol, (MLE) Moringa Leaf Extract, (FFA) Free Fatty Acids, (IV) Iodine Value and (SV) Saponification Value.

## CONCLUSION

The important consideration of this study is the use environmental stimulants to improving the seed yield and fixed oil quality of the black cumin plants. Under the environmental conditions of same study, the highest value of the growth characters, seed yield and fixed oil of *Nigella sativa* plants, were obtained due to foliar spray of moringa leaf extract (MLE), amino acids and  $\alpha$ -tocopherol at high concentration use.

## REFERENCES

- Abd El-Hamied, S.A. and E.I. El-Amary (2015). Improving growth and productivity of pear trees using some natural plants extracts under North Sinai conditions. IOSR Journal of Agriculture and Veterinary Science 8, 1-9.
- Abdel Aziz, N.G., L.S. Taha and S.M.M. Ibrahim (2009). Some studies on effect of putrescine, ascorbic acid and thiamine on growth, flowering and some chemical constituents of gladiolus plants at Nurbaria. Ozean J. Appl. Sci., 2(2): 169-179.
- Abd-Elkader, H.H., H.Y. Massoud, T.T. El-Baz and M.A. El-Erian (2020). Effect of Amino Acids Spray on Growth, Flowering and Keeping Quality of *Gerbera jamesonii* L. as a Pot Plant. J. of Plant Production, Mansoura Univ., 11 (4):333 - 339.
- Abdel-Rahman, S.S.A. and A.A.S. Abdel-Kader (2020). Response of Fennel (*Foeniculum vulgare*, Mill) plants to foliar application of moringa leaf extract and benzyladenine (BA). South African J. of Botany, 129: 113-122.
- Abdou, M.A.H., A.A. El-Sayed, E.T. Ahmed and A.A.M. Abdel Salam (2015). Effect of compost, mineral NPK, effective microorganisms and some vitamin treatments on growth, fruit yield and essential oil content of coriander (*Coriandrum sativum*, L.) plants. Scientific J. Flowers & Ornamental Plants, 2 (3): 203-212.
- Abdou, M.A.H., F.S. Badran, E.T. Ahmed, R.A. Taha and M.A.M. Abdel-Mola, (2018). Effect of compost and some natural stimulant treatments on: II. Corms production and chemical constituents of (*Gladiolus grandiflorus* cv. Peter Pears) plants. Proc. of The 4<sup>th</sup> Conf. of SSSFOP, Scientific J. Flowers & Ornamental Plants, 5(2):115-126.
- Ali, E.F., F.A.S. Hassan and M. Elgimabi (2018). Improving the growth, yield and volatile oil content of *Pelargonium graveolens* L. Herit by foliar application with moringa leaf extract through motivating physiological and biochemical parameters. South African J. of Botany, 119: 383-389.
- Al-Jassir, S.M. (1992). Chemical composition and microflora of black cumin (*Nigella sativa*) seeds growing in Saudi Arabia. Food Chem, 45: 239 - 242.
- Andrews, D. (2006). Nutraceutical moringa composition. <http://www.google.com/patents/US20060222882/>.
- Anwar, F., S. Latif, M. Ashraf and A.H. Gilani (2007). *Moringa oleifera*: a food plant with multiple medicinal uses. Phytotherapy Res., 21: 17-25.
- A.O.A.C. (1990). Association of Official Analytical Chemists. Official Method of Analysis 15<sup>th</sup> Edition, Washington, D.C., USA.
- A.O.A.C. (1995). Association of Official Agricultural Chemists. Official Method of Analysis 14<sup>th</sup> Edition, Washington, A.O.A.C., D.C., USA. 490-510.
- Aono, M., A. Kubo, H. Saji, K. Tanaka and N. Kondo (1993). Enhanced tolerance to photo-oxidative stress of transgenic (*Nicotiana glauca*) with high chloroplastic glutathione reductase activity. Plant Cell Physiol., 34:129-135.
- A.R.C. (2005). Agriculture Research Center. Technical Bulletin No. 984, Cairo, Egypt.
- Ashfaq, M., S. Basra and U. Ashfaq (2012). Moringa: A miracle plant of agro-forestry. Journal of Agriculture and Social Sci., 8: 115-122.
- Aslam, M., B. Sultana, F. Anwar and H. Munir (2016). Foliar spray of selected plant growth regulators affected the biochemical and antioxidant attributes of spinach in a field experiment. Turkish J. of Agriculture and Forestry, 40: 136-145.
- Bowler, C., M.V. Montogu and D. Inze (1992). Superoxide dismutase and stress tolerance. Ann. Rev. Plant Physiol. Plant Mol. Biol., 48: 223-250.
- Burits, M. and F.B. Bucar (2000). Antioxidant activity of *Nigella sativa* essential oil. Phytother. Res., 14: 323 - 328.
- Chapman, H.D. and R.E. Pratt (1975). Methods of Analysis for Soil, Plant and Water. California Univ., Division of Agric. Sci., p.172-173.
- Cottenie, A., M. Nerloo, G. Velghe and L. Kiekens (1982). Biological and analytical aspects of soil pollution. Lab. of Analytical Agro. State Univ. of Calif. Division of Agric. Sci.: 60-69.
- Dickson, C.D., T. Altabella and M.J. Chrispeels (1991). Slow growth phenotype of transgenic tomato expressing plastic invertase. Plant Physiol., 95: 420-425.
- Duke, J.A. (1992). Handbook of Phytochemical Constituents of GRAS Herbs and Other Economic Plants. CRC Press, Boca Raton, Florida.
- El-Awadi, M.E. and E.A. Hassan (2010). Physiological responses of fennel (*Foeniculum vulgare* Mill) plants to some growth substances. The effect of certain amino acids and a pyrimidine derivative. J. Am. Sci., 6: 120-124.
- Epstein, E. and A.J. Bloom (2005). Mineral Nutrition on Plants: Principles and Perspectives. 2<sup>nd</sup> ed. Sinauer Associates, Sunderland, MA, USA.
- Foidl, N., H.P.S. Makkar and K. Becker (2001). The potential of *Moringa oleifera* for agricultural and industrial uses. In: The Miracle Tree: The Multiple Attributes of Moringa, pp: 45-76. (Lowell J. Fuglie, ed.), CTA, Wageningen The Netherlands.
- Fuglie, L.J. (2000). The Miracle Tree: *Moringa oleifera*: Natural Nutrition for the Tropics. The Multiple Attributes of Moringa. p. 172.
- Gamal EL-Din, K.M. and M.S.A. Abd EL-Wahed (2005). Effect of some amino acids on content of chamomile plant. Inter. J. of Agri. Biol., (7): 376-380.
- Guenther, E. (1961). The Essential Oils, 4<sup>th</sup> Ed., Vol. III. D. Van Nostrand Comp., New York, London.
- Hanafy, A.A.H., M.R. Nesiem, A.M. Hewedy and H.S. Sallam (2010). Effect of some simulative compounds on growth, yield and chemical composition of snap bean plants grown under calcareous soil conditions. J. of American Sci., 6 (10): 552-569.
- Hasegawa, P.M., R.A. Bressan, J.K. Zhu and H.J. Bohnert (2000). Plant cellular and molecular responses to high salinity. Annual Review of Plant Physiology, (51): 463-499.
- Jackson, M.L. (1973). Soil Chemical Analysis. Prentice-Hall of India Private Ltd. M-97, New Delhi, India, 498 p.
- Jacob, S.J.P. and S. Shenbagaraman (2011). Evaluation of antioxidant and antimicrobial activities of the selected green leafy vegetables. International J. of Pharm Tech Res., 3: 148-152.

- Leung, A.Y. and S. Foster (1996). Encyclopedia of Common Natural Ingredients Used in Food, Drugs and Cosmetics (2nd edn). Wiley, New York.
- Lopez-Basco, M.A., and M.L. de Castro (2020). Soxhlet Extraction. In Liquid-Phase Extraction. Elsevier, pp. 327-354.
- Mady, M.A. (2009). Effect of foliar application with salicylic acid and vitamin E on growth and productivity of tomato (*Lycopersicon esculentum*, Mill.) plant. J. Agric. Sci. Mansoura Univ., 34 (6): 6715-6726.
- Makkar, H.P.S., G. Francis and K. Becker (2007). Bioactivity of phytochemicals in some lesser known plants and their effects and potential applications in livestock and aquaculture production systems. Animal, 1 (9): 1371-1391.
- Moran, R. (1982). Formula determination of chlorophyll pigment extracted with N,N diethyl formamide. Plant Physiol., 69: 1376-1381.
- Mostafa, G.G. (2015). Improving the growth of fennel plant grown under salinity stress using some biostimulants. American J. of Plant Physiology 10 (2): 77-83.
- Moyo, B., P.J. Masika, A. Hugo and V. Muchenje (2011). "Nutritional characterization of Moringa (*Moringa oleifera* Lam) leaves," African J. of Biotechnology, 10 (60): 12925-12933.
- MSTAT-C (1986): A Microcomputer Program for the Design, Management and Analysis of Agronomic Research Experiments (Version 4.0). Michigan State Univ., U.S.A.
- Munne-Bosche, S. and J. Falk (2004). New insights into the function of tocopherols in plant. Planta, 218: 323-326.
- Nagar, P.K., R.I. Leyer and P.K. Sircar (2006). Cytokinins in developing fruits of Moringa pterigosperma Gaertn. Plant Physiology, 55, 45-50.
- Neely, W.C., J.M. Martin and S.A. Barker (1988). products and relative reaction rates of the oxidation of tocopherols with singlet molecular oxygen. Photochemistry and Photobiology. 48 (4): 423-428.
- Poorghadir, M., A.M. Torkashvand, S.A. Mirjalili and P. Moradi, (2020). Interactions of amino acids (proline and phenylalanine) and biostimulants (salicylic acid and chitosan) on the growth and essential oil components of savory (*Satureja hortensis* L.). Biocatalysis and Agricultural Biotechnology, 30, 101815
- Price, M. (2007). The Moringa Tree. ECHO Technical Note, eBook. Available on <http://www.miracle-trees.org/> Moringa- doc/ebook Moringa.pdf.
- Rai V.K. (2002). Role of amino acids in plant responses to stress. Biol Plant. 45: 471-478.
- Saburi, M.R., H. Sayed, S. Mohammad and D. Taghi (2014). Effects of amino acids and nitrogen fixing bacteria on quantitative yield and essential oil content of basil (*Ocimum basilicum*). Agric.Sci. Dev, (8): 265-268.
- Sadak, M.Sh., M.M. Rady, N.M. Badr and M.S. Gaballah (2010). Increasing sunflower salt tolerance using nicotinamide and  $\alpha$ -tocopherol. Int. J. Acad. Res., 2 (4): 263-270.
- Siddhuraju, P. and K., Becker, (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick tree (*Moringa oleifera* Lam). J. of Agriculture and Food Chem., 15, 2144-2155.
- Solati, Z., B.S. Baharin and H. Bagheri (2014). Antioxidant property, thymoquinone content and chemical characteristics of different extracts from *Nigella sativa* L. seeds. J. Am. Oil Chem. Soc., 91 (2): 295-300.
- Soltani, Y., V.R. Saffari, A.A.M. Moud and M., Mehrabani (2012). Effect of foliar application of  $\alpha$ -tocopherol and pyridoxine on vegetative growth, flowering, and some biochemical constituents of *Calendula officinalis* L. plants. African J. of Biotechnology, 11(56): 11931-11935.
- Taiz, L. and E. Zeiger (2010). Plant Physiology. 5th ed. Sinauer Associates, Sunderland, MA, USA.
- Takruri, H.R.H. and M.A.F. Dameh (1998). Study of nutritional value of black cumin seeds (*Nigella sativa* L.). J. Sci. Food Agric., 76: 404 – 410.
- Tao, Y., J.L. Ferrer, K. Ljung, F. Pojer and F. Hong (2008). Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants. Cell, 133 (1): 164-176.
- Wilde, S.A., B.B. Gorey, J.G. Layer and J.K. Voigt (1985). Soils and Plant Analysis for tree culture. Published by Mohan Primlani, Oxford and IBH publishing Co., New Delhi, p. 1-142.
- Yamego, C.W., Bengaly, M.D., Savadogo, A., Nikiema P.A., Traore S.A. (2011). Determination of chemical composition and nutritional values of Moringa oleifera leaves. Pak. J. Nutr. 10(3):264-268.
- Yasmeen, A. (2011). Exploring the Potential of Moringa (*Moringa oleifera*) Leaf Extract as Natural Plant Growth Enhancer. Ph.D. Thesis, Fac. Agric., Univ. of Agric., Faisalabad, Pakistan, 150 p.
- Zedan, H.M.Z. (2000). Effect of tryptophan and paclobutrazol on caraway (*Carum carvi* L.) and coriander (*Coriandrum sativum* L.) plants. M.Sc. Thesis, Fac. Agric, Cairo Univ., Egypt, 150 p.

**تحسين النمو والانتاجية ومحصول الزيت لنباتات حبة البركة بالرش الورقي ببعض المنشطات**  
**أحمد محمد عياط<sup>1\*</sup>، عادل جلال محمود قناوي<sup>2</sup>، مصطفى عبد المنعم ابو العنين<sup>3</sup> و مصطفى عبده محمود عبد المولى<sup>4</sup>**

<sup>1</sup> قسم النباتات الطبية والعطرية - كلية الزراعة - جامعة بني سويف

<sup>2</sup> قسم بحوث النباتات الطبية والعطرية - معهد بحوث البساتين - مركز البحوث الزراعية

<sup>3</sup> قسم الكيمياء الزراعية - كلية الزراعة - جامعة بني سويف

<sup>4</sup> قسم البساتين - كلية الزراعة - جامعة بني سويف

أجريت تجربة حقلية خلال الموسمين المتعاقبين 2019/2018 و 2020/2019 بالمزرعة البحثية بمحطة بحوث بساتين سدس بمحافظة بني سويف، بهدف تحسين النمو والانتاجية وكذلك محصول الزيت من خلال الرش الورقي بفيتامين هـ والأحماض الأمينية ومستخلص أوراق المورينجا. أوضحت النتائج أن أفضل صفات النمو الخضري (طول النبات وقطر الساق وعدد الأفرع الرئيسية ووزن النبات عند الحصاد)، والصفات المحصولية (عدد الكيسولات للنبات ومحصول البذور للنبات والقدان)، انتاجية الزيت (النسبة المئوية للزيت العطري والثابت ومحصول الزيت العطري والثابت للقدان)، وكذلك الخسائر الكيميائية مثل صبغات التمثيل الضوئي والنسبة المئوية للنتروجين والفوسفور والبوتاسيوم والأحماض الدهنية الحرة والرقم البودي ورقم التصبن للزيت الثابت وذلك نتيجة الرش الورقي بالتركيزات الأعلى من مستخلص أوراق المورينجا (5 مل/لتر) يليها الأحماض الأمينية (2 مل/لتر) ثم فيتامين هـ بمعدل 100 جزء في المليون، مع عدم وجود فروق معنوية في أغلب الصفات بين هذه المعاملات الثلاثة. لذلك يمكن التوصية برش نباتات حبة البركة بالتركيز الأعلى من مستخلص أوراق المورينجا بمعدل 5 مل/لتر لتحسين صفات النمو وإنتاجية القدان من البذور ومحصول الزيت الطيار والثابت تحت نفس الظروف البيئية المماثلة.

**الكلمات المفتاحية:** حبة البركة، مستخلص أوراق المورينجا، فيتامين هـ، الأحماض الأمينية، الزيت العطري والزيت الثابت