PROBABLE GENES FOR YELLOW RUST RESISTANCE IN SOME EGYPTIAN WHEAT CULTIVARS

Youssef, I. A. M.; S. Sherif and S. Negm Wheat disease Res. Dept., Plant Pathology Res. Institute, A.R.C. Giza, Egypt

ABSTRACT

The present work gave evidence to the presence of 11 physiological races of *Puccinia striiformis* West., the causal agent of stripe rust of wheat in Egypt during 2005/2006. These races were identified as: 4E0, 4E2, 4E16, 64E4, 76E4, 134E18, 196E18, 198E150, 198E86, 206E30 and 230E186. These races were identified according to their virulence to both the world and European set of wheat differentials. They showed a wide range of virulence. The obtained results indicated the efficacy of Yr4⁺, YrSP, Yr1, Yr5 and Yr10.

It was recommended to incorporate such genes within the local wheat cultivars aiming to increase their level of resistance against stripe rust.

Matching between cultivars and lines indicated that the wheat cultivar Sakha 61 was postulated to have 9Yr's. Also, the obtained results indicated that matching amongst cultivars clarified the presence of common resistance genes against stripe rust. This may be the reason behind the considerable level of resistance to stripe rust *i.e.*, Yr6 and Yr2 and this would indicate the genetic approximation of such cultivars (Gemmeza 5, Gemmeza 7, Giza 168, Sakha 61 and Sakha 93). This report is considered a primitive indicator to the presence of resistance genes, it could be confirmed by studying the pedigree of each cultivar and/or making crosses to observe the segregation. Consequently, the present work is considered an effective tool in the breeding program of wheat against rust diseases in general.

INTRODUCTION

Wheat (*Triticum aestivum*, L.) stripe rust incited by (*Puccinia striiformis* West.) is an important disease worldwide (Stubbs, 1985 and 1988 and Roelfs *et al.*, 1992). Cool and high humidity are suitable factors to the wide distribution of the disease (Stubbs, 1988, Johnson, 1988 and Danial, 1994). Since the discovery of genetic resistance in stripe rust that reported by Biffen (1905), many papers were reported *i.e.*, Flor (1955) suggested the gene- for gene theory, the occurrence of pyramiding resistance (accumulating multiple genes) as a term was suggested by Watson (1970). In Egypt, a lot of papers were published in the same trend *i.e.*, physiological specialization and varietal resistance (Abdel-Hak *et al.*, 1972; El-Daoudi *et al.*, 1977; Abu El-Naga *et al.*, 1997, 1998, 1999 and 2001, and Youssef *et al.*, 2003; 2006a and 2006b).

The main objective of the present work is the identification of stripe rust physiologic races in some of the northern governorates in Egypt and determining the postulated and common stripe rust genes involved in some Egyptian wheat cultivars.

MATERIALS AND METHODS

The present work was carried out at Sakha Agricultural Research Station, Wheat Disease Research Section in 2005/2006 growing season.

Stripe rust samples were collected from the different susceptible wheat cultivars showing the symptoms of stripe rust disease on leaves and/or

spikes (Hendrix, 1967, Hendrix and Fuchus, 1970; Manners, 1971 and Shaner and Powelson, 1973).

The general survey included certain governorates *i.e.*, Dakahlia, Dumytta and Kafrel-Sheikh in areas grown to wheat cultivars *i.e.*, Sakha 8, Sakha 69, Giza 163 and Rust trap nursery. Samples were left to dry over envelops at room temperature overnight and preserved at a fridge in disseccator containing calcium chloride till usage

Purification: single pustules were picked from each sample and used in the inoculation of eight days old seedlings from the susceptible wheat cultivar *i.e.*, Morocco, Little club or *Triticum spelta saharensis* following the method adopted by (Stubbs, 1988), in which seedlings were rubbed gently between fingers using sterile distilled water plus adhesive material (Tween 20), then inoculated with spore powder from each samples, preserved in humid apparatus at 9-10°C overnight in darkness. Transferred to the benches in controlled cadges at 15-18°C for 8 hrs. at night, and 16 hrs. at 15-18°C at light density of 7500 lux. This diurnal system was applied for 8-10 days. The developed uredial sori were immediately picked and transferred to complete pots of susceptible wheat cvs. to be used.

Race identification: The differential varieties listed in Table (1) in addition to 9 local vars. were artificially inoculated with the purified samples using the technique of Stubbs (1988). Records of stripe rust were recognized as high infection type (HIT) and low infection type (LIT). the high infection for each race was expressed as the decenary value for each differential of both world and European sets.

Postulation of resistance genes within local wheat cvs.

After the record of stripe rust, data on cvs. virsus physiological races, a matching was performed between cvs. (unknown genes) and wheat stripe rust differentials (known genes) to reach to the gene presence (whether it probably present or may be present) the method suggested by Statler (1984) was followed in this regard. This suggested method omitted the races that characterized by complete virulence or avirulence to test cvs. or Yr's.

Likewise, the host lines and /or cvs. Having complete resistance or susceptibility were also omitted from matching. So, the reaction (LIT):(HIT) or (HIT):(LIT) were adequately selected for the matching. The four categories (+, -, -0, and 0) were belt on the following basis:

- 0 : absence of (LIT): (HIT) cultures this means that cultivars involved the same gene(s).
- : the absence of (HIT) : (LIT), indicating that the hosts (Yr's) or (cv.), does not contain the same gene(s).
- 0 : the absence of H : L or L : H cultures indicating that both hosts carry the same resistance gene(s) at least for the used culture.
- + : the occurrence of some H : L cultures and some L : H cultures indicating that hosts do not carry the same resistant gene(s).

Compared 9 cultivars with unknown resistance genes (host B) to 17 isgenic lines each carrying a single known gene for resistance to stripe rust (host A). Because host A carries only a single gene, infection type in categories "0" and "- 0" *i.e.*, no LIT : HIT cultures indicates that cultivars carries the gene present in the isogenic line. Category "- 0" indicated that it

carries only that gene. Category "-" and "+" (*i.e.*, some cultures in the LIT : HIT infection type) indicate that the tested cultivar does not possess the gene carried in the isogenic line. Category "-" means that the cultivar does not carry the gene for resistance in the comparison. While category "+" indicates that there is a different gene for resistance.

Table (1): Identification of 11 physiologic races of stripe rust (Puccini	а
striiformis West.) on 17 wheat differential cultivars and thei	r
specifications in Egypt (2005/2006).	

Differential cultivar	Triticum spella f.sp album SPYr5	Clement Yr9	Suwong2XOmmar Yrsu	Strubs Dikkopf YrSD	Moro Yr10	Vilmorin 23 Yr3V	Heinskolben Yr6	Lee Yr7	Chinese 166 Yr1	World Set	HeinsVII Yr2 ⁺	Spalding prolific YrSP	Carsten V Yrcv	Compaire Yr8	Nord Desperz Yr3N	Heins Peko Yr6*	Reichersberg 42Yr7	Hybride 46 Yr4*	European Set	Race
Abbreviation	SP	CL	SU	SD	Мо	V23	НK	Lee	Ch		VII	SP	CV	Com	ND	Pe	R42	H46		
Yr's	5+	9+	SU	SD	10	3V	6	7	1		2+	SP	CV	8	3N	6+	7+	4+		
Decenary	(2)8	(2)7	(2)6	(2) ⁵	(2)4	(2) ³	(2) ²	(2) ¹	(2) ⁰		(2)7	(2)6	(2) ⁵	(2)4	(2) ³	(2) ²	(2) ¹	(2) ⁰		
value	256	128	64	32	16	8	4	2	1		128	64	32	16	8	4	2	1		
							н			4									0	4E0
							н			4							н		2	4E2
			н				п			4 64				п		н			10	4E10 64E4
			н			н	н			76						н			4	76E4
		н					Н	н		134				Н			н		18	134E18
		н	н				н			196				н			н		18	196E18
		Н	Н				Н	Н		198	Н			Н		Н	Н		150	198E150
1		Н	Н				Н	Н		198	Н		Н	Н	Н		Н		186	198E186
		н	н			н	н	Н		206				Н	н	н	Н		30	206E30
1	<u> </u>	н	н	н	I	L	н	н		230	н		н	н	н	L	н	ļ	186	230E186
L-Uigh inf	l Anti-		<u> </u>					- TI		230	n la		Н	н	п		п		186	230E186

H=High infection type L = Low infection type (Plank)

Postulation of common resistance gene(s) between cvs.

A theoretical matching was performed between cvs. and themselves after the confirmation of postulated genes aiming to limit the similar response for each cv. the following limitations must be taken into considerations: cultivars showing (0 or - 0), (- 0) indicates that both cultivars have the same gene(s) for resistance, while, 0 indicates that cultivar B carries at least one gene present in cultivar A. On the other hand (–) indicates that cultivar A carries at least one gene not present in cultivar B, and category (+) indicates that both cultivars carry genes which are not present in the other.

		Host B (unknown)
		Resistant	Susceptible
Heat A (known)	Resistant	LIT : LIT	LIT : HIT
HUSLA (KIIOWII)	Susceptible	HIT : LIT	HIT : HIT

The matching or comparison was performed in diallelic system and its reciprocal one aiming to search for at least one common gene for resistance at least with the tested isolates or races.

Gene efficacy: was recorded by dividing the resistant response by the total number of responses.

Virulence percentage was computed by dividing the susceptible responses by the total number of responses.

EXPERIMENTAL RESULTS

As regard to the identification of stripe rust (P. striiformis) predominant in Egypt during 2005/2006, data presented in Table (1) revealed such characterization. These data revealed that 12 physiologic race could be identified on 17 universal differential varieties (included 9 European and 8 world sets). These identified races were; 4E0, 4E2, 4E16, 64E4, 76E4, 134 E18, 196E18, 198E150, 198E185, 206E30, and 230E186, respectively. The value relevant to each race was derived from the summation of the dicenary values for each of defeated cultivars in each set (for each high infection type). For example race *i.e.*, 4E0 indicated that the race attacked Heins Kolben (Yr6) only and was avirulent to either of the lines of the European Set. On this foundation the identification was accomplished with the rest of races. It must be remembered that race 230E186 was repeated twice. It is clear from the commulated data that vars. i.e., Triticum spelta album (Yr5). Moro (Yr10), Chinese 166 (Yr1), Spalding prolific (YrSP) and Hybride 46 (Yr4⁺) were not attacked by either of the tested races. Consequently, these vars. were not represented in further matching to postulate gene in the commercials.

Data in Table (2) clarified the matching between 9 local commercial cvs., 16 stripe rust monogenic lines, when inoculated against 11 races of *P. striiformis* was singly used.

These data indicated that cv. Gemmeiza-5 may involve (Yr6 and Yr2), Gemmeiza-7 (Yr6 and Yr2), Gemmeiza-10 (Yr7 and Yr6), Giza 168 (Yr6, SU and 2), Sakha 8 (6 and 2), Sakha 61 (Yr1, 3N, 8, 7, 6, SU, 9, 2 and A⁺), Sakha 93 (8, 6 and 2). It was indicated that no tested Yr's could be detected with Gemmeiza-9 and Sakha-94.

Table (2): Revealed the incidence of cultivars in the LIT : HIT (Low infection type: High infection type) comparison of 16 near-isogenic lines of yellowrust (Yr's) and 9 commercial wheat cultivars inoculated with 11 racesof Puccinia striiformis West. f.sp. tritici at seedling stage undergreenhouse condition in 2005/2006 growing season.

Yr's Cultivars	Yr7+	Yr6⁺	Yr3N	Yr8	Yrcv	Yr2+	Yr7	Yr6	Yr3V	YrSD	YrSU	Yr9+	Yr18	Yr2	Yr19	Yr(A ⁺)
Gemmeiza5	+	+	-	-	•	•	1	0	•	1	+	-	•	0	+	I
Gemmeiza 7	+	+	-	-	I	-	I	0	I	i	+	-	I	0	+	-
Gemmeiza 9	i	-	-	-	I	-	I	+	I	i	I	-	I	+	I	-
Gemmeiza 10	0	+	-	+	I	+	+	0	+	+	+	+	+	+	+	+
Giza 168	+	+	+	+	+	+	+	0	I	+	0	+	+	0	+	+
Sakha 8	+	+	-	-	I	-	I	0	I	i	+	-	I	0	+	-
Sakha 61	0	+	0	0	+	+	0	0	+	+	0	0	+	0	+	0
Sakha 93	+	+	+	0	+	+	+	0	+	+	+	+	+	0	+	+
Sakha 94	+	+	-	-	-	-	-	+	+	-	+	-	+	+	+	-

Data in Table (2a) supported those recorded in table (2) since they indicated that Sakha 61 included *ca.*, 9 Yr's. Both Giza 168 and Sakha 93 included 3 Yr's, *i.e.*, Gemmeiza-5, Gemmeiza-7, Gemmeiza-10 and Sakha 8 included 2 Yr's each conversely, no tested Yr's could be detected in either of Gemmeiza-9 or Sakha 94.

Cultivars	Probability Yr's [*] genes	Efficacy of gene (%)
Gemmeiza5	Yr6 and Yr2	12.5
Gemmeiza 7	Yr6 and Yr2	12.5
Gemmeiza 9	0	0
Gemmeiza 10	Yr7 ⁺ and Yr6	12.5
Giza 168	Yr6, YrSU and Yr2	18.75
Sakha 8	Yr6 and Yr2	12.5
Sakha 61	Yr7+, Yr3N, Yr8, Yr7, Yr6, YrSU, Yr9+, Yr2 and YrA+	56.25
Sakha 93	Yr8, Yr6 and Yr2	18.75
Sakha 94	0	0

Table (2a): Probable resistance genes for yellow rust in some Egyptian commercial wheat cultivars in 2005/2006.

* yellow rust resistance genes

Data in Table (2b) showed the efficacy of Yr genes against the tested stripe rust races during 2005/2006. These data indicated that Yr6 exhibited the highest rate of efficacy (77.7%) followed by Yr2 (66.6%), Yr's (7⁺. 8 and SU) were 22.2% each; Yr's (3N, 7, 9⁺ and A⁺) represented by 11.1% each. finally no efficacy was detected with Yr's 6⁺, cv, 2⁺, 3V, SD, 18 and Yr14. *ca.* 0.0%.

Table (2b): The frequency of identified Yr's genes postulated in 9 commercial wheat cultivars in 2005/2006.

Yr's [*] resistance genes	No. of cultivars carrying Yr's	Efficacy (%)
Yr7+	2	22.2
Yr6⁺	0	0.00
Yr3N	1	11.1
Yr8	2	22.1
Yrcv	0	0.00
Yr2+	0	0.00
Yr7	1	11.1
Yr6	7	77.7
Yr3V	0	0.00
YrSD	0	0.00
YrSU	2	22.2
Yr9⁺	1	11.1
Yr18	0	0.00
Yr2	6	66.6
Yr19	0	0.00
Yr(A+)	1	11.1

* yellow rust resistance genes

Data in Table (3) revealed the interrelationships amongst the commercials and its reciprocal reaction. These data indicated that a common stripe rust gene(s) may be presented between each of Gemmeiza-5 and Sakha 8 and between Sakha 8 and Gemmeiza-5 and Gemmeiza-7.

On the other hand, common genes probably present between Sakha 61 and each of Gemmeiza -5, -7, -9, -10, Giza 168, Sakha 8 and -94, however, Sakha 93 likely has common gene with Gemmeiza -5, -7, -9 Sakha

8 and Sakha 94, also, Giza 168 was postulated to have common gene with Gemmeiza 5, -7 -9 and Sakha 8. Likewise, Gemmeiza -9 probably has common genes with Gemmeiza -5 and -7. Finally, Sakha 8 likely has common gene with Giza 168.

DISCUSSION

Wheat stripe rust incited by *Puccinia striiformis* West., is considered to be one of the distructive diseases of wheat (*Triticum aestivum*, L.) all over the world due to the introduction of new cultivars and/or lines with new genetic make up (Stubbs, 1980 and 1985 and Zadoks, 1961).

It was reported that problem of stripe rust of wheat due to relatively olden period (Hendrix, 1967; Manners, 1971; Abdel-Hak *et al.*, 1972; El-Daoudi, 1977 and Stubbs, 1980 and 1985), since the disease was known as a sporadic one, but its occurrence began to be continuous during 1990's (Abu- El-Naga *et al.*, 2001).

The present work is considered an extension to the trend of stripe rust researchers established in the latter decades especially after the building of stripe rust greenhouse at Sakha, Kafr El-Sheikh governorate.

Cultivars	Gemmeiza5	Gemmeiza 7	Gemmeiza 9	Gemmeiza 10	Giza 168	Sakha 8	Sakha 61	Sakha 93	Sakha 94
Gemmeiza5		- 0	-	+	0	- 0	0	0	+
Gemmeiza 7	- 0		-	+	0	- 0	0	0	+
Gemmeiza 9	0	0		0	0	0	0	0	0
Gemmeiza 10	+	+	-		+	+	0	+	+
Giza 168	-	-	-	+		-	0	+	+
Sakha 8	- 0	- 0	-	+	0		0	0	+
Sakha 61	-	-	-	-	-	-		+	-
Sakha 93	-	-	-	+	+	-	+		-
Sakha 94	+	+	-	+	+	+	0	0	

Table (3): Incidence of cultivars in the LIT : HIT (Low infection type : High infection type) comparison of cultivars inoculated with 11 physiologic races of *Puccinia striiformis* West. f.sp. *tritici*.

The obtained results indicated the presence of 11 physiologic races of *Puccinia striiformis* West. During the elapsed seasons, 2000-2005. These races were identified *i.e.*, 4E0, 4E2, 4E16, 64E4, 76E4, 134 E18, 196E18, 198E150, 198E186, 206E30 and 230E186. Similar results were recorded by Abu- El-Naga *et al.* (2001), Youssef *et al.* (2003, 2006a and 2006b) who mentioned that the races 230E150, 4E0, 4E2 4E16, 198E186 and 198E150 in Egypt. On the other hand, in neighboring countries races; 134E150 230E150, 230E134 and 230E150 were recorded by Hakim *et al.* (2001), added to races 134E22, 2E16, 134E146, 134E150 and 6E16, 134E146 and 134E150 that also, recorded by Louwers *et al.* (1992). It is concluded that

specification of such races were similar in time and space since the simultaneous occurrence may by repeated each season due to the advance in both wheat breading and disease programs as previously mentioned.

The obtained results also, indicated that Yr6 is considered a common gene amongst the tested cultivars followed by Yr2. So, the good performance of our cultivars may be due to the presence of such genes.

Conversely, the occurrence of Yr's *i.e.*, 6⁺, cv, 2⁺, 3V, SD, 18 and 19 indicated inefficacy that may be noticed with only our cultivars, since cv having Yr18 are effective elsewhere, where Yr18 varied from 5R to 40MR in Kazakhstan to 100s in Tajikistan according to Absattarova *et al.* (2001).

As regard to comparison between cultivars and the Yr's (monogenic lines) followed by comparisons between cultivars and cultivars. The obtained results indicated that most of these cultivars exhibited common genes (Yr's). These findings are considered speculation or postulation and/or tentative generalization, and to confirm these results we must see their pedigree or make crosses to confirm the presence of such gene within cultivar depending on their segregations.

Finally, the present work would be an effective tool if it is exploided well in the program of breeding for disease resistance.

REFERENCES

- Abdel-Hak, T. M.; D. M. S. Stewart and A. H. Kamel (1972). The current stripe rust situation in the near east countries regional wheat workshop. Beirut. Lebanon, 1-29.
- Absattarova, A.; S. Babayev; K. Bulatova; M. Karabayav; M. Koishibayev; A. Kokhmetova; V. Kuklachava; A. Morgunov; S. Rsaliv; A. Sarbayev; R. A. Urazaliev; M. Yessimbekova and C. R. Wellings (2001). Improvement of wheat yellow rust resistance in KazaKhstan and UZbakistan through sub-regional Co-operation meeting the challenge of yellow rust in cereal crops. Proceedings of the first Regional Conference on Yellow Rust in the center and West Asia and North Africa Region, 8-14 May 2001, Karaj, Iran.
- Abu el-Naga, S. A.; M. M. Kalifa: S. Sherif; W. A. Youssef; Y. H. El-Daoudi and I. Shafik (2001). Virulence of wheat stripe rust pathotypes identified in Egypt during 1991/2000 and sources of resistance. First Regional Yellow Rust Conference for Central & West Asia and North Africa 8-14 May, 2001, Spll, Karaj, Iran.
- Abu el-Naga, S. A.; M. M. Kalifa; A. A. Bassioni, W. A. Youssef; T. M. Shehab El-Din and A. H. Abd El-Latif (1999). Revised evaluation for Egyptian wheat germplasm against physiologic pathotypes of stripe rust (*Puccinia striiformis* west). J. Agric. Sci. Mansoura Univ.,24(2):477-488.
- Abu el-Naga, S. A.; M. M. Kalifa; W. A. Youssef and A. H. Abd El-Latif (1998). Stripe rust situation during the period (1994-1995) with special reference to designating genes conferring resistance in certain Egyptian wheat germplasm. J. Agric. Sci., Mansoura Univ., 23(3): 1127-1136.

- Abu El-Naga, S. A.; M. M. Kalifa; W. A. Youssef; I. A. Imbaby; M. M. El-Shamy; E. Amer; T. M. Shehab El-Din and M. M. E. Shamy (1997). Effect of stripe rust infection on grain yield in certain wheat cultivars and the economic threshold of chemical contral application in Egypt during 1995/97 growing seasons. National Annual condition metting NVRSRP/Egypt Sp. 11-15, 1997: 81-90.
- Biffen, R. H. (1905). Mendel's low's of inheritance and wheat breeding. J. Agric. Sci.,1: 4-48.
- Danial, D. L. (1994). Aspect of durable resistance in wheat to yellow rust. Ph.D. Thesis Wageningen Agricultural University the Netherland.
- El-Daoudi, Y. H(1977). Studies on stripe rust of wheat in Egypt. Master of science Thesis. Faculty of Agriculture Cairo Univ., 78pp.
- Flor, H. H. (1955). Host-parasite interaction in flax rust- its genetic and after implications. Phytopathology, 45: 680-685.
- Hakim, M. S.; A. Yahyaoui, M. El-Naimi and I. Maaz (2001). Wheat yellow rust pathotypes in Western Asia. Proceedings of the first Regional conference on yellow rust in the central and West Asia Africa Region, 8-14 May 2001, Karaj, Iran.
- Hendrix. J. W (1967). Stripe rust of wheat. Pages 325-332 in K. S. quizenbary and L. P. Reitz, eds. Wheat and wheat Improvement Agronomy Monograph No. 13, Am. Soc. Agron. Madison, WI.
- Hendrix. J. W and E. Fuchs (1970). Influence of fall stripe rust infection on tillering and yield of wheat. Plant Dis. Rep. 54: 347-349.
- Johnson, R. (1988). Durable resistance to yellow (stripe) rust in wheat and its implication in plant breeding. In: N. W. Simmonds and S. Rajaram (eds.) Breeding strategies for Resistance to the rust of wheat, p.63-75 CIMMYT, Mexico.
- Louwers, J. M.; C. H. Van Silfhout and R. W. Stubbs (1992). Race analysis of yellow rust in wheat in developing countries. Report. 1990-1992 IPODLO Report 92-11p. DLO Research Institute for Plant Protection (IPO-DLO) Wageningen. The Netherlands.
- Manners, J. G., (1971). Yellow rust of wheat and barley, pages 232-237 in J. H. Western, ed. Diseases of crop plants. Macmillan Press Itd., London.
- Roelfs, A. P.; R. P. Singh and E. E. Saari (1992). Rust diseases of wheat: concepts and Methods of disease Management, Mexico, D. F. CIMMYT 81pp.
- Shaner, G and R. L. Powelson (1973). The oversummering and dispersal of inoclum of *Puccinia striiformis* in oregon. Phytopath., 63: 13-17.
- Statler, G. D.(1984). Probable genes for leaf rust resistance in several hard red spring wheat. Crop Sci., 24: 883-886.
- Stubbs, R. W (1980). Environmental resistance of wheat yellow rust (*Puccinia striiformis* westend, f.sp. *tritici*). Proceeding of the Fifth European Mediterranean Cereal Rusts Conference. 77-81.
- Stubbs, R. W (1985). The cereal rust (11) (A. P. Roelfs and W. R. Bushnell, edds.) Orlando, Florida Academic Press. 61-101.

- Stubbs, R. W (1988). Pathogenicity analysis of yellow (stripe) rust of wheat and its significance in global context. In: Simonds, N. W.& Rajaram, S. (Eds.) Breeding Strategies for resistance to rust of wheat CIMMYT, Mexico, p. 23-38.
- Watson, I. H. (1970). Changes in virulence and population shifts in plant pathogens. Ann. Rev. Phytopathology, 8: 209-230.
- Youssef, I. A. M., S. S. Negm and Gamalat, A. Hermas (2006a). Physiologic specialization in *Puccinia striiformis* and genes conditioning resistance to yellow rust disease in certain wheat cultivars in Egypt. J. Agric. Sci: Mansoura Univ., 31(6): 3473-3482.
- Youssef, I. A. M.; A. A. M. Abualy; I. A. El- Salamony and Doaa, R. El-Naggar (2006b). Identification of physiologic races of stripe (yellow) rust and postulation of resistance genes in certain Egyptian wheat cultivars. Egypt. J. of Appl. Sci., 21(10B): 404-418.
- Youssef, W. A.; M. A. Najeed; Mateld Fransis and Eetmad, E. Draz (2003). Wheat stripe rust pathotypes their Frequency and Virulence Formulae in Egypt during 2000/2001. J. Agric. Sci. Mansoura Univ., 28(5): 3469-3477.
- Zadoks. J. C. (1961). Yellow rust of wheat: studies in epidemiology and physiologic specialization. Tijdschrift over Planteziekten 67: 69-256.

جينات المقاومة المتوقعة للصدأ الأصفر في بعض أصناف القمح المصرية عصام عبد الحميد محمد يوسف – صلاح الدين شريف عمر – صبحي نجم قسم بحوث أمراض القمح – معهد بحوث أمراض النباتات – مركز البحوث الزراعية – الجيزة – مصر.

دلت نتائج هذا البحث علي وجود احدي عشر سلالة فسيولوجية من فطر الصدأ الأصفر علي أصناف القمح المصرية خلال موسم ٢٠٠٦/٢٠٠ ويتسبب هذا المرض عن فطر بكسينيا 4E0, 4E2, 4E16, ويتسبب هذا المرض عن فطر بكسينيا 64E4, 76E4, 134E18, 196E18, 198E150, 198E86, 206E30 and 230E186

وقد تم تعريف هذه السلالات من منطلق عدوانيتها على كل من مفرقات المجموعة العالمية والأوربية. وقد تراوحت هذه السلالات في مدي عدوانيتها على هذه الأصناف المفرقة. وقد دلت النتائج علي كفاءة الجينات Yr4+, YrSP, Yr1, Yr5 and Yr10.

وقد أوصيناً في هذا البحث بضرورُة إدخال هذه الجينات في الأصناف المنزرعة بغية تحسينها من حيث المقاومة لهذا المرض والتصدي لحالات الوبائية التي قد تظهر فجأة. ومن ناحية أخري دلت النتائج بأن الصنف سخا ٦١ من المحتمل أن يحتوي علي ٩ جينات تقريبا. كما أن هناك جينات مشتركة بين أغلب الأصناف موضع الاختبار وهي Yr6 and Yr2 وهذه الجينات يعزي إليها المستوي العام للمقاومة للأصناف المختبرة ، ويدل أيضا على التقارب الوراثي بينها (جميزة °، جميزة ٢، جيزة ١٦٨، سخا ٢٦، سخا ٩٣.)

جُميزة ٧، جَيزَة ١٦٨، سخا ٦١، سخا ٩٣.) وهذا البحث يعتبر مؤشر مبدئي على نوعية جينات المقاومة في الأصناف السائدة ويتم تأكيد نتائج هذا البحث بدراسة نسب هذه الأصناف وكذلك بعمل تهجينات بين الأصناف ودراسة الانعز الات الناتجة عنها. ومن هنا نقول: أن هذا البحث يعتبر أداه فعالة في برامج التربية للمقاومة للأصداء في مصر.