

Journal of Plant Production

Journal homepage & Available online at: www.jpp.journals.ekb.eg

Response of *Schefflera actinophylla* Plants Grown under Drought to Soil Compost Addition and Foliar Application of Melatonin

Abd-Elhady, W. M. F.¹; A. G. Badour² and M. A. El-Sherpiny^{2*}



¹ Ornamental Plants and Landscape Gardening Research Department, Horticultural Research Institute, Agriculture Research Center, Giza, Egypt

² Soil Fertility & Plant Nutrition Research Department, Soil, Water and Environment Research Institute, Agriculture Research Center, El-Gama St., Giza, 12619 Egypt

ABSTRACT

The provision of irrigation water has become crucial, especially for ornamental plants, due to Egypt's demanding water conditions. The use of inexpensive soil amendments *i.e.*, compost and the application of a newly discovered molecule "melatonin" can mitigate the negative effects of water scarcity. Therefore, this study aimed to evaluate the efficacy of combining compost with melatonin in alleviating the adverse effects of water deficit on *Schefflera actinophylla* L plants. Different irrigation regimes *i.e.*, irrigation every 5 (I₁), 7 (I₂) and 10 (I₃) days were evaluated as the main factor. The final composition of the pots, consisting of compost and sand, with options including 1:3 compost: sand mixtures v/v (C₁), 1:1 compost: sand mixtures v/v (C₂) and 3:1 compost: sand mixtures v/v (C₃) was investigated as a secondary factor. In addition, the application of melatonin by spraying at concentrations of zero (F₁) and 100 μM (F₂) was investigated as a tertiary factor. Plant performance decreased with increasing irrigation intervals. Furthermore, plants grown in pots containing 75%compost+25% sand (C₃) showed the best performance followed by the C₂ treatment and finally the C₁ treatment. Moreover, the application of melatonin led to an increase in all the growth and chemical characteristics studied, as well as in the levels of antioxidant, except for malondialdehyde, which decreased compared to the untreated plants. On another note, the plant performance was better under combined treatments of I₂x C₂ (or C₃) x F₂ than that under combined treatments of I₁x C₁x F₁. Finally, these results confirm the potential role of compost and melatonin in mitigating the adverse effects of water deficit.

Keywords: Water deficit, compost, melatonin



INTRODUCTION

The *Schefflera actinophylla* plant belongs to the Araliaceae family. Native to Australia, it has become a popular houseplant due to its distinctive umbrella-like foliage and adaptability to indoor conditions (Abu-Khalaf and Natsheh (2022)). *Schefflera actinophylla* is favored for its aesthetic appeal, making it a popular choice for interior decoration and office spaces. Its glossy green leaves and unique growth habit have earned it a special place among plant lovers, contributing to its popularity as a decorative addition to homes and workplaces (Ahmed and Shahin (2023)).

Saving irrigation water is critical for a number of reasons, including resource conservation, environmental protection, economic efficiency, food security, climate resilience, and overall sustainability. It's a shared responsibility involving farmers, policy makers, and society at large to ensure the efficient and responsible use of this precious resource in agriculture (Saad Eddin *et al.* 2023). However, it's important to note that while water conservation is essential, it must be balanced with the needs of crop production. In some cases, extreme water conservation can lead to water deficit conditions. These water deficit conditions can trigger the generation of reactive oxygen species (ROS) within plant cells, which, in turn, can cause oxidative damage to cell structures and disrupt normal

cellular processes (Elsherpiny 2023). This highlights the importance of adopting water-saving practices that are both efficient and sensitive to the specific water needs of crops. Sustainable irrigation practices aim to optimize water use, by ensuring that crops receive adequate moisture without excessive water waste, thus promoting both water conservation and healthy plant growth (Moursy *et al.* 2023).

Melatonin is known for its potent antioxidant properties, effectively scavenging ROS. By reducing oxidative stress, melatonin acts as a protective shield, protecting plant cells from damage caused by drought-induced ROS (Arnao and Hernández-Ruiz 2015). The beneficial effects of melatonin can be attributed to its ability to regulate stomatal closure (Ahmad *et al.* 2023), enabling plants to balance water conservation with the maintenance of photosynthesis (Dradrach *et al.* 2022). Additionally, melatonin increases photosynthetic rates during drought conditions by protecting the photosynthetic apparatus within chloroplasts from drought-induced damage (Ahmad *et al.* 2023). Moreover, melatonin is thought to stimulate root growth and development, a key factor in facilitating plant water uptake (Khattak *et al.* 2023). A well-developed root system enables the plant to explore a larger volume of soil, thereby increasing, and its resilience to drought (Khosravi *et al.* 2023). Melatonin plays a key role in modulating the expression of genes and proteins involved in stress response pathways (Langaroudi *et al.* 2023). This includes the

* Corresponding author.

E-mail address: M_elsherpiny2010@yahoo.com

DOI: 10.21608/jpp.2023.249013.1288

activation of genes responsible for the synthesis of protective molecules and proteins that enhance the plant's ability to cope with water deficit (Wang *et al.* 2023). Furthermore, melatonin interacts with several plant hormones, including abscisic acid (ABA), which plays a central role in the plant's response to drought stress (Yang *et al.* 2023).

On the other hand, compost enhances water retention under drought stress by improving soil structure (Shah *et al.* (2023), increasing organic matter content, reducing evaporation, promoting root growth and supporting beneficial microbial activity (Elshepiny 2023). These combined effects make compost a valuable tool for improving the soil's ability to retain water and provide it to plants during periods of water scarcity (Paymaneh *et al.* 2023). Moreover, compost slowly releases nutrients over time as it decomposes. These nutrients include nitrogen (N), phosphorus (P), potassium (K), as well as various micronutrients (Soussani *et al.* 2023).

Therefore, the primary goals of the current research are to provide a thorough understanding of the effects of compost and melatonin on improving the performance and increasing the drought tolerance of the *Schefflera actinophylla* plant. This knowledge is important for improving ornamental plant production and developing sustainable approaches to water conservation, especially in water-scarce regions such as Egypt.

MATERIALS AND METHODS

A pot study was conducted to assess how different irrigation regimes [*i.e.*, irrigation every 5 days (I₁), 7 days (I₂) and 10 days (I₃), as the main factor], different compost ratios [*i.e.*, 1:3 compost: sand mixtures v/v (C₁), 1:1 compost: sand mixtures v/v (C₂) and 3:1 compost: sand mixtures v/v (C₃), as the secondary factor] and melatonin application [*i.e.*, zero (F₁) and 100 µM (F₂), as the tertiary factor] affect the growth performance and chemical content of *Schefflera actinophylla* cv. Compacts plants. These treatments were organized in a split-split-plot design, as each treatment was replicated five times.

1. Experimental site

The location of this pot experiment was Al-Mansoura Agricultural Research Station, Agricultural Research Center., Egypt (31.0500°N latitude and 31.3833°E longitude). The study lasted from February 28th to October 2nd, covering both the 2022 and 2023 seasons, each of which lasted seven months.

2. Plant material

The *Schefflera's* leaves are typically large and compound compared to the leaves of some other indoor plants. The leaves are usually compound, consisting of several separate, twisted segments that resemble fingers. The segments can be semi-circular to oval in shape. Leaves are typically a vibrant green color, although the shade of green may vary slightly depending on soil conditions and light levels. Leaves grow progressively from the main stem of the plant, with the largest and most mature leaves at the top. The leaf surface is generally smooth and glossy, as described by Abu-Khalaf and Natsheh (2022); Ahmed and Shahin (2023).

3. Studied substances

Plant cultivar: The *Schefflera actinophylla* cv. Compacts used in this study was obtained from the private nursery located in El Qanater El Khayreya, Egypt. Similar plants

with a height of 20 cm were selected and their number of compound leaves ranged from 9 to 11.

Melatonin: It was purchased from the Sigma Company (Sigma-Aldrich, St. Louis, MO, USA). A foliar spray of melatonin was prepared by dissolving melatonin in ethanol at a concentration of 10 mM and storing it at -20°C. Prior to application to the plants, this mixture was further diluted to attain a concentration of 100 µM, as described by Dradrach *et al.* (2022).

The medium used to fill the pots: Compost (field crop residues) was obtained from the Nile Compost Company, Egypt. Its characteristics are shown in Table 1. While the characteristics of the sandy soil used in this research are shown in Table 2. The analysis was carried out according to the methodology described by Dewis and Freitas (1970).

Table 1. Some characteristics of the compost used

Property	Values
EC, dSm ⁻¹ (1:10)	2.56
pH (1:10)	7.34
Phosphorus (P ₂ O ₅), %	0.42
Potassium (K ₂ O), %	0.76
Organic matter, %	50.59
Organic carbon, %	29.34
Total nitrogen, %	1.38
C:N ratio	21:1

Table 2. Some characteristics of the sand soil used

Traits and units	Values	
Particles size distribution, g kg ⁻¹	Sand	900
	Silt	45.0
	Clay	55.0
Texture class	Sandy	
O.M, %	0.13	
EC, dSm ⁻¹	0.70	
pH	8.00	

4. Experimental set up

First, the potting medium was prepared, and then the plastic pots were filled according to the specified treatment conditions, each pot being 25 cm in diameter and 18 cm deep. After the plants were obtained from the nursery, they were transplanted into their designated plastic pots (in the last week of February). The plants were placed in a seran house, which allowed 65% shading of light. A fertilizer solution was prepared by dissolving 2.0 grams of a commercial compound known as Kristalon (19-19-19) in one liter of tap water. The NPK fertilizer was applied as a soil drench at the same rate every month until the end of the experiment. The first dose was applied 10 days after transplanting, and subsequent doses were applied at monthly intervals. The foliar application of melatonin was made 8 times, with a 15-day interval between each application until the last week of August, using a plastic atomizer, as the first application of melatonin was made in the first week of May. The application rate was 1.5 liters per pot for the melatonin treatments. Irrigation was carried out as usual for *Schefflera actinophylla* L plants until the first week of May, after which the irrigation process was carried out according to the prescribed irrigation treatments.

5. Measurements

The physical characteristics of a whole plant, leaf chemical constituents, plant pigments, antioxidants and oxidation indicators were determined on 2nd October of each season, as shown in Table 3

Table 3. Parameters, methods and references of measurements

Parameters	Methods	References
Physical characteristics of a whole plant		
1. Plant height (cm)	Manually and visually	-----
2. No. of leaves plant ⁻¹		
3. Leaf area (cm ² plant ⁻¹)		
4. Leaf fresh and dry weights (g plant ⁻¹)		
5. Stem fresh and dry weights (g plant ⁻¹)		
6. Root fresh and dry weights (g plant ⁻¹)		
Plant pigments		
1. Chlorophyll a (mg g ⁻¹ F.W)	Spectrophotometrically	Schoefs (2005)
2. Chlorophyll b (mg g ⁻¹ F.W)		
3. Carotene (mg g ⁻¹ F.W)		
Digestion of plant samples		
Digestion for N,P,K determenations	Using a 1:1 mixture of H ₂ SO ₄ and HClO ₄	Peterburgski (1968)
Nutrient elements		
Nitrogen (%)	Via Kjeldahl method	Harborne (1984)
Phosphorus (%)	Using spectrophotometric apparatus	
Potassium (%)	Using flame photometer apparatus	
Biochemical traits		
Total carbohydrate,%	Using spectrophotometric apparatus	Herbert <i>et al.</i> (1971).
Indicators of oxidation		
Malondialdehyde (MDA, μmol.g ⁻¹ F.W)	Spectrophotometric method	Davey <i>et al.</i> (2005)
Antioxidative enzymes		
1. Peroxidase (POX, unit mg ⁻¹ protein ⁻¹)	Spectrophotometric method	Güneş <i>et al.</i> (2019)
2. Catalase (CAT, unit mg ⁻¹ protein ⁻¹)		

6. Statistical analysis

Statistical analysis of the data was performed using CoStat (Version 6.303, CoHort, USA, 1986). The least significant difference (LSD) test was used to compare means, at a significance level of 0.05, following the approach described by Gomez and Gomez (1984). To further evaluate the means between treatments, the Duncan's test was used, as prescribed by Duncan (1955).

RESULTS AND DISCUSSION

1. Physical characteristics of an entire plant

The data presented in Tables 4 & 5 show the individual effect of different irrigation schedules, different compost ratios and melatonin application as well as their interaction, on various physical characteristics of *Schefflera actinophylla* L plants. These characteristics include plant height (cm), No. of leaves plant⁻¹, leaf area (cm² plant⁻¹) (Table 4), fresh and dry leaf weights (g plant⁻¹), fresh and dry stem weights (g plant⁻¹), fresh and dry root weights (g plant⁻¹) (Table 5).

The values of all the above mentioned characteristics decreased as the intervals between irrigations increased. In other words, the irrigation every 5 days (I1 treatment) was the superior for obtaining the maximum values followed by I2 treatment (irrigation every 7 days), while the lowest values were obtained when *Schefflera actinophylla* L plants were irrigated every 10 days (I3 treatment)

Regarding the effect of compost, the plants grown in pots containing 3:1 compost: sand mixtures v/v (C3) showed the highest values for all the above parameters followed by the C2 treatment and finally the C1 treatment.

Regarding melatonin, its application led to an increase in all the above parameters compared to untreated plants. On the other hand, plant performance was better in the combined treatments of I2xC2 (or C3) xF2 than in the combined treatments of I1xC1xF1. The same trend was observed in both seasons studied.

In general, these results confirm the potential role of compost and melatonin in mitigating the negative effects of water deficit. These results can be explained in the following way; Longer watering intervals, such as in the I3 treatment, can cause water stress in *Schefflera actinophylla* L plants.

Water stress reduces the plant's ability to absorb nutrients and perform photosynthesis, resulting in reduced growth and overall plant health. Frequent irrigation, as in the I1 treatment, ensures consistent soil moisture, which promotes healthy root development in *Schefflera actinophylla* L plants. This, in turn, allows for better nutrient and water uptake, resulting in increased plant growth. Frequent irrigation helps to maintain higher transpiration rates, which can have a positive effect on leaf area and overall plant development.

Compost-rich substrates, such as C3 treatment, typically have a higher nutrient content, providing essential elements for *Schefflera actinophylla* L plant growth. The availability of nutrients in the substrate can improve overall plant development, including leaf area, number of leaves, and biomass. Compost can improve soil structure and water retention, ensuring adequate moisture for plant roots. This leads to better nutrient uptake and overall plant performance. Compost can encourage beneficial microbial activity in the soil, helping to mineralize nutrients and make them available to the plant.

Melatonin is known for its antioxidant properties, which can help protect plant cells from oxidative stress caused by water deficit. Reduced oxidative stress can lead to healthier and larger leaves, as well as improved biomass. Melatonin can enhance a plant's stress response mechanisms, allowing it to better cope with water deficit conditions. This can result in increased growth parameters such as plant height and leaf number. Melatonin has been shown to increase photosynthesis and chlorophyll content in plants, which can contribute to increased leaf area and overall plant vitality (Arnao and Hernández-Ruiz 2015).

The combined treatments of optimal irrigation, nutrient-rich compost, and melatonin are likely to have synergistic effects. Adequate irrigation ensures the availability of nutrients from the compost and facilitates the uptake of melatonin, leading to improved plant performance.

These scientific explanations highlight the complex interplay between irrigation, substrate composition, and melatonin in influencing the physiological and biochemical processes that control plant growth and development under water deficit conditions.

These findings are consistent with those documented by Shah et al. (2023) and Elsherpiny (2023), who both confirmed that organic fertilizers, particularly compost, improve soil water retention and the water-holding capacity of the root zone. Additionally, the results are consistent with studies by Ahmad et al. (2023) and Dradrach et al. (2022), which confirmed that melatonin contributes to increased plant tolerance to drought. These combined studies provide further support for the beneficial effects of compost and melatonin in mitigating the effects of water deficit on plant growth and performance.

2. Chemical contents

The data also show that different irrigation schedules, different compost ratios, and the application of melatonin have a significant effect on the pigments of *Schefflera actinophylla* L plants, including chlorophyll a & b (mg g⁻¹ F.W), carotene (mg g⁻¹ F.W) (Table 6) as well as leaf chemical content of nitrogen, phosphorus, potassium, carbohydrate content (%) (Table 7) in the leaves. The data show that the irrigation every 5 days (I₁ treatment) was the best for obtaining the maximum values of all the above mentioned traits. It was followed by the I₂ treatment (irrigation every 7 days), and finally by the I₃ treatment (irrigation every 10 days). These results indicate that the frequency of irrigation plays a key role in shaping the pigment content and chemical composition of the leaves of *Schefflera actinophylla* L plants. Higher irrigation frequency, as exemplified by the I₁ treatment, seems to enhance the accumulation of pigments and the nutrient content within the leaves of *Schefflera actinophylla* L plants.

Regarding the compost treatments, the plants grown in pots containing 3:1 compost: sand mixtures v/v (C₃) showed the highest values for chlorophyll a & b (mg g⁻¹ F.W), carotene (mg g⁻¹ F.W.) (Table 6) as well as leaf chemical content of nitrogen, phosphorus, potassium, carbohydrate content (%) (Table 7). This was followed by the C₂ treatment and finally the C₁ treatment.

Concerning the compost treatments, the plants that were grown in pots filled with a mixture of compost and sand at ratio of 3:1 v/v (C₃ treatment) showed the highest values for chlorophyll a & b (mg g⁻¹ F.W), carotene (mg g⁻¹ F.W) (Table 6) as well as leaf chemical content of nitrogen, phosphorus, potassium, carbohydrates content (%) (Table 7). After the C₃ treatment in terms of these parameters, the C₂ treatment showed intermediate values, and the C₁ treatment gave the lowest values.

These results suggest that the organic-rich C₃ compost treatment provides optimal conditions for improving plant pigment content and nutrient composition. The higher organic content in this treatment is likely to contribute to improved nutrient availability, which in turn promotes chlorophyll synthesis and overall leaf chemistry, including nitrogen, phosphorus, potassium, and carbohydrates.

Regarding melatonin, its application resulted in an increase in the values of chlorophyll a & b (mg g⁻¹ F.W.), carotene (mg g⁻¹ F.W.) (Table 6) as well as leaf chemical content of nitrogen, phosphorus, potassium, carbohydrate content (%) (Table 7) compared to untreated plants.

In relation to melatonin, its application resulted in an increase in the values for chlorophyll a & b (mg g⁻¹ F.W), carotene (mg g⁻¹ F.W) (Table 6). Additionally, the application of melatonin led to an increase in the chemical content of the leaves in terms of nitrogen, phosphorus, potassium, and carbohydrate content (%) (Table 7), when compared to the plants that were not treated.

Table 6. Effect of irrigation schedules, varying compost ratios and melatonin on chlorophyll a & b (mg g⁻¹ F.W) and carotene (mg g⁻¹ F.W) contents in leaves of *Schefflera actinophylla* L plants during two consecutive seasons (2022-2023)

Treatments	Chlorophyll a		Chlorophyll b		Carotene			
	(mg g ⁻¹ F.W)							
	1 st	2 nd	1 st	2 nd	1 st	2 nd		
Main factor : Irrigation schedules								
I ₁	1.028a	1.080a	0.632a	0.658a	0.516a	0.532a		
I ₂	0.992b	1.042b	0.619b	0.643a	0.501b	0.517b		
I ₃	0.896c	0.940c	0.583c	0.607b	0.464c	0.479c		
LSD at 5%	0.011	0.009	0.012	0.015	0.004	0.014		
Sub main factor: Varying compost ratios								
C ₁	0.935c	0.983c	0.596c	0.620c	0.479c	0.493c		
C ₂	0.973b	1.021b	0.611b	0.636b	0.494b	0.509b		
C ₃	1.008a	1.059a	0.626a	0.652a	0.509a	0.525a		
LSD at 5%	0.010	0.010	0.007	0.010	0.005	0.010		
Sub-sub main factor: Melatonin applications								
F ₁	0.962b	1.010b	0.607b	0.631a	0.490b	0.505a		
F ₂	0.983a	1.032a	0.615a	0.641a	0.498a	0.514a		
LSD at 5%	0.006	0.008	0.002	N.S	0.002	N.S		
Interaction								
I ₁	C ₁	F ₁	0.980	1.034	0.612	0.637	0.498	0.513
		F ₂	0.994	1.043	0.618	0.644	0.503	0.519
	C ₂	F ₁	1.008	1.057	0.621	0.647	0.507	0.522
		F ₂	1.051	1.103	0.641	0.669	0.526	0.541
C ₃	F ₁	1.064	1.117	0.648	0.673	0.529	0.543	
	F ₂	1.073	1.129	0.651	0.681	0.534	0.550	
I ₂	C ₁	F ₁	0.937	0.981	0.603	0.628	0.482	0.496
		F ₂	0.956	1.010	0.606	0.630	0.486	0.500
	C ₂	F ₁	0.969	1.019	0.610	0.634	0.492	0.506
		F ₂	1.018	1.069	0.626	0.650	0.511	0.525
	C ₃	F ₁	1.031	1.079	0.632	0.657	0.517	0.533
		F ₂	1.043	1.096	0.637	0.662	0.521	0.539
I ₃	C ₁	F ₁	0.863	0.906	0.563	0.585	0.448	0.460
		F ₂	0.882	0.924	0.575	0.599	0.456	0.471
	C ₂	F ₁	0.892	0.933	0.582	0.607	0.463	0.478
		F ₂	0.898	0.944	0.588	0.612	0.468	0.482
	C ₃	F ₁	0.912	0.960	0.591	0.615	0.474	0.488
		F ₂	0.927	0.972	0.598	0.622	0.477	0.494
LSD at 5%	0.018	0.023	0.007	0.036	0.006	0.037		

Means within a row followed by a different letter (s) are statistically different at a 0.05 level

Since, I₁: Irrigation every 5 days; I₂: Irrigation every 7 days; I₃: Irrigation every 10; C₁: 1 compost: 3 sand (v/v); C₂: 1 compost: 1 sand (v/v); C₃: 3 compost: 1 sand (v/v); F₁: Without melatonin and F₂: Melatonin at rate of 100 µM

N.S*: Non-significant

This shows that melatonin plays a positive role in improving the pigment content and nutrient composition of the plant. The antioxidant properties of Melatonin are likely to contribute to the reduction of oxidative stress, which may lead to improved chlorophyll synthesis and pigment accumulation. Furthermore, melatonin may influence nutrient uptake and utilization, leading to higher levels of nitrogen, phosphorus, potassium, and carbohydrates in the leaves, ultimately promoting plant growth and stress tolerance. In summary, the effects of melatonin on plant drought tolerance are diverse. It acts as an antioxidant, regulates stomatal behavior, maintains photosynthesis, enhances root development, improves stress signalling, and interacts with plant hormones. Together, these mechanisms contribute to improved drought tolerance in melatonin-treated plants, making it a promising tool for mitigating the adverse effects of drought stress (Ahmad et al. 2023).

On the other hand, the plant performance in the combined treatments of I₂x C₂ (or C₃) x F₂ was better than that in the combined treatments of I₁x C₁x F₁. The same trend was observed in both seasons studied. The same pattern or trend was observed in both of the seasons studied. These results are consistent with those reported by Dradrach et al. (2022); Shah et al. (2023) and Elsherpiny (2023).

Table 7. Effect of irrigation schedules, varying compost ratios and melatonin on the leaf chemical content of nitrogen, phosphorus, potassium, and carbohydrate content (%) in leaves of *Schefflera actinophylla* L plants during two consecutive seasons (2022-2023)

Treatments	N		P		K		Carbohydrates			
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd		
Main factor : Irrigation schedules										
I ₁	3.36a	3.51a	0.525a	0.534a	2.17a	2.27a	27.25a	27.92c		
I ₂	3.22b	3.35b	0.510b	0.519b	1.97b	2.05b	26.38b	27.09b		
I ₃	2.88c	3.00c	0.470c	0.479c	1.76c	1.83c	24.06c	24.69a		
LSD at 5%	0.08	0.05	0.002	0.010	0.10	0.09	0.37	0.28		
Sub main factor: Varying compost ratios										
C ₁	3.02c	3.14c	0.486c	0.495c	1.88b	1.96b	25.03c	25.68c		
C ₂	3.16b	3.29b	0.501b	0.509b	1.91b	1.99b	25.94b	26.62b		
C ₃	3.29a	3.42a	0.518a	0.528a	2.11a	2.21a	26.73a	27.41a		
LSD at 5%	0.05	0.05	0.006	0.006	0.06	0.06	0.42	0.24		
Sub-sub main factor: Melatonin applications										
F ₁	3.12b	3.24b	0.498b	0.506b	1.90b	1.99b	25.66a	26.36b		
F ₂	3.19a	3.32a	0.506a	0.515a	2.03a	2.12a	26.14a	26.77a		
LSD at 5%	0.06	0.07	0.003	0.009	0.05	0.05	N.S	0.1		
Interaction										
I ₁	C ₁	F ₁	3.20	3.32	0.505	0.512	2.00	2.09	26.21	26.91
		F ₂	3.24	3.38	0.507	0.518	2.05	2.15	26.56	27.25
	C ₂	F ₁	3.29	3.44	0.514	0.521	2.09	2.19	26.71	27.46
		F ₂	3.46	3.60	0.536	0.544	2.27	2.36	27.71	28.40
	C ₃	F ₁	3.48	3.62	0.543	0.552	2.30	2.41	28.02	28.58
		F ₂	3.51	3.67	0.548	0.557	2.33	2.45	28.29	28.92
I ₂	C ₁	F ₁	3.03	3.16	0.490	0.499	1.92	2.01	25.02	25.77
		F ₂	3.05	3.17	0.495	0.503	1.96	2.04	25.33	25.92
	C ₂	F ₁	3.12	3.23	0.500	0.510	1.40	1.45	25.94	26.71
		F ₂	3.33	3.46	0.519	0.526	2.14	2.22	27.06	27.64
	C ₃	F ₁	3.37	3.49	0.526	0.534	2.18	2.29	27.35	28.19
		F ₂	3.42	3.56	0.530	0.540	2.20	2.29	27.57	28.28
I ₃	C ₁	F ₁	2.76	2.88	0.456	0.463	1.62	1.69	23.35	23.94
		F ₂	2.83	2.95	0.463	0.471	1.71	1.78	23.68	24.27
	C ₂	F ₁	2.86	2.99	0.467	0.471	1.77	1.84	23.88	24.53
		F ₂	2.90	3.02	0.472	0.483	1.78	1.86	24.33	24.96
	C ₃	F ₁	2.96	3.07	0.478	0.490	1.81	1.90	24.44	25.18
		F ₂	3.00	3.12	0.483	0.495	1.85	1.92	24.70	25.28
LSD at 5%	0.18	0.20	0.008	0.028	0.16	0.14	1.54	0.45		

Means within a row followed by a different letter (s) are statistically different at a 0.05 level

Since, I₁: Irrigation every 5 days; I₂: Irrigation every 7 days; I₃: Irrigation every 10; C₁: 1 compost: 3 sand (v/v); C₂: 1 compost: 1 sand (v/v); C₃: 3 compost: 1 sand (v/v); F₁: Without melatonin and F₂: Melatonin at rate of 100 µM N.S*: Non-significant

3. Indicators of oxidation and antioxidant enzymes

The data presented in Table 8 show the effect of the treatments studied on the indicators of oxidation [malondialdehyde (MDA, µmol. g⁻¹ F.W)] and antioxidant enzymes [i.e., peroxidase (POX, unit mg⁻¹ protein⁻¹) and catalase (CAT, unit mg⁻¹ protein⁻¹)] in the leaves of *Schefflera actinophylla* L plants. The data show that the highest values of MDA, POX and CAT were obtained when *Schefflera actinophylla* L. plants were subjected to the irrigation schedule of every 10 days (I₃ treatment), followed by those irrigated every 7 days (I₂ treatment). In contrast, the conventional irrigation treatment (I₁ treatment) resulted in the lowest levels for MDA, POX, and CAT. These results suggest that less frequent irrigation, as seen in the I₃ and I₂ treatments, induced oxidative stress in the plants, leading to higher levels of MDA, an indicator of lipid peroxidation. In response to this stress, the plants increased the activity of antioxidant enzymes, such as POX and CAT, as a defense mechanism. On the other hand, the I₁ treatment, characterized by more frequent irrigation, maintained lower levels of oxidative stress, resulting in reduced MDA and lower activity of antioxidant enzymes.

The plants that were grown in pots filled with a mixture of compost and sand in a ratio of 3:1 v/v (C₃ treatment) showed the highest values for peroxidase (POX) and catalase (CAT) activity. Following the C₃ treatment in terms of these enzyme activities, the C₂

treatment showed intermediate values, and the C₁ treatment resulted in the lowest activity levels for both POX and CAT. Conversely, for malondialdehyde (MDA) levels, the highest values were recorded in the C₁ treatment, followed by the C₂ treatment, and the lowest MDA levels were observed in the C₃ treatment. These results suggest that the C₃ treatment, which contains a higher proportion of compost, increases the activity of antioxidant enzymes (POX and CAT), probably in response to potential oxidative stress. This suggests that the plants are better equipped to combat oxidative damage when grown in a substrate rich in organic matter. Conversely, the C₁ treatment, with a lower compost ratio, appears to result in higher oxidative stress, as evidenced by elevated MDA levels, which is indicative of lipid peroxidation.

Furthermore, the application of melatonin resulted in an increase in the levels of peroxidase (POX) and catalase (CAT) when compared to untreated plants. Conversely, malondialdehyde (MDA) levels decreased in melatonin-treated plants as compared to untreated plants. These observations suggest that melatonin has a positive effect on the plant's antioxidant defense mechanisms. It increases the activity of antioxidant enzymes (POX and CAT), which are involved in reducing oxidative stress by neutralizing harmful reactive oxygen species (ROS). As a result, the levels of MDA, an indicator of lipid

peroxidation and oxidative damage, are reduced in melatonin-treated plants, indicating improved protection against oxidative stress.

Table 8. Effect of irrigation schedules, varying compost ratios and melatonin on malondialdehyde (MDA, $\mu\text{mol. g}^{-1}$ F.W), peroxidase (POX, unit mg^{-1} protein $^{-1}$) and catalase (CAT, unit mg^{-1} protein $^{-1}$) content in leaves of *Schefflera actinophylla* L plants during two consecutive seasons (2022-2023)

Treatments	MDA, $\mu\text{mol.g}^{-1}$ F.W		POX, unit mg^{-1} protein $^{-1}$		CAT, unit mg^{-1} protein $^{-1}$			
	1 st	2 nd	1 st	2 nd	1 st	2 nd		
Main factor : Irrigation schedules								
I ₁	8.54c	8.90c	0.349c	0.366c	0.030c	0.032c		
I ₂	9.74b	10.13b	0.398b	0.417b	0.048b	0.053b		
I ₃	11.62a	12.10a	0.472a	0.495a	0.080a	0.086a		
LSD at 5%	0.09	0.10	0.005	0.008	0.001	0.003		
Sub main factor: Varying compost ratios								
C ₁	10.42a	10.84a	0.425a	0.445a	0.062a	0.067a		
C ₂	10.00b	10.40b	0.406b	0.427b	0.052b	0.057b		
C ₃	9.49c	9.89c	0.387c	0.406c	0.044c	0.048c		
LSD at 5%	0.06	0.16	0.004	0.004	0.001	0.002		
Sub-sub main factor: Melatonin applications								
F ₁	10.09a	10.51a	0.402b	0.422b	0.050b	0.054b		
F ₂	9.84b	10.24a	0.411a	0.431a	0.055a	0.060a		
LSD at 5%	0.09	N.S	0.003	0.002	0.001	0.002		
Interaction								
I ₁	C ₁	F ₁	8.85	9.22	0.355	0.372	0.036	0.039
		F ₂	8.66	9.01	0.362	0.379	0.038	0.040
	C ₂	F ₁	8.57	8.97	0.348	0.366	0.026	0.028
		F ₂	8.53	8.86	0.350	0.368	0.033	0.037
	C ₃	F ₁	8.37	8.71	0.338	0.354	0.021	0.023
		F ₂	8.24	8.63	0.340	0.359	0.024	0.027
I ₂	C ₁	F ₁	10.27	10.69	0.418	0.440	0.056	0.061
		F ₂	10.02	10.40	0.429	0.450	0.059	0.064
	C ₂	F ₁	9.94	10.34	0.389	0.407	0.045	0.048
		F ₂	9.80	10.14	0.406	0.427	0.052	0.057
	C ₃	F ₁	9.29	9.69	0.369	0.387	0.039	0.041
		F ₂	9.14	9.51	0.378	0.395	0.040	0.044
I ₃	C ₁	F ₁	12.63	13.11	0.489	0.511	0.089	0.096
		F ₂	12.08	12.62	0.496	0.520	0.092	0.099
	C ₂	F ₁	11.57	12.05	0.467	0.492	0.076	0.082
		F ₂	11.57	12.06	0.480	0.503	0.082	0.089
	C ₃	F ₁	11.35	11.83	0.444	0.466	0.066	0.071
		F ₂	10.52	10.95	0.456	0.478	0.072	0.079
LSD at 5%	0.23	1.01	0.008	0.007	0.003	0.006		

Means within a row followed by a different letter (s) are statistically different at a 0.05 level

Since, I₁: Irrigation every 5 days; I₂: Irrigation every 7 days; I₃: Irrigation every 10; C₁: 1 compost: 3 sand (v/v); C₂: 1 compost: 1 sand (v/v); C₃: 3 compost: 1 sand (v/v); F₁: Without melatonin and F₂: Melatonin at rate of 100 μM N.S*: Non-significant

Regarding the interaction effect, the plant performance was better in the combined treatments of I₂x C₂ (or C₃) x F₂ than in the combined treatments of I₁x C₁x F₁. The same pattern was observed for both seasons studied. These results are consistent with those of Arnao and Hernández-Ruiz (2015); Ahmad *et al.* (2023); Dradrach *et al.* (2022); Shah *et al.* (2023); Elshepiny (2023).

CONCLUSION

This study underscores the critical importance of addressing water scarcity challenges, especially in the context of ornamental plant cultivation in Egypt. The findings reveal several key points:

1. The performance of *Schefflera actinophylla* L plants is adversely affected as the intervals between irrigation

increase, highlighting the serious impact of water scarcity on plant growth and health.

2. The combination of compost and melatonin shows promise in mitigating the adverse effects of water deficit. In particular, the treatment with 3:1 compost: sand mixtures v/v (C₃) proves to be highly effective, resulting in the best plant growth and physiological responses.
3. The application of melatonin by spraying at a concentration of 100 μM significantly improves various growth parameters, chemical characteristics, and antioxidant levels, thus contributing to improved plant resistance to water deficit conditions.
4. To save irrigation water, the combination of a 7-day irrigation interval (I₂) with a pot composition of 3:1 compost: sand mixtures v/v (C₃), together with melatonin application at 100 μM (F₂), emerged as the most favorable treatment regimen for *Schefflera actinophylla* L plants.

RECOMMENDATIONS

Based on the results of the study, the following recommendations are proposed:

1. Given the observed decline in plant performance with extended irrigation intervals, it is critical to implement efficient water management practices. Consideration should be given to scheduling regular and consistent irrigation to meet the water needs of ornamental plants.
2. The use of compost, especially at higher ratios such as 3:1 compost: sand mixtures v/v (C₃), should be encouraged in *Schefflera* plant cultivation. Compost improves soil structure, nutrient and water retention, and overall plant health.
3. The incorporation of melatonin into cultivation practices, particularly at a concentration of 100 μM , is recommended to enhance plant growth of *Schefflera* and improve tolerance to water deficit conditions. Further research can explore optimal application methods and timing.
4. In the face of ongoing water scarcity challenges, ornamental plant growers should consider adapting their cultivation techniques to align with more sustainable and water-efficient practices. This may include the use of drought-tolerant plant varieties and advanced irrigation technologies.
5. Ongoing research is needed to explore the long-term effects of compost and melatonin applications on different ornamental plant species. Additionally, research into the economic feasibility of these treatments and their scalability in commercial settings is essential.

By implementing these recommendations, ornamental horticulture in Egypt can become more resilient to water scarcity, and ensure the sustainability of this vital industry.

REFERENCES

Abu-Khalaf, N., & Natsheh, B. (2022). Effect of different hormones on rooting development of *Schefflera actinophylla* plants. *Research on Crops*, 23(2), 407-412.

Ahmad, R., Alsahli, A. A., Alansi, S., & Altaf, M. A. (2023). Exogenous melatonin confers drought stress by promoting plant growth, photosynthetic efficiency and antioxidant defense system of pea (*Pisum sativum* L.). *Scientia Horticulturae*, 322, 112431.

- Ahmad, S., Muhammad, I., Wang, G. Y., Zeeshan, M., Yang, L., Ali, I., & Zhou, X. B. (2021). Ameliorative effect of melatonin improves drought tolerance by regulating growth, photosynthetic traits and leaf ultrastructure of maize seedlings. *BMC Plant Biology*, 21(1), 1-14.
- Ahmed, M. A., & Shahin, S. M. (2023). Effect of magnetite and paclobutrazol on growth and chemical composition of *Schefflera actinophylla* Endl. cv. Gold Capella plant under salt stress conditions. *Egyptian Journal of Chemistry*.
- Arao, M. B., & Hernández-Ruiz, J. (2015). Functions of melatonin in plants: a review. *Journal of pineal research*, 59(2), 133-150.
- Davey, M. W., Stals, E., Panis, B., Keulemans, J., & Swennen, R. L. (2005). High-throughput determination of malondialdehyde in plant tissues. *Analytical biochemistry*, 347(2), 201-207.
- Dewis, J., & Freitas, F. (1970). Physical and chemical methods of soil and water analysis. *FAO soils Bulletin*, (10).
- Dradrach, A., Iqbal, M., Lewińska, K., Jedroszka, N., Rana, M. A. K., & Tanzeem-ul-Haq, H. S. (2022). Effects of soil application of chitosan and foliar melatonin on growth, photosynthesis, and heavy metals accumulation in wheat growing on wastewater polluted soil. *Sustainability*, 14(14), 8293.
- Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics*, 11(1), 1-42.
- Elshepiny, M. A. (2023). Role of compost, biochar and sugar alcohols in raising the maize tolerance to water deficit conditions. *Egyptian Journal of Soil Science*, 63(1), 67-81.
- Gomez, K. A., & Gomez, A. A. (1984). "Statistical procedures for agricultural research". John Wiley and Sons, Inc., New York. pp:680.
- Güneş, A., Kordali, S., Turan, M., & Bozhüyük, A. U. (2019). Determination of antioxidant enzyme activity and phenolic contents of some species of the Asteraceae family from medicinal plants. *Industrial crops and products*, 137, 208-213.
- Harborne, J. B. (1984). *Methods of plant analysis. In Phytochemical methods: a guide to modern techniques of plant analysis* (pp. 1-36). Dordrecht: Springer Netherlands.
- Herbert, D., Philipps, P. J., & Strange, R. E. (1971). Carbohydrate analysis. *Methods Enzymol. B*, 5 (680), 265-27.
- Khattak, W. A., Sun, J., Abbas, A., Hameed, R., Jalal, A., Niaz, N., ... & Wang, Y. (2023). Melatonin alleviating drought stress in plants: A review. *South African Journal of Botany*, 161, 192-201.
- Khosravi, S., Haghighi, M., & Mottaghipisheh, J. (2023). Effects of melatonin foliar application on hot pepper growth and stress tolerance. *Plant Stress*, 9, 100192.
- Langaroudi, I. K., Piri, S., Chaeikar, S. S., & Salehi, B. (2023). Evaluating drought stress tolerance in different *Camellia sinensis* L. cultivars and effect of melatonin on strengthening antioxidant system. *Scientia Horticulturae*, 307, 111517.
- Moursy, M. A. M., ElFetyany, M., Meleha, A. M. I., & El-Bialy, M. A. (2023). Productivity and profitability of modern irrigation methods through the application of on-farm drip irrigation on some crops in the Northern Nile Delta of Egypt. *Alexandria Engineering Journal*, 62, 349-356.
- Paymaneh, Z., Sarcheshmehpour, M., Mohammadi, H., & Hesni, M. A. (2023). Vermicompost and/or compost and arbuscular mycorrhizal fungi are conducive to improving the growth of pistachio seedlings to drought stress. *Applied Soil Ecology*, 182, 104717.
- Peterburgski, A. V. (1968). "Handbook of Agronomic Chemistry". Kolos Publishing House, Moscow, (in Russian, pp. 29-86).
- Saad Eddin, M. R., Hassan, A. M., Hegazi, A. A., Eid, S. F., & Mokhtar, A. (2023). Effects of root watering system on yield, water use efficiency and fruit quality of date palm (cv Siwi): a case study in the arid climate, Egypt. *Irrigation Science*, 1-13.
- Schoefs, B. (2005). Plant pigments: properties, analysis, degradation. *Advances in food and nutrition research*, 49, 42-92.
- Shah, M. N., Wright, D. L., Hussain, S., Koutroubas, S. D., Seepaul, R., George, S., ... & Eswaramoorthy, R. (2023). Organic fertilizer sources improve the yield and quality attributes of maize (*Zea mays* L.) hybrids by improving soil properties and nutrient uptake under drought stress. *Journal of King Saud University-Science*, 35(4), 102570.
- Soussani, F. E., Boutasknit, A., Ben-Laouane, R., Benkirane, R., Baslam, M., & Meddich, A. (2023). *Arbuscular mycorrhizal* fungi and compost-based biostimulants enhance fitness, physiological responses, yield, and quality traits of drought-stressed tomato plants. *Plants*, 12(9), 1856.
- Wang, D., Chen, Q., Guo, Q., Xia, Y., Jing, D., & Liang, G. (2023). Metabolomic analyses reveal potential mechanisms induced by melatonin application for tolerance of water deficit in loquat (*Eriobotrya japonica* Lindl.). *Scientia Horticulturae*, 308, 111569.
- Yang, X., Ren, J., Li, J., Lin, X., Xia, X., Yan, W., ... & Ke, Q. (2023). Meta-analysis of the effect of melatonin application on abiotic stress tolerance in plants. *Plant Biotechnology Reports*, 17(1), 39-52.

استجابة نباتات الشفليرا النامية تحت ظروف العجز المائي للإضافات الأرضية للكمبوست والرش الورقي للميلاتونين

وليد محمد فهمي عبد الهادي¹، احمد جمال الدين عبد الخالق بدور² محمد عاطف الشربيني²

¹ قسم بحوث نباتات الزينة وتنسيق الحدائق، معهد بحوث البساتين، مركز البحوث الزراعية، الجيزة، مصر

² قسم بحوث خصوبة التربة وتغذية النبات، معهد بحوث الأراضي والمياه والبيئة، مركز البحوث الزراعية، ش الجامعة، الجيزة، 12619 مصر

الملخص

أصبح توفير مياه الري أمرًا بالغ الأهمية وخاصة عندما نتحدث عن نباتات الزينة، نتيجة للظروف المائية الصعبة في مصر. إن استخدام التعديلات الأرضية الفعالة منخفضة التكلفة، مثل الكمبوست، واستخدام جزيء "الميلاتونين" المكتشف حديثًا، يمكن أن يخفف من الآثار الضارة لندرة المياه. ولذلك هدفت هذه الدراسة إلى تقييم مدى فعالية دمج الكمبوست مع الميلاتونين في التخفيف من الآثار الضارة للعجز المائي على نبات الشفليرا. تم تقييم فترات ري مختلفة والتي كانت كما يلي، الري كل 5 أيام (I₁)، 7 أيام (I₂)، 10 أيام (I₃)، كعامل رئيسي. أيضا تم دراسة التركيبة النهائية لبيئة النمو داخل لأصيص، التي تتضمن الكمبوست والرمل، مع خيارات تشمل خليط من الكمبوست والرمل بنسبه 1:3 (حجم/حجم) (C₁)، وخليط من الكمبوست والرمل بنسبه 1:1 (حجم/حجم) (C₂)، وخليط من الكمبوست والرمل بنسبه 1:3 (حجم/حجم) (C₃)، كعامل منشق أول. بالإضافة إلى ذلك، تم دراسة رش الميلاتونين بتركيزات صفر (F₁) و 100 ميكرومول (F₂) كعامل منشق ثاني. شهد أداء النباتات انخفاضًا مع زيادة فترات الري. علاوة على ذلك، أظهرت النباتات النامية بالأصص التي تحتوي على خليط من الكمبوست والرمل بنسبه 1:3 (حجم/حجم) (C₃) أفضل أداء، تليها معاملة C₂، وأخيرًا C₁. علاوة على ذلك، أدى رش الميلاتونين إلى زيادة في جميع الصفات المدروسة من حيث النمو والخصائص الكيميائية، ومستويات مضادات الأكسدة، باستثناء المالنونيدالدهيد، الذي انخفض مقارنة بالنباتات غير المعاملة. وفي سياق مختلف، أظهر أداء النباتات تحت المعاملة المشتركة I₂ x C₂ أو C₃ x F₂ تحسنًا أفضل من الأداء تحت المعاملة المشتركة I₁ x C₁. وأخيرًا، تؤكد هذه النتائج الدور المحتمل للكمبوست والميلاتونين في التخفيف من التأثيرات الضارة لندرة المياه.