

Journal of Plant Production

Journal homepage & Available online at: www.jpp.journals.ekb.eg

Efficiency of Glycinebetaine in Increasing Potato (*Solanum tuberosum* L.) Plant Tolerance to Drought

Metwaly, E. E. ¹; W. A. El-Saady¹; Z. S. A. El-Shal² and Mona A. A. S. Moussa²



Cross Mark

¹Vegetable and Ornamental Dept., Agric. Faculty, Mansoura Univ., 35516, Mansoura, Egypt

²Potato and Vegetatively propagated vegetable Research. Department, Horticulture Research. Institute., Agriculture Research. Center, Giza, Egypt

ABSTRACT

Improving plant water use efficiency (WUE) by osmoprotectant represents the main approach to sustainable productivity. Field trials were conducted to elucidate the efficiency of glycinebetaine (GlyBet) concentrations (0, 200, 400, 600 mg/l) for overcoming drought injury (1500, 1200, and 900 m³/fed) on potato plant productivity. Drought stress significantly decreased potato plant growth (plant height, and leaves number, leaf area, axillary stem number/plant, as well as foliage fresh weight, leaves dry matter), photosynthetic pigment (chlorophyll a, chlorophyll b, and carotenoid), ion% (nitrogen, phosphorous, and potassium), and yield components (tuber weight plant⁻¹, tuber number plant⁻¹, marketable and non-marketable yield). Alternatively, drought significantly increased antioxidant enzyme activity, non-marketable yield, tuber dry matter, tuber hardness and density, vitamin C, and total soluble solids associated with boosting WUE. The greatest reduction was documented under severe drought (900 m³/fed). Foliar spraying with GlyBet significantly increased all growth and yield traits except non-marketable yield which is decreased. Additionally, GlyBet spraying increased photosynthetic pigment and proline concentration, ion percentage, and activity of antioxidant enzymes associated with improving relative water content and WUE. The greatest values were recorded with 600 mg/l GlyBet application over other concentrations or nontreated plants. Regarding the interaction effects, the current findings revealed that GlyBet supplementation at all concentrations alongside 1200 and 900 m³/fed as consumptive water, nullifies the drastic impact on plant growth and productivity as well as some biochemical traits. Accordingly, 600 mg/l GlyBet foliar spraying as an eco-friendly and cost-effective osmoprotectant twice has the potential to mitigate drought injury and increase WUE.

Keywords: Glycine betaine, drought, potato, yield and antioxidant enzymes

INTRODUCTION

Potato (*Solanum tuberosum* L.) ranks as the 1st highest-produced non-cereal food crop and the 4th supreme imperative food crop next to wheat, corn, and rice worldwide (FAOSTAT, 2021). It is grown in 100 countries and their worldwide production boosted from 270 million tonnes in 1961 to 370 million tonnes in 2019, nourishing more than a billion people (Nasir and Toth 2022). Potato is designated a vigorous source of carbohydrates, dietetic fibers, protein, vitamins, and nutrients (Beals, 2019). As a result of its potential to offer a remarkable production per unit input, boosting potato production in developed and developing countries may make a substantial contribution to meeting the nutritional needs of the growing population (Devaux et al., 2014). Conversely, because of their thin and shallow roots, utmost potato cultivars are considered drought-sensitive, which is categorized with low yield and quality within drought occurs (Cho et al., 2016).

Water scarcity represents the key restricting aspect for sustainable crop productivity worldwide (Alam et al., 2022; Metwaly et al., 2022; Farouk et al., 2023). Drought usually affects 40-60% of worldwide cultivated lands with 30-44 US\$ billion in economic losses (Gupta et al., 2020). Drought strongly induces several physio-biochemical and molecular changes, leading to poor plant performance and yield reduction by 10-60% (Metwaly et al., 2022; Farouk et al.,

2023). The magnitude of their losses, however, mostly depends on drought sternness and crop cultivars (Plich et al., 2020; Hill et al., 2021). Reduction in vegetative growth is the first visible sign of drought followed by senescence and yield reduction (Metwaly et al., 2022; Farouk et al., 2023). Drought may upset potato yield by declining vegetative growth, disturbing metabolic pathways, and decreasing leaf photosynthesis or leaf area duration (Deblonde and Ledent, 2001; Metwaly et al., 2022; Farouk et al., 2023). Besides vegetative growth, drought may shorten the growth cycle, reduce the tubers' size and numbers, and reduce tuber quality (Eiasu et al., 2007; Kumar et al., 2007) and occasionally the plant death (Saleem et al., 2020). Drought-affected plants have developed adaptation strategies for coping with water deficiency, i.e., avoidance, escape, and dehydration tolerance of protoplasts, leading to maintaining water status, ion and redox homeostasis, stabilization of membrane integrity, and hyper-accumulation of osmoprotectants (Saleem et al., 2020; Farouk et al., 2023). Egypt's agricultural extension requires an enormous amount of irrigation water, which is scarce to satisfy the predicted demand along with poor irrigation management. These conditions promoted improved water use efficiency (WUE) and increased plant productivity via cost-effective novel approaches including cultivation of drought tolerance cultivars, and effective agricultural performance (Metwaly et al., 2022; Farouk et al., 2023), and recently using drought-induced tolerant mediators like phytohormones and

* Corresponding author.

E-mail address: elsaady_2003@mans.edu.eg

DOI: 10.21608/jpp.2024.281709.1323



osmoprotectants (Metwaly et al., 2022; Farouk et al., 2023). One of the osmoprotectants used to regulate crop production under normal or stressful conditions is glycinebetaine (GlyBet).

Nowadays, GlyBet is used as an easy method for decreasing the injuries of stress factors on plant productivity and generating drought-stress tolerant crops (Metwaly et al., 2022; Niu et al., 2023). GlyBet is a low-cost, eco-friendly, and effective osmoprotectant, that is widely distributed in a wide range of biota (Chen and Murata, 2008). The efficiency of GlyBet depends on plant species, and developmental stage at the time of application, in addition, to their concentration used, and the number of applications (Metwaly et al., 2022). GlyBet application improves growth, survival, and crop production in the accumulator and non-accumulator plants in normal or stressful conditions (Yang et al., 2022), via regulating several physio-biochemical pathways, enhancing net CO₂ assimilation rates and photosynthetic capacity, maintaining ion homeostasis, and activation reactive oxygen species (ROS) scavenging system (Bai et al., 2022; Niu et al., 2023). In this regard, Shemi et al. (2021) mentioned that GlyBet increased vegetative features, and yield as well as

organic and non-organic solutes, steadying the secondary structure of both enzymes and proteins.

Some non-accumulator economical crops including potatoes can tolerate drought better when GlyBet is applied. So it has been hypothesized that exogenous application of GlyBet could lessen the destructive effects of drought on potatoes. Hence, the objective of this study was to assess the promising role of GlyBet in improving potato plant drought tolerance, in terms of plant growth, some physio-biochemical attributes, and plant productivity. This finding may suggest a novel water-saving strategy under water scarcity circumstances.

MATERIALS AND METHODS

Experimental layout

Two field experiments were done in clay loamy soil (Table 1) of the experimental station of the Vegetables and Floriculture Dept., Fac. Agric., Mansoura Univ. throughout the 2022 and 2023 seasons, to study the effect of irrigation regimes and GlyBet concentration on potato plant (*Solanum tuberosum*, Cv. Spunta) growth and yield as well as some physio-biochemical attributes under drip irrigation system.

Table 1. Experimental soil characteristics of 2022 and 2023 seasons.

Seasons	Silt %	Clay %	Sand %	Texture soil	F.C %	W.P %	AW %	PH	E.C (dSm-1)	O.M %	CaCO ₃ %	N ppm	P ppm	K ppm
2022	41.9	36.4	21.7	Clay loamy	36.1	19.0	17.1	8.02	1.48	1.6	3.11	50.9	6.1	291
2023	42.6	35.9	21.5	Clay loamy	35.5	18.8	16.7	8.03	1.51	1.8	3.32	52.0	6.4	296

F.C : Field Capacity; W.P.: Welting point; AW: Available water; OM: Organic matter

A strip plot on a randomized complete block design was utilized for the existing experiments. The vertical plots were assigned to irrigation regimes: well watering (1500 m³/fed), moderate drought (1200 m³/fed), and severe drought (900 m³/fed). Otherwise, the horizontal -plots were assigned to GlyBet application (0.0, 200, 400, and 600 mg/L GlyBet). The twelve treatments were replicated thrice making 36 experimental units (10.5 m² with 1.5 m extensive bounded districts).

Crop Husbandry

The experimental site was intensely ploughed and formerly added 20 m³/fed farmyard manure. On the 27th and 29th of December in the first and second years, correspondingly, individual tuber seeds (20-25g) were manually planted at a depth of 12-15 cm and 25 cm apart in each plot. Each plot involves 5 dripper lines with a 3 m length for each and 0.7 m space among two dripper lines. Two lines were utilized for the assessment of the morpho-physiological attributes and the other three lines were utilized for yield and quality assessment. Furthermore, a protective zone was omitted from each of the two experimental units to prevent irrigation water infiltration from overlapping

The experimental units took an equivalent volume of irrigation water throughout the emergence stage via furrow irrigation (using a water counter) for identical plant establishment. The rest volumes of the irrigation water (m³/fed.) were supplemented by using a water counter at 1.5 bar, water flowed through the drippers (4 liter/h.), depending on the growth stage at two days' intervals beginning on the 5th and 8th February, and finished 10 and 15 April in the 1st and 2nd seasons, correspondingly. The GlyBet concentrations with 0.01% (v/v) Tween-20 were sprayed twice (40 and 55 days from planting), to an overflow in the early morning using a

back-sprayer at the rate of 300 and 400 l/fed in the 1st and 2nd spraying respectively.

After two weeks from planting, 120, 125, and 100 kg fed⁻¹ of potassium sulfate (50% K₂O), phosphoric acid (50% P₂O₅), and ammonium nitrate (33.5% N) were added throughout the fertigation system at 2-day intervals, Following the Ministry of Agriculture and Land Reclamation, Egypt's approval

Data Recording

At 85 days from planting, five erratically selected plants plot⁻¹ were harvested and designated for assessing growth and physiological attributes in the shoot system.

1. Vegetative growth features

Plant height (cm), number of leaves and axillary stem as well as leaves area (cm²) /plant, besides foliage fresh weight (g/plant), leaves dry matter were determine.

2. Photosynthetic pigment concentration

Chlorophyll a, chlorophyll b, and carotenoids were extracted from the 4th upper leaves using Lichtenthaler (1987) procedure with ice-cold methanol for 48 h. The absorbance of methanolic extract using a spectrophotometer at 470, 652, and 665 nm was recorded, and formerly its concentration was designed.

3. Shoot ion%

Ion % was appraised constantly with AOAC (1990). Ground dried shoots were wet digested with HClO₃/H₂SO₄ till clearness, and then brought to 100 ml with deionized water and retained for estimations. Nitrogen was measured by micro-kjldahl, in the meantime P was measured spectrophotometrically (Spekol 11, UK) via ammonium molybdate with ascorbic acid. K was assessed Flame photometrically.

4. Leaf relative water content (RWC) and proline concentration

The RWC (fresh mass – dry mass)/ (turgid mass – dry mass) × 100 according to Farouk et al. (2020). Meanwhile, proline (µmol g⁻¹ FW) was measured spectrophotometrically at 520 nm by the improved ninhydrin protocol of Magne and Larher (1992).

5. Antioxidant enzymes:

Catalase (CAT) activity was assessed spectrophotometrically at lab. conditions by checking the reduction in absorbance at 240 nm following the protocol of Aebi (1983). The peroxidase (POD) activity was evaluated via 4-methyl catechol as substrate (Onsa et al., 2004). Superoxide dismutase (SOD) activity was decided by quantifying the reticence in the photoreduction of nitroblue tetrazolium by the SOD enzyme (Kumar et al., 2012).

6. Yield, and its components

At 110 days after planting, tubers weight (g) and tubers number/ plant were designated by taking the mean of 5 hills, dry matter %, hardness (kg cm⁻²), density (tubers weight (g) per plant/tubers volume (cm³) per plant), marketable tubers yield (ton/fed.) as healthy tuber with a weight above or equal to 50 g without (growth cracks, irregularly curved shape, rotten, bottleneck shape, diseased, insect attacked, misshapen tuber, those having a weight less than 50 g and two or more knobs) and total tubers yield (ton/fed.) was (total fresh weight of tubers per plot/plot area) x 4200.

7. Tuber Quality

Quality traits of tuber, i.e., ascorbic acid (vitamin C), total soluble solids (TSS), and total carbohydrates were considered (AOAC, 1990). TSS (°Brix) was estimated using a digital refractometer (Model HI96801, Hanna Instruments). Alternatively, ascorbic acid (mg/100 g tuber fresh weight) was appraised via 2,6-dichlorophenol indophenol reagent.

Specific gravity was designed in line with Abdel-Aal (1971) protocol, specific gravity (g/cm³) = Tuber mass (g)/ tuber volume (cm³)

The oven-dried tuber was extracted by sulphuric acid (1N) in a boiling water bath for 5 h., and then filtrated using Whatman number 42 filter papers. The total carbohydrates were estimated using phenol-sulphuric acid (Sadasivam and Manickam, 1996).

8. Water use efficiency (WUE)

The WUE was deliberated by Howell (1994) equation as follow,

$$WUE = \frac{\text{Total tuber yield (ton/red)}}{\text{Crop water consumption (m}^3\text{/red/season)}}$$

Statistical Analysis

Exploiting Costat software, a two-way ANOVA was done for statistical analysis (CoHortSoftware, 2006). The means were separated at p ≤ 0.05 using the LSD pair-wise comparison test.

RESULTS AND DISCUSSION

1. Growth parameters

Table (2) indicates that potato plant growth is significantly (p ≤ 0.05) influenced by irrigation regimes, GlyBet spraying, and their interaction either in the 1st or 2nd years. Compared with well-watered plants, water deficit significantly declined growth trials. Hence, the lowest plant height (57.22 and 58.06 cm), leaves number plant⁻¹ (86.58, and 86.82), leaves area plant⁻¹ (1304 and 1318 cm²), number of axillary stems plant⁻¹ (7.32, and 7.56), foliage fresh weight (367.0 and 372.8 g), and leaf dry matter percentage (12.81, and 12.98%) were noted within severe drought in both seasons respectively over well-watered plants.

Table 2. Response of potato plant growth attributes by glycinebetaine and irrigation regimes in both experimental years.

Treatments	Plant height (cm).		Leaves No /plant		Leaves area (cm ²) / plant		Number of axillary stem/plant		Foliage FW g/ plant		leaves DM %	
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2
Water quantity (I, m ³ /fed)												
1500 (I ₁₅₀₀)	67.47	67.84	100.47	100.75	1602	1624	6.18	6.38	444.7	454.7	11.08	11.23
1200 (I ₁₂₀₀)	59.95	60.27	89.92	90.17	1487	1505	6.91	7.14	403.4	411.0	11.51	11.66
900 (I ₉₀₀)	57.72	58.06	86.58	86.82	1304	1318	7.32	7.56	367.0	372.8	12.81	12.98
LSD 5%	5.39	5.41	7.80	7.82	30	29	0.16	0.17	16.9	17.0	1.05	1.07
Glycinebetaine (GlyBet, ppm)												
0 (GlyBet ₀)	58.76	59.06	88.14	88.34	1368	1385	5.69	5.87	379.8	387.4	11.41	11.55
200 (GlyBet ₂₀₀)	59.82	60.15	89.74	89.97	1461	1479	6.47	6.68	392.9	400.6	11.48	11.63
400 (GlyBet ₄₀₀)	63.18	63.54	94.48	94.75	1492	1511	7.20	7.44	413.6	421.5	12.02	12.18
600 (GlyBet ₆₀₀)	65.07	65.47	96.93	94.75	1534	1554	7.85	8.12	433.8	441.9	12.28	12.45
LSD 5%	3.98	4.00	6.17	6.18	19	18	0.04	0.05	6.8	6.9	0.73	0.74
Interaction Effects												
I ₁₅₀₀ +GlyBet ₀	63.35	63.66	95.03	95.23	1475	1496	5.05	5.21	420.8	430.5	10.62	10.76
I ₁₅₀₀ +GlyBet ₂₀₀	64.97	65.31	97.45	97.71	1590	1611	5.75	5.93	444.1	454.0	10.70	10.85
I ₁₅₀₀ +GlyBet ₄₀₀	69.00	69.39	102.62	102.92	1630	1653	6.68	6.90	444.7	454.7	11.48	11.64
I ₁₅₀₀ +GlyBet ₆₀₀	72.55	72.99	106.78	107.12	1712	1736	7.23	7.48	469.4	479.7	11.51	11.66
I ₁₂₀₀ +GlyBet ₀	57.59	57.87	86.40	86.59	1411	1427	5.87	6.06	294.9	402.3	11.06	11.20
I ₁₂₀₀ +GlyBet ₂₀₀	58.73	59.03	88.10	88.33	1503	1521	6.66	6.87	397.6	405.1	11.27	11.42
I ₁₂₀₀ +GlyBet ₄₀₀	60.71	61.05	91.08	91.34	1505	1523	7.26	7.50	404.3	414.0	11.65	11.81
I ₁₂₀₀ +GlyBet ₆₀₀	62.75	63.12	94.13	94.43	1528	1547	7.88	8.12	414.5	422.4	12.04	12.21
I ₉₀₀ +GlyBet ₀	55.34	55.64	83.01	83.20	1219	1233	6.14	6.33	323.7	329.2	12.54	12.70
I ₉₀₀ +GlyBet ₂₀₀	55.78	56.10	83.67	83.89	1291	1305	7.02	7.24	337.0	242.6	12.47	12.63
I ₉₀₀ +GlyBet ₄₀₀	59.83	60.19	89.75	90.01	1342	1357	7.67	7.92	389.8	395.8	12.93	13.10
I ₉₀₀ +GlyBet ₆₀₀	59.92	60.31	89.89	90.18	1363	1378	8.47	8.75	417.5	423.7	13.29	13.47
LSD 5%	8.16	8.18	12.45	12.49	42	41	0.18	0.20	19.7	19.8	1.64	1.67

Foliar spraying of GlyBet in both seasons improved significantly growth attributes (Table 2). The supreme plant height (65.07 and 65.47 cm), leaves number plant⁻¹ (96.93, and 94.75), leaves area plant⁻¹ (1534 and 1554 cm²), the

number of axillary stems plant⁻¹ (7.85, and 8.12), foliage fresh weight (433.8 and 441.9 g), and leaf dry matter percentage (12.28, and 12.45%) in both seasons respectively were

documented once 600 mg/l GlyBet treatments over the rest of GlyBet concentrations or nontreated plants.

Foliar application of GlyBet with 1200 or 900 m³/fed irrigation regimes nullifies drought injuries over nontreated well-watered plants. Spraying of 600 mg/l GlyBet under severe drought boosting plant height by 8.27 and 8.39%; leaves number plant⁻¹ by 8.28 and 8.38%; leaves area by 11.81, and 11.75%, number of axillary stems plant⁻¹ by 37.94 and 38.23%, foliage fresh weight by 28.97 and 28.70%, and leaves DM% by 5.98 and 6.06% in both seasons over nontreated, severely drought-affected plants (Table, 2).

Drought substantially declined vegetative features, along with the former research (Alam et al., 2022; Metwaly et al., 2022; Farouk et al., 2023). Water scarcity has been recorded to have a deleterious impact on potato stem length (Chang et al., 2018), leaves number plant⁻¹ (Aliche et al., 2018), stem number plant⁻¹ (Chang et al., 2018), and leaf area (Kesiime et al., 2016). These drastic effects may be owing to accelerating numerous metabolic disorders and molecular modifications (Alam et al., 2022), disturbing water and nutrient uptake, and hyperaccumulation of ROS (Farouk et al., 2023). Moreover, drought induces phytohormon imbalance that slows down cell turgidity and impairs nutrient balance along with a decline in growth attributes (Saleem et al., 2020). Additionally, Hussain et al. (2012) indicate that water deficit raises root respiration rate, carbon resource utilization to drop, declined adenosine triple phosphate production as well as over-accumulation of ROS. Moreover, drought diminishes photosynthesis processes via a decline in chlorophyll assimilation or damages its molecules (Saleem et al., 2020). Drought also reduces RWC and increases the intercellular ionic concentration, which impedes ATP assimilation and ribulose biphosphate production capacity

(the primary acceptor of CO₂ within photosynthesis), which reduces photoassimilate production needed for plant development (Flexas et al., 2006).

Application of GlyBet especially 600 mg/l under normal or drought conditions considerably boosted growth attributes over untreated plants within irrigation regimes (Shemi et al., 2021). The enhancement of GlyBet on plant growth under normal or stressful conditions may be resulted from, maintaining water status (Shemi et al., 2021; Metwaly et al., 2022), hormonal balance, and boosting either cell division or elongation (Niu et al., 2023). Moreover, GlyBet application improved stomatal conductance, and sub-stomatal CO₂ concentration, which in turn enhances CO₂ diffusion along with photosynthesis improvement and plant development (Niu et al., 2023). As for the interactions, a few researchers are verifying the existing findings that demonstrate the use of GlyBet reduces the detrimental impacts of drought on plant establishment (Shemi et al., 2021).

2. Photosynthetic pigment:

Table 3 shows that the concentration of chlorophyll a, chlorophyll b, and carotenoids in potato leaves expressively decreases under drought conditions. The greatest concentration in both experimental seasons was recorded in 1500 m³/fed irrigation regimes, which was followed by 1200 m³/fed and then 900 m³/fed irrigation regimes.

Table 3 ascertains that spraying GlyBet on potato plants exhibited a positive effect on chlorophylls and carotenoids concentration over nontreated plants. The main effectiveness was 600 mg/L, which increased Chl a (11.30 and 11.43%), Chl b (12.17 and 11.95%), and carotenoids (8.77 and 8.91%) relative to untreated plants.

Table 3. Chlorophyll a, chlorophyll b, and carotenoid concentration of potato leaves as affected by glycinebetaine and irrigation regimes in both experimental years.

Treatments	Chl. a mg/100 FW		Chl.b mg/100 FW		Carotenoids mg/100g FW	
	S1	S2	S1	S2	S1	S2
Water quantity (I, m ³ /fed)						
1500 (I ₁₅₀₀)	33.65	34.08	16.06	16.94	13.23	14.06
1200 (I ₁₂₀₀)	29.97	30.36	14.91	15.44	12.40	13.18
900 (I ₉₀₀)	28.86	29.23	13.08	13.46	10.65	11.31
LSD 5%	2.66	2.70	0.30	0.31	0.38	0.41
Glycinebetaine ppm						
0 (GlyBet ₀)	29.38	29.74	13.72	14.30	11.62	12.34
200 (GlyBet ₂₀₀)	29.91	30.29	14.66	15.25	12.03	12.78
400 (GlyBet ₄₀₀)	31.32	31.72	14.97	15.57	12.08	12.84
600 (GlyBet ₆₀₀)	32.70	33.14	15.39	16.01	12.64	13.44
LSD 5%	2.21	2.24	0.19	0.20	0.68	0.73
Interaction Effects						
I ₁₅₀₀ +GlyBet ₀	31.67	32.06	14.79	15.64	12.98	13.78
I ₁₅₀₀ +GlyBet ₂₀₀	32.48	32.90	15.94	16.81	13.03	13.85
I ₁₅₀₀ +GlyBet ₄₀₀	33.68	34.12	16.35	17.24	13.07	13.89
I ₁₅₀₀ +GlyBet ₆₀₀	36.77	37.26	17.18	18.08	13.84	14.72
I ₁₂₀₀ +GlyBet ₀	28.79	29.15	14.15	14.66	11.62	12.34
I ₁₂₀₀ +GlyBet ₂₀₀	29.37	29.74	15.08	15.61	12.49	13.27
I ₁₂₀₀ +GlyBet ₄₀₀	30.36	30.75	15.09	15.63	12.53	13.33
I ₁₂₀₀ +GlyBet ₆₀₀	31.37	31.80	15.33	15.87	12.97	13.79
I ₉₀₀ +GlyBet ₀	27.67	28.01	12.23	12.60	10.27	10.91
I ₉₀₀ +GlyBet ₂₀₀	27.89	28.24	12.96	13.33	10.57	11.23
I ₉₀₀ +GlyBet ₄₀₀	29.91	30.30	13.46	13.85	10.64	11.31
I ₉₀₀ +GlyBet ₆₀₀	29.96	30.37	13.67	14.07	11.12	11.82
LSD 5%	4.16	4.21	0.42	0.43	0.87	0.93

The existing investigation established that GlyBet application raised chlorophyll a, chlorophyll b, and carotenoid levels in potato leaves compared to non-treated plants under these irrigation regimes. By enhancing chlorophyll a (8.27

and 8.42%), chlorophyll b (10.86 and 11.66%), and carotenoid (8.27 and 8.34%) content above nontreated severe drought-affected plants, spraying of 600 mg/L GlyBet lessens the destructive effects of drought.

The current finding and earlier research proved that drought induces a substantial deterioration in chlorophylls and carotenoids, that was lessened via GlyBet application (Table 3, Alam et al., 2022, Shemi et al., 2021). The drastic impact of drought on chlorophyll and carotenoid concentrations may be attributable to (1) acceleration of the malformation of chloroplast, and plastoglobules over-production (Farouk and Omar 2020); (2) activation of chlorophyllase biosynthesis (Haider et al., 2017); (3) devastation of light-harvesting pigment-protein complexes chiral macro-aggregates (Lai et al., 2007).

Alternatively, GlyBet supplementation increased photosynthetic pigments under normal or stressful conditions (Shemi et al., 2021). These rises might be resulted from, (1) well-organized ROS mitigation capacity via motivation of antioxidant enzymes and osmolyte buildup (Shemi et al., 2021); (2) recitals of a vital occupation i.e., cytokinin in enhancing chlorophyll assimilation, and raising chloroplast number cell⁻¹ (Hasanuzzaman et al., 2019); (3) inducing the accumulation of carotene that protect chloroplasts and chlorophyll against ROS; (4) preventing photoinhibition capacity along with increasing photosynthetic capacity (Niu et al., 2023).

3. ion percentage:

Drought stress expressively decreased the percentages of N, P, and K, and the supreme drops of N (13.62 and 13.63%), P (18.53 and 18.59%), and K (34.17 and 33.80%) noted within severe drought over controls in the 1st and 2nd years, correspondingly (Table 4).

Table 4 designates that GlyBet application expressively elevates the shoot's N, P, and K% relative to nontreated plants, and 600 mg/L GlyBet generated the highest ion %.

Table 4. Ion percentage of potato shoots as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	N %		P %		K %	
	S1	S2	S1	S1	S2	S2
Water quantity (I, m ³ /fed)						
1500 (I ₁₅₀₀)	3.685	3.959	0.464	0.484	4.77	4.91
1200 (I ₁₂₀₀)	3.306	3.551	0.410	0.428	3.76	3.89
900 (I ₉₀₀)	3.183	3.419	0.378	0.394	3.14	3.25
LSD 5%	0.300	0.322	0.035	0.037	0.13	0.13
Glycinebetaine (GlyBet, ppm)						
0 (GlyBet ₀)	3.240	3.479	0.399	0.416	3.39	3.44
200 (GlyBet ₂₀₀)	3.299	3.542	0.407	0.424	3.88	3.93
400 (GlyBet ₄₀₀)	3.454	3.711	0.426	0.445	3.96	4.14
600 (GlyBet ₆₀₀)	3.572	3.839	0.438	0.457	4.32	4.56
LSD 5%	0.215	0.23	0.026	0.028	0.25	0.23
Interaction Effects						
I ₁₅₀₀ +GlyBet ₀	3.493	3.750	0.441	0.460	4.25	4.31
I ₁₅₀₀ +GlyBet ₂₀₀	3.582	3.847	0.452	0.472	4.75	3.81
I ₁₅₀₀ +GlyBet ₄₀₀	3.715	3.991	0.469	0.489	4.86	5.29
I ₁₅₀₀ +GlyBet ₆₀₀	3.951	4.247	0.492	0.513	5.22	5.23
I ₁₂₀₀ +GlyBet ₀	3.176	3.410	0.395	0.412	3.01	3.05
I ₁₂₀₀ +GlyBet ₂₀₀	3.238	3.478	0.402	0.419	3.36	3.41
I ₁₂₀₀ +GlyBet ₄₀₀	3.348	3.597	0.416	0.434	4.08	4.13
I ₁₂₀₀ +GlyBet ₆₀₀	3.460	3.719	0.430	0.448	4.60	4.97
I ₉₀₀ +GlyBet ₀	3.051	3.276	0.363	0.378	2.92	2.95
I ₉₀₀ +GlyBet ₂₀₀	3.076	3.303	0.365	0.381	3.54	3.59
I ₉₀₀ +GlyBet ₄₀₀	3.299	3.544	0.392	0.409	2.95	2.99
I ₉₀₀ +GlyBet ₆₀₀	3.304	3.551	0.393	0.410	3.14	3.49
LSD 5%	0.468	0.503	0.058	0.061	0.36	0.38

GlyBet supplementation specifically 600 mg/L with 1200 or 900 m³/fed irrigation water invalidated the declined injury of water deficit over untreated plants within each irrigation regime.

Earlier studies have formerly confirmed the current outcomes that exhibit a drop in nutrient percentage once drought occurs (Metwaly et al., 2022). The defeat in N % under drought can result from an inhibition in nitrate reductase activity (Farahani et al., 2009). The deterioration in K % with water deficit may be elucidated by interrupting water status in stomata and plant cells (Sarani et al., 2014). Alternatively, the decline in nutrient percentage within drought conditions was mitigated by GlyBet supplementation. The role of GlyBet in boosting nutrients% is not implicit and some similar investigators have confirmed current findings (Khoshkham et al., 2021). This promotion may be resulted from a better development of root system, and stabilizing membrane permeability (Hasanuzzaman et al., 2019).

4. Relative water content (RWC), and proline concentration

Significant differences were established amongst water regimes for RWC, and proline content (Table 5). Gradually drought stress expressively reduced leaf RWC accompanied by proline over-accumulation relative to control plants. Moderate drought expressively improved proline (28.83% in the 1st year and 27.42% in the 2nd year) over control plants.

Table 5. Relative water content, and proline concentration of potato shoots as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	Relative water content %		Proline (μmol g-lfw)	
	S1	S2	S1	S2
Water quantity (I, m ³ /fed)				
1500 (I ₁₅₀₀)	75.58	76.59	20.39	21.84
1200 (I ₁₂₀₀)	71.34	72.29	26.27	27.83
900 (I ₉₀₀)	66.03	66.91	22.90	24.35
LSD 5%	1.78	1.80	3.03	3.03
Glycinebetaine (GlyBet, ppm)				
0 (GlyBet ₀)	68.92	69.81	21.99	23.24
200 (GlyBet ₂₀₀)	70.96	71.89	22.84	24.21
400 (GlyBet ₄₀₀)	71.56	72.53	23.37	24.93
600 (GlyBet ₆₀₀)	72.48	73.48	24.54	26.31
LSD 5%	2.79	2.83	1.61	1.60
Interaction Effects				
I ₁₅₀₀ +GlyBet ₀	74.39	75.34	19.16	20.36
I ₁₅₀₀ +GlyBet ₂₀₀	75.30	76.28	19.83	21.26
I ₁₅₀₀ +GlyBet ₄₀₀	75.53	76.55	20.52	22.02
I ₁₅₀₀ +GlyBet ₆₀₀	77.12	78.18	22.04	23.74
I ₁₂₀₀ +GlyBet ₀	67.75	68.62	24.62	25.86
I ₁₂₀₀ +GlyBet ₂₀₀	71.36	72.29	26.05	27.35
I ₁₂₀₀ +GlyBet ₄₀₀	72.75	73.74	26.54	28.24
I ₁₂₀₀ +GlyBet ₆₀₀	73.47	74.49	27.88	29.88
I ₉₀₀ +GlyBet ₀	64.62	65.45	22.20	23.50
I ₉₀₀ +GlyBet ₂₀₀	66.24	67.09	22.63	24.03
I ₉₀₀ +GlyBet ₄₀₀	66.40	67.30	23.04	24.54
I ₉₀₀ +GlyBet ₆₀₀	66.85	67.77	23.72	25.31
LSD 5%	3.61	3.66	4.03	4.04

Application of GlyBet significantly improved RWC, and proline over nontreated plants. The spraying with GlyBet at 600 mg/l enhanced RWC (5.16 and 5.25%), and proline (11.59 and 11.49%) over control in the 1st and 2nd seasons respectively.

Regarding the interaction, GlyBet spraying in moderate and severe drought abolishes the destructive impact

of water stress on RWC over nontreated drought-affected plants in both years (Table 5). Furthermore, GlyBet spraying to moderate or severe drought-affected plants significantly induces the over-accumulation of proline corresponding to nontreated control plants (Table 5). Conversely, the maximum proline (27.88 and 29.88 $\mu\text{mol g}^{-1}\text{fw}$) was achieved once 600 mg/l GlyBet spraying with 1200 m^3/fed irrigation water.

RWC was familiarized as a decent measure for crop water status within drought conditions. The injuries of drought on RWC % were lessened by GlyBet application, leading to an improvement in RWC %, relative to untreated drought-affected plants (Table 5). Earlier, it was discovered that RWC % declined under drought (Alam et al., 2022, Shemi et al., 2021; Metwaly et al., 2022). The RWC of GlyBet-treated plants within well-watered or water deficient was retained at greater altitudes to an outstanding level over stressed plants without GlyBet application, a motivation that GlyBet will enhance potato water status under water deficit. Regularly, GlyBet spraying lessened the dropping of potatoes RWC % under drought, as confirmed by Yang et al. (2022). Due to its solid hydrophilicity and solubility, GlyBet has been shown to maintain plant water status in arid regions as well as contribute to the osmotic adjustment capability of plant tissues (Genard et al., 1991). Alasvandyari et al. (2017) verified that GlyBet may increase K^+ accumulation in drought-affected plants, hence supporting the plants' capability to preserve the leaf's water content. Furthermore, GlyBet treatments invigorated root system establishment and then reinforced the capability of water absorption, and upregulation of aquaporin genes, to enhancement water conservation and boost WUE (Yang et al., 2022, Hasanuzzaman et al., 2019). This designates that GlyBet spraying might endure potato leaf membrane stabilities under stressful conditions.

A pervasive phenomenon of stress-affected plants is the buildup of a superior amount of compatible osmolytes, that are a lesser molecular weight tremendously soluble, and usually harmless at a great level without affecting metabolic pathways (Ashraf and Foolad, 2007). Proline represents a signaling molecule that impacts cell multiplication and elicits definite gene expression, which is needed for plant endurance within water scarcity or GlyBet supplementation (Metwaly et al., 2022). Proline buildup in plants plays a decisive function in preserving osmotic adjustment, so, plant cells can absorb extra water from the soil (Ullah et al., 2012). Additionally, proline has been deliberated as a carbon and nitrogen source for speedy rescue from stressful circumstances and performing as a preservative for membrane and macromolecules, as well as a ROS mitigator (Mousavi et al., 2009). This increase may be owing to a decline in proline oxidase and proline catabolizing enzymes (Debnath, 2008). Conversely, proline assimilation once GlyBet is applied under normal or stressful conditions is not entirely recognized and requires extra investigation.

5. Antioxidant Enzyme activities

Table 6 ascertains that drought ominously accelerated antioxidant enzyme activity (SOD, Cat, and POD) over well-watered plants. The supreme activity of each enzyme throughout the experimental time was documented under moderate water stress compared with severe drought or well-watered plants.

Application of GlyBet concentration raised progressively SOD, CAT, and POD activity in potato shoots in either the 1st or 2nd season. The utmost effective concentration in this regard is 600 mg/l GlyBet which accelerates SOD, CAT, and POD activities by 11.68, 14.89, and 41.38% in the first season, and by 7.68, 14.00, and 33.34% in the second season over non-treated plants.

The data in the same table indicated that GlyBet spraying under moderate or severe stress significantly improved antioxidant enzyme activities relative to nontreated plants under such irrigation regimes. The maximum activities were documented once treated plants with 600 mg/l GlyBet under moderate drought compared with other concentrations and water stress levels.

Table 6. Antioxidant enzyme activities of potato shoots as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	SOD (Unit mg-1 protein)		Catalase (Unit mg-1 protein)		Peroxidase (Unit mg-1 protein)	
	S1	S2	S1	S2	S1	S2
	Water quantity (m^3/fed)					
1500 (I ₁₅₀₀)	2.253	2.350	4.954	5.053	2.272	2.389
1200 (I ₁₂₀₀)	2.732	2.767	6.118	6.199	2.609	2.768
900 (I ₉₀₀)	2.613	2.647	5.552	5.626	2.371	2.497
LSD 5%	0.124	0.190	0.225	0.325	0.107	0.111
Glycinebetaine ppm						
0 (GlyBet ₀)	2.353	2.473	5.088	5.198	2.032	2.237
200 (GlyBet ₂₀₀)	2.547	2.580	5.550	5.622	2.368	2.478
400 (GlyBet ₄₀₀)	2.604	2.630	5.682	5.758	2.396	2.507
600 (GlyBet ₆₀₀)	2.628	2.663	5.846	5.926	2.873	2.983
LSD 5%	0.064	0.090	0.258	0.304	0.216	0.209
Interaction						
I ₁₅₀₀ +GlyBet ₀	2.091	2.387	4.455	4.645	1.900	2.101
I ₁₅₀₀ +GlyBet ₂₀₀	2.269	2.298	4.945	5.009	2.243	2.333
I ₁₅₀₀ +GlyBet ₄₀₀	2.358	2.389	5.304	5.375	2.250	2.339
I ₁₅₀₀ +GlyBet ₆₀₀	2.296	2.327	5.112	5.183	2.696	2.785
I ₁₂₀₀ +GlyBet ₀	2.490	2.520	5.475	5.546	2.150	2.386
I ₁₂₀₀ +GlyBet ₂₀₀	2.729	2.764	6.265	6.346	2.540	2.672
I ₁₂₀₀ +GlyBet ₄₀₀	2.809	2.845	6.119	6.201	2.600	2.732
I ₁₂₀₀ +GlyBet ₆₀₀	2.902	2.940	6.613	6.704	3.146	3.281
I ₉₀₀ +GlyBet ₀	2.479	2.510	5.334	5.403	2.046	2.224
I ₉₀₀ +GlyBet ₂₀₀	2.643	2.676	5.439	5.510	2.323	2.430
I ₉₀₀ +GlyBet ₄₀₀	2.644	2.679	5.623	5.699	2.340	2.449
I ₉₀₀ +GlyBet ₆₀₀	2.686	2.722	5.812	5.892	2.776	2.885
LSD 5%	0.115	0.216	0.587	0.822	0.301	0.302

Reactive oxygen species as signaling molecules are produced in plants via photorespiration and oxidative processes in mitochondria; for controlling developmental and metabolic pathways under normal or stressful conditions (Farouk et al., 2023). The over-production can reach toxic levels in the plants to counter the harmful impacts of ROS, and plants establish scavenging strategies i.e., to mitigate the ROS level including antioxidant enzymes and molecules (Hossain et al., 2013). The degree of ROS injury depends on the equilibrium amongst the assembly and elimination of ROS. Antioxidant enzymes can alleviate the drastic impact of ROS molecules (Mane et al., 2008). A current study proved that SOD, CAT, and POD activity was improved by drought, GlyBet, and their interactions, relative to untreated plants under such treatments (Table 6), thus conferring higher drought resistance

The CAT and POD convert H₂O₂ into water and molecular oxygen, and are supposed to limit cellular injury and enhance oxidative stress tolerance (Kusvuran et al., 2013). Superoxide radicals that emerge with metabolic processes are transformed into hydrogen peroxide and molecular oxygen by the SOD enzyme (Kusvuran et al., 2013). The metabolism of hydrogen peroxide is dependent on several functionally interrelated antioxidant enzymes like CAT and POD. As CAT and POD synchronize with SOD and play a prime defensive function in O₂⁻ and H₂O₂ scavenging process (Hoque et al., 2007). GlyBet spraying significantly increased antioxidant enzyme activity under

normal or stressful conditions. Since GlyBet-treated plants can sustain maximum antioxidant enzyme activity at a reasonable level to avert impairment induced by ROS accumulation (Hoque et al., 2007; Farooq et al., 2010).

6. Yield and its components:

Tables (7,8) outcomes clearly show that drought triggered an extreme drop in yield and components. Relative to control plants, there was a considerable reduction in tuber weight and tuber number plant⁻¹, marketable and non-marketable yield, total yield, tuber dry matter, tuber hardness, and tuber density in both years.

Table 7. Potato yield as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	Tubers weight (g) per plant		Tubers No./ plant		Marketable yield (ton/ fed.)		Non Marketable yield (ton/ fed.)		Total yield (ton/ fed.)	
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2
Water quantity (I, m ³ /fed)										
1500 (I ₁₅₀₀)	633.6	662.4	5.50	5.75	19.42	19.68	0.549	0.556	19.97	20.23
1200 (I ₁₂₀₀)	591.3	618.2	4.99	5.12	16.45	16.67	0.662	0.671	17.12	17.34
900 (I ₉₀₀)	543.3	568.0	4.81	4.93	13.24	13.41	0.845	0.856	14.09	14.27
LSD 5%	5.6	5.9	0.44	0.45	0.57	0.59	0.011	0.012	0.58	0.60
Glycinebetaine (GlyBet, ppm)										
0 (GlyBet ₀)	552.8	577.7	4.89	4.95	14.43	14.61	0.791	0.803	15.22	15.41
200 (GlyBet ₂₀₀)	571.4	597.2	4.99	5.08	15.80	16.00	0.724	0.734	16.53	16.74
400 (GlyBet ₄₀₀)	602.8	630.4	5.21	5.37	16.91	17.13	0.654	0.662	17.57	17.80
600 (GlyBet ₆₀₀)	630.6	659.5	5.45	5.65	18.34	18.58	0.574	0.580	18.91	19.17
LSD 5%	6.6	7.0	0.36	0.38	0.25	0.26	0.006	0.007	0.25	0.26
Interaction Effects										
I ₁₅₀₀ +GlyBet ₀	565.4	590.9	5.27	5.33	18.55	18.78	0.625	0.634	19.17	19.41
I ₁₅₀₀ +GlyBet ₂₀₀	588.7	615.3	5.41	5.52	18.86	19.09	0.577	0.585	19.43	19.68
I ₁₅₀₀ +GlyBet ₄₀₀	669.7	700.3	5.62	5.78	19.29	19.51	0.529	0.536	19.79	20.05
I ₁₅₀₀ +GlyBet ₆₀₀	710.6	743.2	6.12	6.35	21.02	21.30	0.467	0.472	21.48	21.78
I ₁₂₀₀ +GlyBet ₀	478.9	604.9	4.79	4.84	13.91	14.08	0.773	0.783	14.69	14.87
I ₁₂₀₀ +GlyBet ₂₀₀	582.9	609.2	4.89	4.99	15.94	16.15	0.709	0.718	16.65	16.87
I ₁₂₀₀ +GlyBet ₄₀₀	595.6	622.9	5.05	5.21	17.54	17.77	0.622	0.630	18.16	18.40
I ₁₂₀₀ +GlyBet ₆₀₀	607.7	635.6	5.22	5.42	18.41	18.66	0.546	0.553	18.95	19.21
I ₉₀₀ +GlyBet ₀	514.1	537.2	4.61	4.65	10.83	10.96	0.977	0.990	11.81	11.95
I ₉₀₀ +GlyBet ₂₀₀	542.5	567.0	4.65	4.74	12.61	12.78	0.885	0.896	13.50	13.67
I ₉₀₀ +GlyBet ₄₀₀	543.1	567.9	4.98	5.13	13.94	14.12	0.810	0.820	14.75	14.94
I ₉₀₀ +GlyBet ₆₀₀	573.4	599.8	4.99	5.17	15.58	15.79	0.708	0.717	16.29	16.51
LSD 5%	11.6	12.2	0.69	0.71	0.63	0.64	0.011	0.012	0.64	0.65

Table 8. Potato yield charctaritics as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	Tuber dry matter %		Hardness kg cm ⁻²		Density (g/ cm ³)	
	S1	S2	S1	S2	S1	S2
Water quantity (I, m ³ /fed)						
1500 (I ₁₅₀₀)	19.51	19.76	5.211	5.354	0.886	0.896
1200 (I ₁₂₀₀)	20.62	20.89	5.785	5.944	0.983	0.994
900 (I ₉₀₀)	21.50	21.79	6.853	7.041	1.173	1.186
LSD 5%	1.05	1.03	0.205	0.211	0.040	0.041
Glycinebetaine (GlyBet, ppm)						
0 (GlyBet ₀)	18.46	18.59	5.501	5.648	0.950	0.960
200 (GlyBet ₂₀₀)	19.95	20.15	5.779	5.935	0.991	1.001
400 (GlyBet ₄₀₀)	21.37	21.69	6.102	6.270	1.038	1.049
600 (GlyBet ₆₀₀)	22.38	22.83	6.418	6.597	1.078	1.090
LSD 5%	0.51	0.56	0.557	0.572	0.089	0.091
Interaction Effects						
I ₁₅₀₀ +GlyBet ₀	17.74	17.86	4.921	5.053	0.851	0.860
I ₁₅₀₀ +GlyBet ₂₀₀	18.52	18.71	5.061	5.198	0.865	0.874
I ₁₅₀₀ +GlyBet ₄₀₀	20.10	20.40	5.356	5.504	0.911	0.920
I ₁₅₀₀ +GlyBet ₆₀₀	21.65	22.09	5.505	5.661	0.919	0.930
I ₁₂₀₀ +GlyBet ₀	18.65	18.78	5.351	5.494	0.922	0.932
I ₁₂₀₀ +GlyBet ₂₀₀	20.42	20.62	5.744	5.899	0.982	0.992
I ₁₂₀₀ +GlyBet ₄₀₀	21.38	21.70	5.864	6.026	0.997	1.008
I ₁₂₀₀ +GlyBet ₆₀₀	22.03	22.47	6.183	6.355	1.032	1.043
I ₉₀₀ +GlyBet ₀	19.01	19.14	6.230	6.397	1.078	1.089
I ₉₀₀ +GlyBet ₂₀₀	20.91	21.11	6.533	6.710	1.126	1.138
I ₉₀₀ +GlyBet ₄₀₀	22.62	22.69	7.086	7.281	1.207	1.220
I ₉₀₀ +GlyBet ₆₀₀	23.47	23.94	7.564	7.775	1.282	1.296
LSD 5%	1.77	1.73	0.524	0.538	0.093	0.095

GlyBet application ominously increased potato yield and their components over untreated treatment. The most effective concentration was 600 mg/L GlyBet, which increased tuber weight and tuber number plant⁻¹, marketable yield, total yield, tuber dry matter, tuber hardness, and tuber density by 14.07, 11.45, 27.09, 24.24, 21.23,16.66, and 13.47% in the first year and 14.15, 14.14, 27.17, 24.52, 22.80, 16.80, 13.54% in the 2nd year respectively, over untreated plants (Tables 7,8).

Potato yield components are less deleteriously impacted by drought when GlyBet is sprayed. With untreated drought-affected plants, yield components in both experimental years have declined as the addition of 600 mg/L GlyBet under severe drought

As indicated previously and in the current investigation, drought markedly reduced crop yield and its quantity, meanwhile GlyBet application nullifies the injury of drought and boosted yield attributes with normal conditions (Khoshkharam et al., 2021; Shemi et al., 2021; Metwaly et al., 2022). Potato yield impressively depends on its canopy structure as well as on yield contributing trials like tuber number plant⁻¹, average tuber size, tuber fresh and dry mass, and tuber yield plant⁻¹. All these trials were declined by drought which eventually dropped marketable yield (Aliche et al., 2018). The decline in yield and its quality under drought

may result from reduced canopy growth (branches and either leaf number and leaf area per plant) associated with dropping in photosynthesis rate and Calvin cycle enzyme activity (Shemi et al., 2021). Additionally, drought stress upset assimilates production and portioning into developed tuber (Shemi et al., 2021; Alam, et al., 2022), as well as reducing the number of stolons per plant (Schafleitner et al., 2007). Lastly, drought accelerates energy imbalance in chloroplasts that evokes ROS production along with crop senescence and yield reduction (Farouk et al., 2023).

GlyBet motivates several physio-biochemical pathways and morpho-anatomical alterations that may be the reason for the rise in crop yield (Shemi et al., 2021; Metwaly et al., 2022). These outcomes were approved with the values provided previously (Shemi et al., 2021, Khoshkharam et al., 2021). GlyBet attributed the yield improvement to both the defense of a higher net photosynthetic rate and an enhancement in the source-sink relationship (Shemi et al., 2021, Hasanuzzaman et al., 2019). GlyBet motivates growth and yield attributes due to its osmoprotective function and regulates ion homeostasis (Niu et al., 2023) accompanied by improving drought-affected plant CO₂ assimilation (Shemi et al., 2021; Niu et al., 2023), and owing to its role in biosynthesis and transport of hormones such as cytokinins that could have a role in photoassimilates transportation (Taiz and Zeiger, 2006).

7. Tubers chemical quality:

Table 9 shows that drought improved vitamin C and TSS but reduced total sugar and protein over control plants. The supreme vitamin C was eminent during moderate drought. Alternatively, the highest concentration of TSS was gained under severe drought, meanwhile, the maximum total sugars and protein were noted within well-watered conditions either in 1st or 2nd years.

Table 9 shows that GlyBet spraying substantially affected tuber chemical quality attributes. The maximum values were acquired through 600 mg/L GlyBet spraying in 1st or 2nd year, over former concentrations or nontreated plants.

Results in Table 9 display that spraying of GlyBet concentration within irrigation regimes expressively improved vitamin C, TSS, total sugars, and protein as related to nontreated plants in each irrigation regime. The maximum concentration of vitamin C was attained under moderate drought with 600 mg/L GlyBet, but the uppermost TSS was noted under severe drought and spraying 600 mg/l GlyBet. Otherwise, the highest concentration of total sugars and protein was documented under normal conditions and sprayed with 600 mg/L GlyBet.

There are scant reports on the effect of GlyBet and/or drought on tuber quality. Accordingly, Khoshkharam et al. (2021); Shemi et al. (2021); Metwaly et al. (2022) establish that TSS and vitamin C declined with water deficit, but increased with GlyBet spraying. Moreover, it is recorded that drought stress significantly declines protein and carbohydrate percentage in potato tuber, which might be triggered by the decline of CO₂ fixation, and a drop of assimilates translocation, as well as a deterioration in the starch and protein assimilates gene expression (Younis et al., 2000). On

the other hand, the raising in carbohydrate% of GlyBet pretreatment might be attributed to the rise of photosynthetic pigments (Table 3), leading to the upgrading of carbohydrate assimilation and build-up.

Table 9. Potato tuber quality as affected by glycinebetaine under irrigation regimes in both experimental years.

Treatments	Vit. C		TSS,%		Total sugars,%		Protein,%	
	S1	S2	S1	S1	S2	S2	S1	S2
Water quantity (L, m ³ /fed)								
1500 (I ₁₅₀₀)	13.28	13.46	4.03	4.25	5.18	4.40	13.32	13.49
1200 (I ₁₂₀₀)	15.82	16.03	4.56	4.81	4.65	4.85	12.41	12.56
900 (I ₉₀₀)	14.94	15.13	4.86	5.13	4.14	4.32	10.82	10.97
LSD 5%	1.06	1.07	0.37	0.39	0.09	0.10	0.14	0.15
Glycinebetaine (GlyBet, ppm)								
0 (GlyBet ₀)	12.29	12.44	4.19	4.36	4.38	4.56	11.40	11.54
200 (GlyBet ₂₀₀)	13.99	14.17	4.48	4.70	4.53	4.72	12.18	12.33
400 (GlyBet ₄₀₀)	15.48	15.68	4.54	4.82	4.74	4.95	12.42	12.58
600 (GlyBet ₆₀₀)	16.97	17.20	4.72	5.06	4.98	5.19	12.73	12.90
LSD 5%	1.15	1.16	0.35	0.37	0.10	0.11	0.09	0.10
Interaction Effects								
I ₁₅₀₀ +GlyBet ₀	10.91	11.05	3.72	3.87	4.85	5.05	12.31	12.46
I ₁₅₀₀ +GlyBet ₂₀₀	12.42	12.58	3.95	4.14	5.12	5.34	13.26	13.43
I ₁₅₀₀ +GlyBet ₄₀₀	14.17	14.35	4.10	4.34	5.13	5.35	13.57	13.74
I ₁₅₀₀ +GlyBet ₆₀₀	15.64	15.85	4.35	4.56	5.64	5.85	14.13	14.32
I ₁₂₀₀ +GlyBet ₀	13.26	13.43	4.33	4.50	4.56	4.75	11.77	11.92
I ₁₂₀₀ +GlyBet ₂₀₀	15.16	15.36	4.61	4.84	4.58	4.78	12.55	12.70
I ₁₂₀₀ +GlyBet ₄₀₀	16.57	16.79	4.62	4.89	4.68	4.88	12.56	12.72
I ₁₂₀₀ +GlyBet ₆₀₀	18.30	18.54	4.69	5.02	4.78	4.99	12.75	12.92
I ₉₀₀ +GlyBet ₀	12.69	12.84	4.53	4.71	3.73	3.89	10.12	10.25
I ₉₀₀ +GlyBet ₂₀₀	14.39	14.57	4.88	5.12	3.88	4.05	10.72	10.86
I ₉₀₀ +GlyBet ₄₀₀	15.69	15.90	4.92	5.21	4.42	4.61	11.14	11.28
I ₉₀₀ +GlyBet ₆₀₀	16.99	17.22	5.14	5.50	4.54	4.74	11.32	11.47
LSD 5%	2.01	2.04	0.70	0.75	0.17	0.19	0.21	0.22

Within drought conditions, photosynthesis is principally controlled by stomatal and mesophyll restrictions, i.e., in how far CO₂ remains accessible for chloroplasts when stomatal and mesophyll conductance is retained small to evade extreme transpiration. Only at high-stress levels non-stomatal metabolic limitations, such as reduced ribulose biphosphate carboxylase regeneration and ATP assimilation inflict carbon assimilation under drought. Conversely, at high irradiances, RuBP exists in excess, and CO₂ should endure the restrictive aspect for photosynthesis (Parry et al., 2007). Little CO₂ accessibility for chloroplasts favors the oxygenase reaction of ribulose biphosphate carboxylase and photorespiration, resulting in up to 50% decline in carbon gain over well-watered plants.

8. Water use efficiency (WUE)

Figure (1) shows that drought expressively improved WUE associated with control plants. The maximum WUE (17.85, and 17.65%) was gained under severe stress over control plants in both years respectively.

WUE significantly improved by GlyBet spraying over nontreated plants (Figure 1), where the supreme WUE was achieved by supplementation of 600 mg/l GlyBet which raised it by 26.43% (1st year) and by 26.57% in 1st and 2nd year.

Additionally, Figure (1 C,D) indicates that GlyBet spraying under 1200 or 900 m³/fed irrigation regimes enhanced WUE over nontreated well-watered plants. The maximum WUE (41.62% and 41.73% during 1st and 2nd years) was achieved through 600 mg/l GlyBet spraying under severe drought correspondingly.




Figure 1. Potato water use efficiency as affected by irrigation regimes (A), glycinebetaine concentration (B), and their interactions (C,D) in both seasons. I 1500, well-watered (1500 m³/fed); I 1200, moderate drought (1200 m³/fed); I 900, severe drought (900 m³/fed); fed., feddan; GlyBet0, non spraying with glycinebetaine; GlyBet 200, spraying with 200 ppm glycinebetaine; GlyBet 400, spraying with 400 ppm glycinebetaine; GlyBet 600, spraying with 600 ppm glycinebetaine.

One of the most imperative physio-biochemical strategies used by drought-affected plants is the enhancement of WUE (Metwaly et al., 2022). Current findings revealed that drought and/or GlyBet considerably improved potato plant WUE in both experimental years over non-treated plants. GlyBet application raises potato WUE via manipulating photosynthesis and improving root establishment, leading to restored water absorption (Ahmed et al., 2019). The improvement in WUE resulted from lessening transpiration water loss and concurrently preserving leaf water status as well as raising root/shoot ratio along with inducing drought tolerance (Boguszewska-Mankowska et al., 2020). The goal is to raise plant WUE in drought-influenced plants. This may be achieved in two ways: by making the crop more capable of producing biomass per unit of water, or by improving the plant's ability to adapt. However, the effect of a drought on a plant's WUE often varies depending on the cultivar of the plant and the intensity of the drought (Gholami Zali et al., 2018).

CONCLUSION

Current findings explicitly that the application of GlyBet is an efficacious approach for improving plant performance under drought. Generally, spraying drought-affected potato plants two times at 40 and 55 days from cultivation with 600 mg/L GlyBet might be a prospective manner for dropping the effects of drought and consequently refining WUE and yield components and their quality.

REFERENCES

Abdel-Aal, R.M. (1971). Effect of groundwater and parent material on different soil characteristics in the North Nile Delta. Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Aebi, H.E. (1983). Catalase. In: Methods of enzymatic analysis. Bergmeyer, H.U. (Ed.). Verlag Chemie Weinheim. Pp. 273-286.

Ahmed, N., Zhang, Y., Li, K., Zhou, Y., Zhang, M., Li, Z. (2019). Exogenous application of glycine betaine improved water use efficiency in winter wheat (*Triticum aestivum* L.) via modulating photosynthetic efficiency and antioxidative capacity under conventional and limited irrigation conditions. *Crop J.* 7, 635–650.

Alam, A., Ullaha, H., Thuenproma, N., Tisarumc, R., Chaumc, S., Datta, A. (2022). Seed priming with salicylic acid enhances growth, physiological traits, fruit yield, and quality parameters of cantaloupe under water-deficit stress. *S. Afr. J. Bot.* 150, 1–12.

Alasvandyari, F., Mahdavi, B., Hosseini, S.M. (2017). Glycine betaine affects the antioxidant system and ion accumulation and reduces salinity-induced damage in safflower seedlings. *Arch. Biol. Sci.* 69, 139–147.

Aliche, E.B., Oortwijn, M., Theeuwens, T.P., Bachem, C.W., Visser, R.G., van der Linden, C.G. (2018). Drought response in field grown potatoes and the interactions between canopy growth and yield. *Agric. Water Manag.* 206, 20–30.

AOAC (1990). Official Methods of Analysis, 20th ed.; Association of Official Analytical Chemists: Arlington, VA, USA.

Ashraf, M., Foolad, M.R. (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environ. Exp. Bot.* 59, 206-216.

Bai, M., Zeng, W., Chen, F., Ji, X., Zhuang, Z., Jin, B. (2022). Transcriptome expression profiles reveal response mechanisms to drought and drought-stress mitigation mechanisms by exogenous glycine betaine in maize. *Biotechnol. Lett.* 44, 367–386. doi: 10.1007/s10529-022-03221-6

Beals, K.A. (2019). Potatoes, nutrition and health. *Am. J. Potato Res.* 96, 102–110. doi: 10.1007/s12230-018-09705-4

Boguszewska-Mańkowska, D., Zarzyńska, K., Nosalewicz, A. (2020). Drought differentially affects root system size and architecture of potato cultivars with differing drought tolerance. *Am. J. Potato Res.* 97, 54–62.

Chang, D.C., Jin, Y.I., Nam, J.H., Cheon, C.G., Cho, J.H., Kim, S.J., Yu, H.-S. (2018). Early drought effect on canopy development and tuber growth of potato cultivars with different maturities. *Field Crops Res.* 215, 156–162.

Chen, T.H.H., Murata, N. (2008). Glycine betaine: an effective protectant against abiotic stress in plants. *Trends Plant Sci.* 13, 499-505.

- Cho, K.S., Han, E.H., Kwak, S.S., Cho, J.H., Im, J.S., Hong, S.Y., Sohn, H.B., Kim, Y.H., Lee, S.W. (2016) Expressing the sweet potato orange gene in transgenic potato improves drought tolerance and marketable tuber production. *C R Biol.* 339(5–6), 207–213
- Deblonde, P.M.K., Ledent, J.F. (2001) Effects of moderate drought conditions on green leaf number, stem height, leaf length and tuber yield of potato cultivars. *Eur. J. Agron.* 14, 31–41.
- Debnath, M. (2008). Responses of *Bacopa monnieri* to salinity and drought stress in vitro. *J. Medicinal Plants Res.* 11, 347-351.
- Devaux, A., Kromann, P., Ortiz, O. (2014). Potatoes for sustainable global food security. *Potato Res.* 57, 185-189.
- Eiasu, B.K., Soundy, P., Hammes, P.S. (2007). Response of potato (*Solanum tuberosum*) tuber yield components to gelpolymer soil amendments and irrigation regimes. *N.Z.J. Crop Hort.* 35, 25–31. doi: 10.1080/01140670709510164
- FAOSTAT (2021). Available online: <https://www.fao.org/faostat/en/#data/QCL> (accessed on 23 November 2021).
- Farahani, H.A., Valadabadi, S.A., Daneshian, J., Shiranirad, A.H., Khalvati, M.A. (2009). Medicinal and aromatic plants farming under drought conditions. *J. Hortic. For.* 1, 86–92.
- Farooq, M.S.M., Basra, A., Wahid, A., Cheema, Z.A., Cheema, M.A., Khaliq, A. (2008). Physiological role of exogenously applied glycinebetaine to improve drought tolerance in fine grain aromatic rice (*Oryza sativa* L.). *J. Agron. Crop Sci.* 194, 325–333.
- Farouk, S., AL-Huqail, A.A., El-Gamal, S.M.A. (2023). Potential role of biochar and silicon in improving physio-biochemical and yield characteristics of borage plants under different irrigation regimes. *Plants* 12, 1605. <https://doi.org/10.3390/plants12081605>
- Farouk, S., Elhindi, K.M., Alotaibi, M.A. (2020). Silicon supplementation mitigates salinity stress on *Ocimum basilicum* L. via improving water balance, ion homeostasis, and antioxidant defense system. *Ecotoxicol. Environ. Saf.* 206, 111396.
- Farouk, S., Omar, M.M. (2020). Sweet basil growth, physiological and ultrastructural modification, and oxidative defense system under water deficit and silicon forms treatment. *J. Plant Growth Regul.* 39, 1307–1331.
- Flexas, J., Bota, J., Galmes, J., Medrano, H., Ribas-Carbó, M. (2006). Keeping a positive carbon balance under adverse conditions: Responses of photosynthesis and respiration to water stress. *Physiol. Plant.* 127, 343–352.
- Genard, H., Le Saos, J., Billard, J.P., Tremolieres, A., Boucaud, J. (1991). Effect of salinity on lipid composition, glycine betaine content and photosynthetic activity in chloroplasts of *Suaeda maritima*. *Plant Physiol. Biochem.* 29, 421–427.
- Gholami Zali, A., Ehsanzadeh, P., Szumny, A., Matkowski, A. (2018). Genotype-specific response of *Foeniculum vulgare* grain yield and essential oil composition to proline treatment under different irrigation conditions. *Ind. Crops Prod.* 124, 177–185.
- Gupta, A., Rico-Medina, A., Caño-Delgado, A.I. (2020). The physiology of plant responses to drought. *Science* 368, 266–269.
- Haider, M.S., Zhang, C., Kurjogi, M.M. (2017). Insights into grapevine defense response against drought as revealed by biochemical, physiological and RNA-Seq analysis. *Sci. Rep.* 7, 13134.
- Hasanuzzaman, M., Rehman, K., Nahar, H.K., Alharby, H.F. (2019). Plant abiotic stress tolerance. Agronomic, molecular and biotechnological approaches. Springer Nature Switzerland AG.
- Hill, D., Nelson, D., Hammond, J., Bell, L. (2021). Morphophysiology of potato (*Solanum tuberosum*) in response to drought stress: paving the way forward. *Front. Plant Sci.* 15, 675690. doi: 10.3389/fpls.2021.675690
- Hoque, M.A., Okcma, E., Banu, M.N.A., Nakamura, Y., Shimoishi, Y., Murata, Y. (2007). Exogenous proline mitigates the detrimental effects of salt stress more than exogenous betaine by increasing antioxidant enzyme activities. *J. of Plant Physiology* 164, 553-561.
- Hossain, M.A., Mostofa, M.G., Fujita M. (2013). Cross protection by cold-shock to salinity and drought stress-induced oxidative stress in mustard (*Brassica campestris* L.) seedlings. *Molecular Plant Breeding* 4, 50-70.
- Howell, T. (1994). Irrigation engineering, evapotranspiration. In: Arntzen, C.J., Ritter, E.M. (Eds.), *Encyclopedia of Agricultural Science*. Academic Press, New York.
- Hussain, S., Ali, A., Ibrahim, M., Saleem, M.F., Alias Haji, M.A., Bukhsh, A. (2012). Exogenous application of abscisic acid for drought tolerance in sunflower (*Helianthus annuus* L.): A review. *The J. of Animal and Plant Sci.* 22, 806-826.
- Kesiime, V.E., Tusiime, G., Kashaija, I.N., Edema, R., Gibson, P., Namugga, P., Kakuhenzire, R. (2016). Characterization and evaluation of potato genotypes (*Solanum tuberosum* L) for tolerance to drought in Uganda. *Am. J. Potato Res.* 93, 543–551.
- Khoshkharam, M., Shahrajabian, M.H., Esfandiari, M. (2021). The effects of methanol and amino acid glycine betaine on qualitative characteristics and yield of sugar beet (*Beta vulgaris* L.) cultivars. *Notulae Sci. Biol.* 13, 10949. <https://doi.org/10.15835/nsb13210949>
- Kumar, A., Dutt, S., Bagler, G., Ahuja, P.S., Kumar, S. (2012). Engineering a thermo-stable superoxide dismutase functional at subzero to >50°C, which also tolerates autoclaving. *Sci. Rep.* 2, 387.
- Kumar, S., Asrey, R.A.M., Mandal, G. (2007). Effect of differential irrigation regimes on potato (*Solanum tuberosum*) yield and postHarvest attributes. *Indian J. Agric. Sci.* 77, 366–368.
- Kusvuran, S., Ellialtioglu, S., Polat, Z. (2013). Antioxidative enzyme activity, lipid peroxidation, and proline accumulation in the callus tissues of salt and drought tolerant and sensitive pumpkin genotypes under chilling stress. *Hort. Environ. Biotechnol.* 54, 319-325.
- Lai, Q., Zhi-yi, B., Zhu-Jun, Z., Qiong-Qiu, Q., Bi-Zeng, M. (2007). Effects of osmotic stress on antioxidant enzymes activities in leaf discs of PSAG12-IPT modified gerbera. *J. Zhejiang Univ. Sci.* 8(7), 458-464. doi:10.1631/jzus.2007.B0458.
- Lichtenthaler, H.K. (1987). Chlorophylls and carotenoids: Pigments of photosynthetic biomembrane. *Methods Enzymol.* 148, 350–352.
- Magne, C., Larher, F. (1992). High sugar content of extracts interferes with colorimetric determination of amino acids and free proline. *Anal. Biochem.* 200, 358-362. DOI:10.1016/0003-2697(92)90285-f
- Mane, S.P., Robinet, C.V., Ulanov, A., Schafleitner, R., Tincopa, L., Gaudin, A., Novberto, G., Alvarado, C., Solis, C., Bolivar, L.A. (2008). Molecular and physiological adaptation to prolonged drought stress in the leaves of two andean potato genotypes. *Funct. Plant Biol.* 35, 669–688.

- Metwaly, E.S.E., Al-Yasi, H.M., Ali, E.F., Farouk, H.A., Farouk, S. (2022). Deteriorating harmful effects of drought in cucumber by spraying glycinebetaine. Agriculture 12, 2166. <https://doi.org/10.3390/agriculture12122166>
- Mousavi, E.A., Kalantari, K.M., Jafari, S.R. (2009). Change of some osmolytes accumulation in water stressed colza (*Brassica napus* L.) as affected by 24-epibrassinolide. Iranian J. of Sci. and Technology, Transaction A, 33, A1.1-11
- Nasir, M.W., Toth, Z. (2022). Effect of drought stress on potato production: A review. Agronomy 12, 635. <https://doi.org/10.3390/agronomy12030635>
- Niu, T., Zhang, J., Li, J., Gao, X., Ma, H., Gao, Y., Chang, Y., Xie, J. (2023). Effects of exogenous glycine betaine and cycloleucine on photosynthetic capacity, amino acid composition, and hormone metabolism in *Solanum melongena* L. Scientific Reports 13, 7626. <https://doi.org/10.1038/s41598-023-34509-w>
- Onsa, G.H., Saari, N., Selamat, J., Bakar, J. (2004). Purification and characterization of membrane-bound peroxidases from *Metroxylon sagu*. Food Chem. 85, 365-376.
- Parry, M.A.J., Madgwick, P.J., Carvalho, J.F.C., Andralojc, P.J. (2007). Prospects for increasing photosynthesis by overcoming the limitations of Rubisco. J. Agric. Sci. 145, 31-43.
- Plich, J., Boguszewska-Mankowska, D., Marczewski, W. (2020). Relations between photosynthetic parameters and drought-induced tuber yield decrease in Katahdin-derived potato cultivars. Pot. Res. 63, 463-477. doi: 10.1007/s11540-020-09451-3
- Sadasivam, S., Manickam, A. (1996). Biochemical methods. 2nd edition, New Age International. India.
- Saleem, M.H., Fahad, S., Khan, S.U., Ahmar, S., Khan, M.H.U., Rehman, M., Maqbool, Z., Liu, L. (2020). Morpho-physiological traits, gaseous exchange attributes, and phytoremediation potential of jute (*Corchorus capsularis* L.) grown in different concentrations of copper-contaminated soil. Ecotoxicol. Environ. Saf. 189, 109915.
- Sarani, M., Namrudi, M., Hashemi, S.M., Raoofi, M.M. (2014). The effect of drought stress on chlorophyll content, root growth, glucosinolate and proline in crop plants. Int. J. Farming Allied Sci. J. 3, 994-997.
- Schafleitner, R., Rosales, R.O.G., Gaudin, A., Aliaga C.A.A., Martinez, G.N., Marca, L.R.T., Bolivar, L.A., Delgado, F.M., Simon, R., Bonierbale, M. (2007). Capturing candidate drought tolerance traits in two native Andean potato clones by transcription profiling of field grown plants under water stress. Plant Physiol. Biochem. 45, 673-690.
- Shemi, R., Wang, L., Gheith, E.M., Hussain, H.A., Hussain, S., Irfan, M., Cholidah, L., Zhang, K., Zhang, S., Wang, L. (2021). Effects of salicylic acid, zinc and glycine betaine on morphophysiological growth and yield of maize under drought stress. Sci. Rep. 11, 3195.
- Taiz, L., Zeiger, E. (2006). Plant physiology, 4th ed.; Sinauer Associates, Inc.:Sunderland, MA,USA.
- Ullah, F., Bano, A., Nosheen, A. (2012). Effects of plant growth regulators on growth and oil quality of canola (*Brassica napus* L.) under drought stress. Pak. J. Bot. 44(6),1873-1880
- Yang, Y., Xie, J., Li, J., Zhang, J., Zhang, X., Yao, Y., Wang, C., Niu, T., Bakpa, P.E. (2022). Trehalose alleviates salt tolerance by improving photosynthetic performance and maintaining mineral ion homeostasis in tomato plants. Front. Plant Sci. 13, 974507. <https://doi.org/10.3389/fpls.2022.974507>
- Younis, M.E., El-Shahaby, O.A., Abohamed, S.A., Ibrahim, A.H. (2000). Effects of water stress on growth, pigments and 14CO₂ assimilation in three sorghum cultivars. J. Agron. Crop Sci. 185, 73-82. <https://doi.org/10.1046/j.1439-037x.2000.00400.x>

كفاءة الجليسين بيتين في زيادة تحمل نبات البطاطس للجفاف

السعيد السيد متولى¹، وليد على السعدى¹، زيدان شهاب احمد الشال² و منى عبد العزيز عبد الحليم سالم موسى²

¹ قسم الخضار والزينة - كلية الزراعة - جامعة المنصورة - 35516 - المنصورة - مصر
² قسم بحوث البطاطس والخضار خضرية التكاثر - معهد بحوث البساتين - مركز البحوث الزراعية - الجيزة - مصر

الملخص

تحسين كفاءة استخدام النبات للمياه من خلال المواد المنظمة للاسموزية يمثل النهج الرئيسي لإستدامة الانتاجية. أجريت تجارب حقلية لتقييم كفاءة استخدام تراكيز مادة الجليسين بيتين (0، 600، 400، 200 جزء في المليون) في التغلب على التأثير الضار للجفاف (1500، 1200، 900 م³/فدان) على انتاجية البطاطس. أدى الجفاف إلى انخفاض معنوي في نمو البطاطس (ارتفاع النبات، عدد الأوراق لكل نبات، المساحة الورقية/النبات، عدد الافرع/النبات، الوزن الطازج للمجموع الخضري، المادة الجافة)، وتركيز صبغات البناء الضوئي (الكلوروفيل أ، والكلوروفيل ب، والكاروتينويد) والنسبة المئوية للأيونات (النيتروجين، الفوسفور، والبوتاسيوم)، ومكونات المحصول (وزن الدرنة/نبات، عدد الدرنة/نبات، المحصول القابل وغير قابل للتسويق). ايضا، أدى الجفاف إلى زيادة كبيرة في نشاط الإنزيمات المضادة للأكسدة، ومحصول الدرنة غير القابل للتسويق، والمادة الجافة للدرنات، وصلابة الدرنة وكثافتها، وفيتامين C، والنسبة المئوية للمادة الصلبة، وزيادة كفاءة استخدام المياه. وكانت أكثر الإنخفاضات المسجلة تحت ظروف الجفاف الشديد. أدى الرش الورقي بالجليسين بيتين إلى زيادة معنوية في جميع صفات النمو والمحصول، باستثناء المحصول غير القابل للتسويق والذي ينخفض عند الرش بالجليسين بيتين. ايضا، أدى رش الجليسين بيتين إلى زيادة تركيز صبغات البناء الضوئي، ونسبة الأيونات، وتركيز البرولين، ونشاط الإنزيمات المضادة للأكسدة، تحسين المحتوى المائي النسبي وكفاءة استخدام المياه. وكانت أعلى القيم لتلك الصفات تم تسجيلها مع 600 ملجم/لتر من الجليسين بيتين مقارنة بالتركيز الأخرى أو النباتات غير المعاملة. أوضحت النتائج أن الرش بالجليسين بيتين بجميع تركيزاته يقلل من التأثيرات الضارة للجفاف المعتدل أو الشديد، على نمو النبات وإنتاجيته وكذلك بعض الصفات البيوكيميائية وبالتالي، فإن استخدام 600 ملجم/لتر من الجليسين بيتين مرتين باعتباره مادة حافظة للأسموزية، وصديقة للبيئة، ومنخفضة التكلفة، لديه القدرة على التخفيف من أضرار الجفاف وزيادة كفاءة استخدام المياه.

الكلمات الدالة: الجليسين بيتين، الجفاف، البطاطس، المحصول، الإنزيمات المضادة للأكسدة