

# Journal of Plant Production

Journal homepage & Available online at: [www.jpp.journals.ekb.eg](http://www.jpp.journals.ekb.eg)

## Increasing Growth, Yield and Bioactive Compounds of Strawberry by Bio-Stimulants

Mohamed I. A. Mohamed\* and M. K. F. El-Tawashy

Department of Vegetable Crops, Faculty of Agriculture, Cairo University, Giza, Egypt.



### ABSTRACT

Bio-stimulants have obtained global attention due to their beneficial impacts on the growth and development of several of crops. The current experiment was carried out during two winter seasons to assess the impact of different bio-stimulant substances on plant vigor, yield, and quality attributes in two strawberry cultivars (Fortuna and Festival) in a plastic greenhouse. Our treatments were: 1) humic acid 15% (10 ml L<sup>-1</sup>); 2) fulvic acid 15% (10 ml L<sup>-1</sup>); 3) potassium citrate (2 g L<sup>-1</sup>); 4) potassium silicate (2 g L<sup>-1</sup>); 5) salicylic acid (1 g L<sup>-1</sup>); 6) calcium carbonate (2 g L<sup>-1</sup>); and 7) control (water). Strawberry plants were treated weekly with each bio-stimulant, started after 15 days of transplanting and repeated 10 times. The most effective bio-stimulants were calcium carbonate and potassium silicate. Calcium carbonate enhanced all studied traits except dry matter, alkaloids concentration, antioxidant activity in leaves, and titratable acidity, while potassium silicate improved all traits except dry matter and alkaloids in leaves. Potassium citrate showed the highest value of flavonoids in leaves, firmness, and TSS. Humic acid exhibited the highest antioxidant activity in leaves and titratable acidity. Fulvic acid showed the highest total carbohydrates. The interaction between the cultivars and bio-stimulants was different in all traits. Significant differences were found among cultivars in all traits except total alkaloids, phenolics compounds, antioxidant activity in leaves, early yield, total yield, average fruit weight, TSS, and vitamin C in fruits (1st year) and number of leaves/plants, antioxidant activity, carbohydrates in leaves and total yield (2nd year).

**Keywords:** strawberry; bio-stimulants; productivity and quality.



### INTRODUCTION

The strawberry plant (*Fragaria × ananassa* Duch.) is considered one of the most globally distributed crops (Chandler et al., 2012). In addition to being a natural source of nutrients and phytochemicals like folic acid, potassium, ascorbic acid, and fiber, strawberries also have health benefits that include preventing cancer, heart disease, blood sugar and pressure regulation, inflammation reduction, cholesterol reduction, and aging prevention (AbdusSamee et al., 2023). Egypt is one of the significant players in the world strawberry market, ranked as the fourth largest strawberry producing country after China, USA, and Turkey. In 2022 Egypt produced 637842.16 tons and exported 45210.15 tons with an export value of about 117540000 USD, making Egypt the sixth largest strawberry exporter worldwide. The Ministry of Agriculture in Egypt announced that the production of strawberry plants during 2019/2020 season was 433945 tons, cultivated in 26756 feddans, while average production was 17.08 tons / feddan.

Bio-stimulants have obtained global attention due to their beneficial impacts on the growth and development of several crops. Bio-stimulants involve the utilization or use of chemical substances, physical methods, or biological agents to induce a positive response to improve growth, development, and quality of several crops, and induce biotic and abiotic stress tolerance alongside increased nutrient uptake (Soppelsa et al., 2019; Garza-Alonso et al., 2022). Recently, many former studies documented the positive effects of the application of bio-stimulants on different crops including chemical and biochemical substances.

Humic substances are complex organic molecules, naturally developed in soil as a result of the long-term microbial degradation of plant and animal waste. Humic substances are categorized based on solubility characteristics of humin, fulvic, and humic acids (Ore et al., 2023). Zydlik and Zydlik (2023) reported that humic acid application was a suitable alternative for mineral fertilization, which led to maximized soil enzyme activity, an increase in leaf area by 60%, and a significant enhancement in flower number, fruit number, and fruit set, which positively affected on yield over mineral fertilizers. Ullah et al. (2017) evaluated the effect of three levels of humic acid (1.5, 3.0, and 4.5 ml L<sup>-1</sup>), and found that using 3 ml L<sup>-1</sup> produced more strawberry fruits and the highest value of total carotenoids. Chakraborty et al. (2023) studied the response of two cultivars of strawberry (Camarosa and Nabila) to humic acid and seaweed extract and found that foliar application of 2.0 ml L<sup>-1</sup> humic acid with 1.0 ml L<sup>-1</sup> seaweed extract significantly enhanced the plant growing, number of leaves, flowers and fruits, as well as, fruit weight, fruits length and yield than control, while the maximum total soluble solids (TSS %), vitamin c, total sugar were exhibited with 2.0 ml L<sup>-1</sup> humic acid + 2.0 ml L<sup>-1</sup> seaweed extract.

Salicylic acid is a plant hormone and produced naturally in plants, and is considered a phenolic compound (Hassoon and Abduljabbar, 2020). Salicylic acid enhances plant vigor and fruit quality traits (Mohamed et al., 2017; Youssef et al., 2017) and improves the strawberry plant's tolerance for different abiotic stresses such as salinity stress (Roshdy et al., 2021), drought stress (Dakheel et al., 2022), and enhanced strawberry plant resistance for biotic stresses such as gray mold disease (Emam et al., 2023) and spider mite (De Resende et al., 2021).

\* Corresponding author.

E-mail address: [Mohamed.abdallah@agr.cu.edu.eg](mailto:Mohamed.abdallah@agr.cu.edu.eg)

DOI: 10.21608/jpp.2024.282168.1325

Potassium citrate as a bio-stimulate enhanced strawberry plant's growth parameters such as plant length, number of crowns, runners, and leaves as well as, leaves fresh, and dry weight for leaves and enhanced the early, total, and marketable yield, especially with using mineral source for nitrogen over compost manure, while the Spraying of potassium citrate improved fruit weight, TSS%, and vitamin C with compost manure (Awad et al., 2010). Potassium citrate induced relative resistance to the two-spotted spider mite in strawberry plants (Abdelwines and Ahmed, 2022). Foliar application with potassium citrate improves the growth and productivity of several vegetable crops like potatoes (Mansour and El-Metwaly, 2019), hot peppers (Ghoname et al., 2009), tomatoes (Zakher and Elashry, 2016) and peas (El-Sayed et al., 2019).

Potassium silicate played an important role as a bio-stimulant in strawberry plants, achieved significant plant growth improvement, maximized fruit yield, and enhanced fruit quality when used at 0.6 g/L as a foliar application (Nada, 2020). EL-Sayed et al., (2023) found that utilizing potassium silicate on strawberry plants as foliar application significantly improved leaf number, chlorophyll, biomass, dry matter, and leaf area, in addition to fruit length, weight, and diameter, which led to a significant increment in both total and marketable yield. Hussain et al. (2023) reported that spraying strawberry plants three times with potassium Silicate decreases whitefly and aphid incidence by an increasing of the epidermal cuticle layer thickness and leaf thickness. Also, potassium silicate promoted different abiotic and biotic stress tolerances, like salinity (Yaghubi et al., 2019), water deficit (Dehghanipoodeh et al., 2018), and leaf blight disease resistance (Abd-El-Kareem et al., 2019).

A few former studies were found on the effect of foliar spray of calcium carbonate on plants. Foliar spray with calcium carbonate on gooseberry plants significantly increased fruit number/plant, weight, and size of fruit (Shukla et al., 2011). Calcium carbonate induced relative resistance to the two-spotted spider mites in strawberry plants (Abdelwines and Ahmed, 2022). Calcium carbonate was utilized as a biological control agent for powdery mildew in strawberry plants and found a reduction in disease incidence (Pertot et al., 2008). This experiment aimed at comparing the effects of different bio-stimulants i.e., humic acid, fulvic acid, salicylic acid, potassium citrate, potassium silicate, and calcium carbonate on plant vigor, yield, and quality attributes of strawberry.

## MATERIALS AND METHODS

### 1. Plant materials and the experimental layout.

Our experiment was performed during the two winter seasons of 2017/2018 and 2018/2019 to assess and compare the effects of different bio-stimulants on strawberry plant vigor, yield, and quality attributes in a plastic greenhouse of the vegetable crops department, Cairo University, Giza province, Egypt. The two strawberry cultivars (Fortuna and Festival) seedlings were transplanted in the first week of October in four rows/terraces with 25 cm between transplants and between rows on each terrace. Each experimental plot was about 6 m<sup>2</sup> (5 m length x 1.2 m width). Terraces were covered with black plastic mulch. Drip irrigation was applied with two drip lines/ terrace; also spry irrigation system was used for 10 days only after transplanting to accelerate plant development. Common agricultural practices including

diseases and pest management were applied according to the Ministry of Agriculture and Land Reclamation, Egypt.

### 2. Treatments

The experimental treatments were: 1) humic acid 15% (10 ml L<sup>-1</sup>); 2) fulvic acid 15% (10 ml L<sup>-1</sup>); 3) potassium citrate (2 g L<sup>-1</sup>); 4) potassium silicate (2 g L<sup>-1</sup>); 5) salicylic acid (1 g L<sup>-1</sup>); 6) Calcium carbonate (2 g L<sup>-1</sup>); and 7) control (water). All treatments were added as foliar application except, humic acid was added manually as soil application. Strawberry plants were sprayed weekly, starting 15 days after the transplanting, and repeated 10 times. The experiment was performed using a randomized complete block design with 4 replicates in a split-plot design. Two strawberry cultivars as main plots, while bio-stimulants as split-plots.

### 3. Data recorded

#### Vegetative traits and leaves chemical attributes

Agronomic traits were measured after three months of strawberry transplanting, except leaf area/plant measured after two months to measure the quick response. Five plants were randomly selected to assess the number of leaves/plants, leaf dry matter, and leaf area/ plant (Pandey and Singh, 2011). Photosynthetic pigments (chlorophyll a, chlorophyll b, total chlorophyll, and total carotenoids) were assessed spectrophotometrically following the method of Mitic et al. (2013). The Folin-Ciocalteu method was used to determine phenolic concentration (Meda et al., 2005); meanwhile, the colorimetric method using aluminum chloride was used to measure flavonoids concentration (Chang et al., 2002). The phenol-sulfuric acid method was applied to assess the total carbohydrates (Dubois et al., 1956). Antioxidant activity was evaluated by the method of stable 2,2-diphenyl-1-picrylhydrazyl radical (Kedare and Singh, 2011).

#### Yield and fruit quality attributes

Strawberry fruits were harvested at full mature stage from the whole plot for each treatment and the first 5 pickings were expressed as early yield (kg/m<sup>2</sup>). Total yield (kg/m<sup>2</sup>) was accumulatively determined from each plot for each treatment during the harvesting season. Average fruit weight was assessed as the mean weight of ten fruits. Fruit firmness was determined by a food pressure tester model FT011. Total soluble solids (TSS) were measured by a digital refractometer (model PR101, Co. Ltd., Tokyo, Japan). Titratable acidity was assessed in fresh strawberry fruits using 0.1 N NaOH solution with phenolphthalein as an indicator (AOAC, 1990). Ascorbic acid (Vitamin C) was assessed in ripe strawberry fruits using 2, 6 dichlorophenol indophenol dye by a titrimetric method (AOAC, 1990). Anthocyanin was measured in strawberry fruits according to Pirie and Mullins's method 1976.

### 4. Statistical analysis

Data were statistically analyzed by MSTAT-C v. 2.1 software program (Michigan State University, Michigan, USA) using two-way ANOVA. The means of treatment were compared by Duncan's multiple range test at the 0.05 level of probability (Gomez and Gomez, 1984).

## RESULTS AND DISCUSSION

Increasing demand for bio-stimulants due to their positive effects on plant growth parameters, especially in poor soils, and unsuitable conditions such as abiotic and biotic stresses, and reduce the amounts of required fertilizers for plants and improve their productivity and enhance their tolerance to different stresses (De Vasconcelos and Chaves, 2019).

### **Effect of bio-stimulants, cultivars and their interaction on number of leaves/plants and leaf area.**

The individual effect of stimulants on strawberry plants is shown in Figure 1. Potassium silicate demonstrated the highest number of leaves/plants in both seasons with no significant difference with salicylic acid and calcium carbonate in the second season (Figure 1A). Fulvic acid caused a significant reduction in the number of leaves/plants in both seasons. Potassium silicate recorded the highest value of leaf area in the first season compared to all other treatments followed by calcium carbonate (Figure 1B). Calcium carbonate and potassium citrate recorded the largest leaf area in the second season with no significant differences. The lowest value of leaf area per plant was recorded in fulvic acid treatment in the first season and control treatment in the second season.

Concerning the interaction between cultivars and bio-stimulants, Table 1 revealed that significant differences were found among bio-stimulants in the two cultivars for the number of leaves and leaf area per plant. Strawberry plants treated with potassium silicate displayed the highest number of leaves/ plants in the Fortuna cultivar, followed by calcium carbonate in Festival cultivar in the first season. In the second season, the highest significant no. of leaves/plants was recorded in salicylic acid and potassium silicate in the Fortuna cultivar and calcium carbonate in Festival cultivar without significant differences among them.

The plants of the Fortuna cultivar treated with potassium silicate exhibited the largest leaf area in the first season, followed by plants of Festival cultivar treated with calcium carbonate. While the treatment of fulvic acid, potassium citrate, and calcium carbonate recorded the largest leaf area in Festival cultivar in the second season compared to all other treatments without significant differences among them.

Cultivars exhibited a significant effect on the number of leaves/plants in the first year only, while Cultivars displayed a significant effect on leaf area in both years.

Potassium ( $K^+$ ) as a cation, and calcium ( $Ca^{2+}$ ) are the most prevalent inorganic compounds in plant cellular media (Sardans and Peñuelas, 2021). Many former studies confirmed potassium's role in numerous physiological processes such as control of cellular growth, controlling water balance, in addition to metabolite and nutrient transport (Sardans and Peñuelas, 2021). Potassium application led to improved plant growth characteristics like number of leaves, biomass of the plant, leaf area, and dry matter %. Growth improvement by K could be due to their role in physiological processes such as nitrogen usage, protein biosynthesis, and cell development (Coskun et al., 2017). EL-Sayed et al (2023) reported that improvement in strawberry growth parameters such as the number of leaves and leaf area by using three concentrations of potassium silicate as a foliar application. Nada (2020) reported an enhancement in the number of leaves and leaf area due to spraying strawberry plants with four concentrations of potassium silicate. Previous studies confirmed an increment in both leaf area and the number of leaves after foliar spraying with potassium silicate separately and in mixed treatment with acetyl-salicylic acid on tomato plants (El-Sayed et al., 2023). In contrast, no effects on leaf area and leaf number were shown in strawberry plants using potassium silicate under water stress (Dehghanipoodeh et al., 2018). Also, potassium citrate alone and combined with boron led to improved vegetative parameters in Sweet Charlie strawberry cultivar (Awad et al.,2010).

A vital role was played by calcium as an essential element required for plant development in normal and stressful conditions. It is important for membrane stability and cell wall and plays a second messenger in many physiological processes (Thor, 2019). Furthermore, the transport of carbohydrates that are produced during photosynthesis to sinks is accelerated by calcium (Navazio et al.,2020). Former studies in agreement with our results, such as Rozbiany and Taha, (2020) who found that positive increment in all vegetative characters due to foliar application of calcium on strawberry plants. El-Sayed et al. (2023) reported that leaf area and number of leaves were enhanced as a result of calcium foliar application on strawberry plants.

### **Effect of bio-stimulants, cultivars and their interaction on leaves chemical attributes.**

#### **a- Photosynthetic pigments concentration:**


Data in Figure 2 showed the individual effects of bio-stimulants on strawberry plants for chlorophyll a, b, total chlorophyll, and total carotenoids. Calcium carbonate recorded the highest value of chlorophyll a content in both seasons, with no significant differences with salicylic acid and potassium silicate in the second season (Figure 2A). In contrast, fulvic acid showed a reduction in chlorophyll a over control plants in the first season. Calcium carbonate in both seasons, potassium citrate, and potassium silicate in the second season exhibited the highest content of chlorophyll b (Figure 2B). Regarding the total chlorophyll concentration, calcium carbonate in both seasons and potassium silicate in the second season gave the highest value compared to all other treatments (Figure 2C). Fulvic acid showed a reduction in total chlorophyll over control plants in the first season. Regarding total carotenoids, all bio-stimulants caused a significant reduction in total carotenoid concentration, except fulvic acid in the second season (Figure 2D).

Regarding the interaction between the cultivars and bio-stimulants, Table 2 displayed that significant differences were found among the bio-stimulants in the two cultivars for chlorophyll a, b, total chlorophyll and total carotenoids. Plants of the Festival cultivar treated with calcium carbonate exhibited the maximum value of chlorophyll a in both seasons. Foliar application of Fortuna plants with potassium citrate recorded the highest chlorophyll a concentration after calcium carbonate in first season, while foliar application of Festival plants with potassium silicate and potassium citrate in second season ranked second in chlorophyll a concentration. Concerning chlorophyll b concentration, in the first season spraying Fortuna plants treated with calcium carbonate recorded the highest content, and spraying Festival plants with calcium carbonate ranked the second. Meanwhile, the foliar application of both cultivars with potassium citrate, calcium carbonate, and potassium silicate on Festival showed the highest concentration of chlorophyll b in the second season. concerning total chlorophyll concentration, foliar application with calcium carbonate in both cultivars during the first season and in Festival cultivar in the second season, in addition to potassium silicate in Festival cultivar in the second season displayed the highest significant concentration of total chlorophyll. Concerning total carotenoids, plants of control in both cultivars in first season and control plants in Fortuna, in addition to spraying Festival with fulvic acid showed the highest concentration of total carotenoids in second season, while other treatments caused a significant reduction in total carotenoids in both cultivars.

Cultivars demonstrated significant effects on chlorophyll a, b, total chlorophyll, and total carotenoids in both years.

Our results agree with former studies, which found that foliar spraying on two different cultivars of strawberry (Albion and Festival) with calcium significantly improved chlorophyll concentration (Rozbiany and Taha, 2020). Foliar spraying with potassium silicate under normal or water stress conditions on strawberry achieved a significant increment in chlorophyll index (Dehghanipoodeh et al., 2018). Also, foliar

spraying with three concentrations (4, 6, and 8%) of potassium silicate and three concentrations of calcium significantly increased chlorophyll percentage in strawberry leaves (El-Sayed et al., 2023). Nada (2020) reported that using potassium silicate with three concentrations (0.2, 0.4 and 0.6 g L<sup>-1</sup>) enhanced chlorophyll a, b, and total, in addition to total carotenoids in strawberry leaves. In potatoes, foliar spraying with potassium citrate and potassium silicate enhanced total chlorophyll content in leaf tissues (Mousa et al., 2023).




Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

Fig. 1. Canopy parameters (A: number of leaves/plant and B: leaf area) of two strawberry cultivars treated with some stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Table 1. Canopy attributes (number of leaves/plant and leaf area) of strawberry cultivars Fortuna and Festival treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Stimulants / Cultivars	No. of Leaves/plant				Leaf area/ plant (cm <sup>2</sup> )			
	Fortuna		Festival		Fortuna		Festival	
	2017-2018							
Fulvic acid	17.6 $\pm$ 0.83	h	29.76 $\pm$ 1.26	g	219.17 $\pm$ 2.64	l	370.99 $\pm$ 4.12	j
Humic acid	31.16 $\pm$ 1.27	f	32.51 $\pm$ 2.09	e	388.48 $\pm$ 4.15	i	404.31 $\pm$ 7.55	g
Salicylic acid	36.85 $\pm$ 1.27	c	36.15 $\pm$ 1.52	cd	460.28 $\pm$ 4.76	c	450.65 $\pm$ 5.03	e
Potassium Citrate	31.91 $\pm$ 1.84	ef	35.17 $\pm$ 1.70	d	396.55 $\pm$ 5.75	h	437.94 $\pm$ 5.48	f
Potassium Silicate	40.63 $\pm$ 0.74	a	36.27 $\pm$ 1.73	cd	508.70 $\pm$ 2.82	a	451.63 $\pm$ 5.59	de
Calcium Carbonate	36.50 $\pm$ 1.50	c	38.30 $\pm$ 2.01	b	455.00 $\pm$ 4.90	d	476.94 $\pm$ 7.00	b
Water (Control)	29.27 $\pm$ 1.88	g	32.41 $\pm$ 1.45	e	363.49 $\pm$ 6.07	k	403.76 $\pm$ 4.69	g
Mean	31.98	b	34.37	a	398.81	b	428.03	a
	2018-2019							
Fulvic acid	18.43 $\pm$ 1.68	g	29.51 $\pm$ 2.58	f	314.20 $\pm$ 33.27	g	436.40 $\pm$ 18.33	a-c
Humic acid	30.91 $\pm$ 2.60	ef	34.27 $\pm$ 4.20	b-e	414.72 $\pm$ 24.37	cd	365.60 $\pm$ 34.90	ef
Salicylic acid	37.85 $\pm$ 2.59	ab	35.83 $\pm$ 3.11	bc	376.70 $\pm$ 41.79	d-f	373.20 $\pm$ 35.64	ef
Potassium Citrate	31.64 $\pm$ 3.86	d-f	34.87 $\pm$ 3.52	b-d	359.70 $\pm$ 40.68	f	457.70 $\pm$ 26.89	ab
Potassium Silicate	41.20 $\pm$ 1.52	a	35.95 $\pm$ 3.58	b	370.30 $\pm$ 35.12	ef	418.90 $\pm$ 28.94	bc
Calcium Carbonate	36.19 $\pm$ 3.07	b	40.10 $\pm$ 4.02	a	401.90 $\pm$ 32.66	c-e	469.5 $\pm$ 23.72	a
Water (Control)	31.13 $\pm$ 7.60	d-f	32.15 $\pm$ 3.01	c-f	353.20 $\pm$ 33.32	fg	378.50 $\pm$ 32.57	d-f
Mean	32.47	a	34.66	a	370.09	b	414.25	a

<sup>a</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Mean value  $\pm$  standard error.



Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

Fig. 2. Leaf pigments content (chlorophyll a (A), chlorophyll b (B), and total chlorophyll (C); and carotenoids (D)) of two strawberry cultivars treated with some stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Table 2. Leaf photosynthetic pigments concentration (chlorophylls a, b, and total chlorophyll; and carotenoids) of strawberry cultivars Fortuna and Festival treated with some stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.


Stimulants / Cultivars	Chlorophyll A (mg. 100 g <sup>-1</sup> FW)		Chlorophyll B (mg. 100 g <sup>-1</sup> FW)		Total chlorophyll (mg. 100 g <sup>-1</sup> FW)		Total carotenoids (mg. 100 g <sup>-1</sup> FW)	
	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival
	2017-2018							
Fulvic acid	70.29±2.96	h 65.12±2.74	j 34.76±1.46	i 32.21±1.36	j 105.05±4.42	j 97.32±4.10	k 23.67±1.00	c 23.32±0.98
Humic acid	73.93±3.11	e 71.31±3.00	g 36.57±1.54	g 35.26±1.48	h 110.50±4.65	f 106.58±4.49	i 22.91±0.96	e 22.31±0.94
Salicylic acid	76.32±3.21	d 72.40±3.05	f 37.74±1.59	f 36.81±1.55	g 114.10±4.80	e 109.22±4.60	g 21.97±0.93	g 19.20±0.81
Potassium Citrate	80.54±3.39	b 76.91±3.24	d 39.83±1.68	c 38.28±1.61	e 120.40±5.07	b 115.20±4.85	d 20.61±0.87	i 21.73±0.91
Potassium Silicate	70.67±2.98	h 79.01±3.33	c 39.01±1.64	d 39.08±1.65	d 109.70±4.62	fg 118.10±4.97	c 21.64±0.91	h 20.68±0.87
Calcium Carbonate	78.73±3.32	c 81.73±3.44	a 42.77±1.80	a 40.42±1.70	b 121.50±5.12	a 122.10±5.14	a 22.46±0.95	f 19.20±0.81
Water (Control)	70.67±2.98	h 68.08±2.87	i 34.94±1.47	hi 39.78±1.68	c 105.60±4.45	j 107.90±4.54	h 25.37±1.07	a 24.45±1.03
Mean	74.45	a 73.51	b 37.94	a 37.4	b 112.39	a 110.91	b 22.66	a 21.55
	2018-2019							
Fulvic acid	73.64±4.13	h 76.93±3.51	g 33.85±2.77	f 38.22±1.67	cd 109.80±6.80	fh 115.10±5.17	df 19.86±1.74	fg 24.67±1.08
Humic acid	79.68±4.49	e-g 78.26±3.33	fg 37.53±2.27	c-e 38.15±2.31	cd 113.50±6.48	e-g 115.10±5.54	df 21.26±1.29	df 19.82±1.25
Salicylic acid	80.65±3.86	ef 82.19±3.28	de 36.64±3.00	de 39.67±1.73	bc 119.40±6.75	bd 123.70±5.11	bc 18.26±1.60	gh 23.02±1.01
Potassium Citrate	76.76±4.12	gh 86.19±3.25	bc 41.59±2.23	ab 41.72±1.89	ab 118.40±6.28	ce 125.10±5.00	b 20.83±1.12	ef 22.03±1.01
Potassium Silicate	84.67±4.28	cd 87.93±3.05	b 37.92±2.18	cd 42.08±2.29	a 122.60±6.38	bc 131.70±5.39	a 18.49±1.06	gh 20.67±1.12
Calcium Carbonate	82.79±4.00	de 93.81±3.05	a 40.63±2.18	ab 40.70±3.40	ab 120.00±5.96	bd 130.90±6.24	a 20.97±1.12	ef 16.92±1.52
Water (Control)	70.26±2.80	i 76.53±3.55	gh 35.30±1.48	ef 33.99±3.33	f 107.10±4.37	h 108.60±6.19	gh 24.07±1.00	ab 22.74±2.23
Mean	78.34	b 83.11	a 37.63	b 39.22	a 115.82	b 121.44	a 20.53	b 21.41

<sup>2</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Mean value  $\pm$  standard error.

**b- Dry matter and carbohydrates in leaves**

Results in Figure 3 revealed the individual effects of bio-stimulants on strawberry plants for dry matter and carbohydrates. Fulvic acid only showed a significant increment in dry matter % in the first season. In the second season, fulvic acid and the control treatment showed the highest dry matter % while other treatments reduced dry matter % in strawberry leaves (Figure 3A). Regarding carbohydrates concentration, fulvic acid in both seasons showed the highest value followed by potassium citrate and calcium carbonate in both seasons with no significant difference between them, in addition to humic acid in the second season.

Regarding the interaction between cultivars and bio-stimulants, Table 3 demonstrated that significant differences were found among the effects of the bio-stimulants in the two cultivars for dry matter and carbohydrates. All bio-stimulants led to a significant reduction in dry matter % in both cultivars, except fulvic acid in both cultivars without a significant difference between fulvic acid treatment and control plants in the Fortuna cultivar only. The spraying Fortuna plants with fulvic acid in both seasons and Festival plants with potassium citrate in the second season displayed the highest carbohydrate concentration. Spraying Fortuna plants with humic acid in both seasons and Festival plants with potassium citrate in first the season ranked second in carbohydrate content.



Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

**Fig. 3.** Leaf dry matter % and carbohydrate concentration of strawberry cultivars Fortuna and Festival treated with some stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

**Table 3.** Leaf dry matter % and carbohydrate concentration of strawberry cultivars Fortuna and Festival treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Stimulants / Cultivars	Dry matter (%)				Total Carbohydrates (g.100g <sup>-1</sup> DW)			
	Fortuna		Festival		Fortuna		Festival	
	2017-2018							
Fulvic acid	39.72±0.38	a	36.14±1.59	bc	68.72±0.91	a	66.16±0.75	ef
Humic acid	36.29±0.73	b	34.41±0.14	de	67.39±0.74	bc	65.01±0.62	i
Salicylic acid	34.81±0.39	b-d	34.42±0.11	de	61.58±1.02	l	65.56±1.06	g-i
Potassium Citrate	34.71±0.35	cd	33.71±0.17	d-f	65.31±0.53	hi	67.67±0.79	b
Potassium Silicate	34.63±0.32	d	32.53±0.15	fg	63.50±0.77	k	64.17±0.73	j
Calcium Carbonate	33.12±0.19	e-g	32.12±0.80	g	66.57±0.70	de	67.07±0.56	cd
Water (Control)	40.04±0.25	a	33.46±0.17	d-g	65.59±0.69	gh	65.95±0.75	fg
Mean	36.18	a	33.82	b	65.52	b	65.94	a
	2018-2019							
Fulvic acid	40.07±0.18	a	36.04±1.60	bc	71.60±0.70	a	69.31±0.51	cd
Humic acid	36.75±0.61	b	34.32±0.11	de	70.48±0.50	b	67.60±0.42	e
Salicylic acid	35.07±0.27	cd	34.35±0.09	de	65.85±0.69	f	67.33±0.60	e
Potassium Citrate	34.48±0.27	de	33.82±0.12	de	67.52±0.36	e	70.99±0.54	ab
Potassium Silicate	34.42±0.24	de	32.44±0.11	fg	66.73±0.52	e	67.21±0.49	e
Calcium Carbonate	33.25±0.14	ef	31.59±0.61	g	69.50±0.47	c	69.42±0.38	c
Water (Control)	40.21±0.19	a	33.35±0.14	ef	68.48±0.47	d	69.08±0.51	cd
Mean	36.32	a	33.70	b	68.59	a	68.70	a

<sup>a</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Mean value  $\pm$  standard error.

Cultivars exhibited a significant effect on dry matter in both years, but carbohydrates in the first season only.

Our results agree with those of Abou El Hassan and Husein (2016) and Suh et al., (2014) who found that foliar

spray with fulvic acid increased plant dry weight in tomato plants. Aminifard et al., (2012) reported positive enhancement in total carbohydrate concentration in pepper due to fulvic acid treatment. Concerning humic acid,

Chakraborty et al., (2023) tested two concentrations of humic acid (1.0 and 2.0-ml L<sup>-1</sup>) on strawberry plants and found that both concentrations increased the total sugar concentration. Regarding potassium citrate, foliar application with a combination of potassium citrate 1% and boron 50 ppm on strawberry plants improved carbohydrate concentration under both mineral nitrogen and compost manure application as organic fertilizer (Awad et al., 2010). EL-Sayed et al (2023) reported a significant increase in total sugar due to foliar spraying with calcium and potassium silicate separately in strawberry plants.


**c- Non-enzymatic components**

Data in Figure 4 displayed the individual effects of bio-stimulants on strawberry plants for non-enzymatic components, which include alkaloids, flavonoids, total phenolics, and antioxidant activity. All bio-stimulants showed negative effects on alkaloid concentration in leaves in both

seasons (Figure 4A). Potassium citrate exhibited the highest significant flavonoid content followed by calcium carbonate in both seasons compared to other treatments (Figure 4B).

All bio-stimulants impaired total phenolics, except calcium carbonate and potassium citrate. Calcium carbonate showed the highest value of total phenolics, followed by potassium citrate in both seasons (Figure 4C). All bio-stimulants enhanced the antioxidant activity, except calcium carbonate, which caused a significant reduction. Humic acid recorded the highest significant antioxidant activity, followed by fulvic acid (Figure 4D).

Concerning the interaction between cultivars and bio-stimulants, Table 4 showed that significant differences were found among the effects of the bio-stimulants in the two cultivars for non-enzymatic components.



Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

**Fig. 4. Non-enzymatic antioxidant leaf concentration (A: total alkaloids; B: total flavonoids; C: total phenolics; and D: antioxidant activity) of two strawberry cultivars treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.**

All bio-stimulants reduced alkaloids in leaves over control plants in both cultivars, which showed the highest significant value of alkaloids. Regarding flavonoids, foliar application of Festival plants with potassium citrate in both seasons, calcium carbonate of Festival plants in the first season displayed the highest flavonoids content, meanwhile

spraying Fortuna plants with potassium citrate in both seasons and calcium carbonate of Festival plants in the second season ranked second in flavonoids concentration.

Concerning total phenolics, all bio-stimulants had a negative effect on total phenolic concentration, except calcium carbonate and potassium citrate. Spraying the

Fortuna cultivar with calcium carbonate recorded the maximum value, while spraying the festival cultivar with calcium carbonate ranked second in both seasons, in addition to potassium citrate in festival cultivar in the first season and in Fortuna in the second season. Concerning antioxidant activity, all bio-stimulants enhanced antioxidant activity,

except potassium citrate and calcium carbonate which are equal to control treatments in both cultivars throughout both seasons. Humic acid foliar application on Fortuna cultivar demonstrated the highest antioxidant activity in both seasons, meanwhile, humic acid application in festival cultivar ranked second in both seasons.

**Table 4. Leaf concentration of non-enzymatic components (alkaloids, flavonoids, total phenolics, and antioxidant activity) of strawberry cultivars Fortuna and Festival treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.**

Stimulants / Cultivars	Total Alkaloids (mg HE.g <sup>-1</sup> DW)		Total flavonoids (mg Catechin.g <sup>-1</sup> DW)		Total phenolics (mg GAE.g <sup>-1</sup> DW)		Antioxidant activity DPPH IC <sub>50</sub> (µg/g Extract)	
	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival
2017-2018								
Fulvic acid	1.00±0.03 e	0.99±0.03 e	2.41±0.07 f	2.81±0.11 de	6.52±0.19 f	6.47±0.16 fg	3.35±0.03 c	3.39±0.04 c
Humic acid	1.21±0.08 e	1.23±0.05 e	1.70±0.05 h	1.84±0.06 gh	4.64±0.07 i	5.09±0.15 h	4.85±0.03 a	4.63±0.05 b
Salicylic acid	2.44±0.17 b	2.42±0.15 b	3.00±0.07 c	2.80±0.08 de	6.17±0.15 g	6.65±0.22 f	2.40±0.09 e	2.26±0.10 e
Potassium Citrate	2.01±0.05 c	1.93±0.11 c	3.76±0.16 b	4.13±0.26 a	8.48±0.05 c	9.13±0.29 b	1.84±0.06 f	1.05±0.11 g
Potassium Silicate	1.50±0.18 d	1.54±0.19 d	2.51±0.21 f	2.72±0.08 e	7.88±0.20 d	7.15±0.10 e	2.99±0.08 d	3.28±0.12 c
Calcium Carbonate	1.99±0.08 c	2.03±0.09 c	3.01±0.14 c	4.06±0.09 a	9.76±0.19 a	8.98±0.20 b	0.90±0.09 h	0.79±0.10 h
Water (Control)	2.87±0.10 a	2.95±0.08 a	1.96±0.16 g	2.96±0.06 cd	7.87±0.20 d	8.50±0.14 c	1.07±0.11 g	1.13±0.09 g
Mean	1.860	1.867	2.62	3.04	7.33	7.42	2.48	2.36
2018-2019								
Fulvic acid	1.16±0.02 g	1.16±0.03 g	2.33±0.25 ef	3.23±0.05 cd	6.94±0.18 d	5.68±0.15 e	3.49±0.02 cd	3.55±0.03 c
Humic acid	1.44±0.02 f	1.31±0.11 fg	1.78±0.11 g	2.08±0.03 fg	4.41±0.11 f	4.38±0.13 f	4.99±0.02 a	4.61±0.20 b
Salicylic acid	3.02±0.06 b	2.60±0.20 c	3.02±0.22 d	3.11±0.04 d	5.46±0.13 e	5.59±0.20 e	2.78±0.06 e	2.44±0.15 f
Potassium Citrate	2.30±0.04 d	2.37±0.15 d	4.12±0.18 b	5.09±0.12 a	8.23±0.05 b	7.74±0.26 c	2.09±0.04 g	1.49±0.18 h
Potassium Silicate	2.06±0.06 e	2.07±0.06 e	3.05±0.16 d	3.03±0.04 d	6.90±0.18 d	6.68±0.09 d	3.34±0.06 d	3.56±0.19 c
Calcium Carbonate	2.40±0.06 d	2.38±0.14 d	3.54±0.07 c	4.38±0.04 b	8.86±0.17 a	8.01±0.18 bc	1.27±0.06 ij	1.15±0.08 j
Water (Control)	3.32±0.07 a	3.25±0.10 a	2.57±0.08 e	3.18±0.03 d	6.90±0.18 d	7.81±0.13 c	1.51±0.07 h	1.43±0.15 hi
Mean	2.242	2.161	2.91	3.44	6.81	6.55	2.78	2.60

<sup>a</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test (p < 0.05). Mean value ± standard error.

Cultivars displayed a significant effect on flavonoids in both seasons and on total phenolics in the second season only, but they had a non-significant effect on alkaloids and antioxidant activity in both seasons.

Non-enzymatic components are produced by plants to defend against different stresses. In former studies, El-Sayed et al., (2023) revealed that foliar spraying with salicylic acid reduced the non-enzymatic components in tomato leaves such as total sugars and total phenolics. Aminifard et al., (2012) found that fulvic acid treatment increased total phenol and antioxidant activity while having no effects on total flavonoids in pepper plants. El-Sayed et al., (2023) found that calcium and potassium silicate didn't affect the total phenol concentration, while increased antioxidant capacity in strawberry plants. Aghaeifard et al., (2016) found a positive effect of salicylic acid on strawberry plants, that increased antioxidant capacity, while humic acid didn't affect this trait.

**Effect of bio-stimulants, cultivars, and their interaction on yield and its quality**

Results in Figure 5 revealed the individual effects of bio-stimulants which displayed significant differences between bio-stimulants for early yield, total yield, average fruit weight, and fruit firmness. Potassium silicate in both seasons showed the highest significant early yield with no significant differences with calcium carbonate in the first season, and salicylic acid in the second season (Figure 5A). Fulvic acid showed the lowest early yield in the first season compared to all other treatments and was equal to the control treatment in the second season. Calcium carbonate in both seasons, potassium silicate in the first season, and potassium citrate in the second season recorded the highest total yield, while fulvic acid produced the lowest total yield in both seasons (Figure 5B). Fulvic and humic acid in the first season and all bio-stimulants in the second season caused a significant reduction in average fruit weight (Figure 5C). Regarding firmness, potassium citrate, potassium silicate, and

calcium carbonate showed the highest values in both seasons with no significant difference (Figure 5D).


Regarding the interaction between cultivars and bio-stimulants, Table 5 showed significant differences were found among the bio-stimulants in the two cultivars for early yield, total yield, average fruit weight and fruit firmness. Concerning the early yield, spraying the festival cultivar with potassium silicate in both seasons, calcium carbonate in the first season, and salicylic acid in the second season showed the highest early yield. Humic acid and potassium citrate ranked second in this trait in festival cultivar in the second season. Fulvic acid caused a significant reduction in early yield in both cultivars in the first year, while other bio-stimulants did not differ from plants of control.

Regarding total yield, calcium carbonate in both cultivars during the two seasons, potassium silicate in both cultivars in the first season, and humic acid in festival cultivar produced the highest total yield. Fulvic acid reduced total yield in both cultivars in both seasons. Other treatments did not differ from plants of the control.

Concerning the average fruit weight, fulvic acid in both cultivars and humic acid in Fortuna caused a significant reduction in this trait, while other treatments did not differ from plants of control in both cultivars in the first season. Fulvic acid and humic acid in both cultivars, in addition to salicylic acid and potassium citrate in the Fortuna caused a significant reduction in average fruit weight in the second season, while other treatments did not differ from plants of the control in both seasons. Concerning fruit firmness, all treatments except fulvic acid and salicylic acid in both cultivars, in addition to humic acid in Festival cultivar improved fruit firmness in the first season. Also, all treatments, except salicylic acid in both cultivars, in addition to humic and fulvic acid in Festival cultivar enhanced fruit firmness in the second season.



Cultivars displayed a significant effect on fruit weight in the second season only, but they had a non-significant effect on total yield in both years.



Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

Fig. 5. Yield component (A: early yield and B: total yields  $\text{kg/m}^2$ ) and physical fruit quality parameters (C: average fruit weight and D: fruit firmness) of two strawberry cultivars treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Table 5. Yield component and physical fruit quality parameters of strawberry cultivars Fortuna and Festival treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

Cultivars (C)	Early yield( $\text{kg/m}^2$ )		Total yield( $\text{kg/m}^2$ )		Average fruit weight(g)		Fruit firmness( $\text{kg/cm}^2$ )	
	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival
Stimulants (S)								
	2017-2018							
Fulvic acid	0.39 $\pm$ 0.06 e	0.67 $\pm$ 0.02 e	0.91 $\pm$ 0.13 d	1.32 $\pm$ 0.04 d	14.64 $\pm$ 0.60 e	15.45 $\pm$ 1.02 e	2.35 $\pm$ 0.02 b-e	2.14 $\pm$ 0.04 e
Humic acid	1.34 $\pm$ 0.14 b-d	1.42 $\pm$ 0.05 a-d	3.25 $\pm$ 0.33 bc	3.36 $\pm$ 0.13 ab	18.92 $\pm$ 0.48 d	23.03 $\pm$ 0.73 a-c	2.43 $\pm$ 0.06 a-c	2.22 $\pm$ 0.13 c-e
Salicylic acid	1.44 $\pm$ 0.12 a-d	1.39 $\pm$ 0.07 a-d	3.25 $\pm$ 0.35 bc	3.19 $\pm$ 0.15 bc	22.74 $\pm$ 1.07 bc	25.86 $\pm$ 0.28 a	2.19 $\pm$ 0.06 de	2.16 $\pm$ 0.09 e
Potassium Citrate	1.37 $\pm$ 0.09 b-d	1.29 $\pm$ 0.05 cd	3.18 $\pm$ 0.20 bc	3.07 $\pm$ 0.09 bc	21.30 $\pm$ 0.43 cd	24.48 $\pm$ 0.53 ab	2.53 $\pm$ 0.06 ab	2.41 $\pm$ 0.11 a-d
Potassium Silicate	1.67 $\pm$ 0.08 ab	1.59 $\pm$ 0.05 a-c	3.84 $\pm$ 0.19 ab	3.71 $\pm$ 0.14 ab	24.88 $\pm$ 2.14 ab	22.66 $\pm$ 1.57 bc	2.45 $\pm$ 0.03 a-c	2.54 $\pm$ 0.07 ab
Calcium Carbonate	1.73 $\pm$ 0.05 a	1.72 $\pm$ 0.13 a	4.14 $\pm$ 0.10 a	4.12 $\pm$ 0.28 a	24.69 $\pm$ 0.88 ab	24.02 $\pm$ 1.15 a-c	2.62 $\pm$ 0.13 a	2.40 $\pm$ 0.06 a-d
Water (Control)	1.66 $\pm$ 0.32 ab	1.22 $\pm$ 0.07 d	3.82 $\pm$ 0.75 ab	2.45 $\pm$ 0.15 c	23.54 $\pm$ 1.78 a-c	23.75 $\pm$ 0.23 a-c	2.16 $\pm$ 0.08 e	2.15 $\pm$ 0.06 e
Mean	1.37 a	1.32 a	3.19 a	3.02 a	21.53 a	22.74 a	2.38 a	2.28 b
	2018-2019							
Fulvic acid	0.28 $\pm$ 0.05 f	0.83 $\pm$ 0.17 de	1.13 $\pm$ 0.08 g	1.86 $\pm$ 0.27 f	13.24 $\pm$ 0.09 f	14.25 $\pm$ 0.74 f	2.08 $\pm$ 0.02 ab	1.86 $\pm$ 0.03 d
Humic acid	0.58 $\pm$ 0.05 ef	1.56 $\pm$ 0.07 b	3.69 $\pm$ 0.01 a-c	3.27 $\pm$ 0.01 c-e	19.52 $\pm$ 0.01 e	22.57 $\pm$ 0.44 d	2.10 $\pm$ 0.01 ab	1.84 $\pm$ 0.05 d
Salicylic acid	0.62 $\pm$ 0.19 e	2.39 $\pm$ 0.19 a	2.82 $\pm$ 0.28 e	3.28 $\pm$ 0.14 c-e	21.64 $\pm$ 0.49 d	25.43 $\pm$ 0.09 b	1.84 $\pm$ 0.04 d	1.81 $\pm$ 0.06 d
Potassium Citrate	1.25 $\pm$ 0.06 bc	1.33 $\pm$ 0.17 b	3.81 $\pm$ 0.26 a-c	3.73 $\pm$ 0.23 a-c	21.89 $\pm$ 0.14 d	24.79 $\pm$ 0.53 bc	2.20 $\pm$ 0.01 a	2.01 $\pm$ 0.06 bc
Potassium Silicate	0.99 $\pm$ 0.07 cd	2.16 $\pm$ 0.16 a	3.47 $\pm$ 0.41 b-d	3.29 $\pm$ 0.19 c-e	25.33 $\pm$ 1.96 b	24.96 $\pm$ 0.74 bc	2.11 $\pm$ 0.01 ab	2.14 $\pm$ 0.03 a
Calcium Carbonate	0.78 $\pm$ 0.13 de	0.84 $\pm$ 0.06 de	4.01 $\pm$ 0.09 ab	4.28 $\pm$ 0.24 a	23.30 $\pm$ 0.12 cd	24.85 $\pm$ 0.58 bc	2.16 $\pm$ 0.07 a	2.09 $\pm$ 0.04 ab
Water (Control)	0.74 $\pm$ 0.19 de	0.59 $\pm$ 0.01 ef	3.22 $\pm$ 0.44 c-e	2.84 $\pm$ 0.08 de	25.04 $\pm$ 0.61 bc	30.06 $\pm$ 0.13 a	1.89 $\pm$ 0.06 cd	1.88 $\pm$ 0.02 d
Mean	0.74 b	1.38 a	3.16 a	3.22 a	21.42 b	23.84 a	2.05 a	1.94 b

<sup>a</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Mean value  $\pm$  standard error

Previous works, in agreement with our results, found that foliar spray with calcium and potassium silicate enhanced average fruit weight, fruit firmness, marketability, and total yield in strawberry (EL-Sayed et al., 2023). Nada (2020) reported a significant increment in fruit weight, fruit firmness, and early and total yield in strawberry due to foliar spraying with potassium silicate. Rozbiany and Taha (2020) found that fruit fresh weight and yield/plant were significantly improved due to foliar Spraying with 500 ppm of calcium in strawberry cv. Festival. Also, there was a positive impact on early and total yield in tomatoes after foliar treatment with salicylic acid and potassium silicate (EL-Sayed et al., 2023). Youssef et al., (2017) tested the effect of eight concentrations of salicylic acid and found that salicylic acid at 4.0 mM produced the biggest average fruit weight and the highest early and total yield.

**Effect of bio-stimulants, cultivars, and their interaction on fruit quality traits**


Data in Figure 6 exhibited that significant differences were found between bio-stimulants on strawberry plants for TSS, titratable acidity, ascorbic acid and anthocyanin concentration. Concerning TSS, potassium citrate, potassium silicate, and calcium carbonate demonstrated the highest content of TSS in both seasons. Concerning titratable acidity, humic acid, salicylic acid, and potassium silicate showed the highest value of titratable acidity in fruits in both seasons, while potassium citrate and calcium carbonate decreased the titratable acidity value. Concerning ascorbic acid, no significant differences were found

among bio-stimulants in the first season. Meanwhile, potassium citrate, potassium silicate, and calcium carbonate displayed the highest ascorbic acid in fruits in the second season. Regarding anthocyanin concentration, calcium carbonate, potassium citrate and potassium silicate in both seasons exhibited the highest significant value.

Concerning the interaction between cultivars and bio-stimulants, Table 6 displayed that significant differences were found among the effects of the bio-stimulants in the two cultivars for titratable acidity, TSS, anthocyanin in both seasons and ascorbic acid in second season only.

Potassium citrate, potassium silicate in both cultivars and calcium carbonate in Fortuna cultivar showed the highest TSS content in first season, while all bio-stimulants enhanced the TSS content in Fortuna cultivar, except salicylic acid, while potassium silicate and calcium carbonate in both seasons showed the highest TSS in Festival cultivar, and potassium citrate ranked second in TSS in Festival cultivar in the second season. Concerning titratable acidity, humic acid in both cultivars in addition to salicylic acid and potassium silicate in Festival cultivar during both seasons showed the highest value of titratable acidity in fruits.

Concerning ascorbic acid in second season, potassium silicate and calcium carbonate in both cultivars, and humic acid and potassium citrate in Fortuna cultivar exhibited the highest ascorbic acid concentration in fruits during the second season.



Columns with the same letter represent values that are not significantly different according to Duncan's multiple range test ( $p < 0.05$ ). Vertical bars represent  $\pm$  standard error of the mean.

**Fig. 6.** Fruit chemical composition (A: TSS; B: titratable acidity; C: ascorbic acid; and D: anthocyanin) of two strawberry cultivars treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.

**Table 6. Fruit chemical composition of strawberry cultivars Fortuna and Festival treated with some bio-stimulants in greenhouse conditions during the 2017-2018 and 2018-2019 winter seasons.**

Cultivars (C)	TSS (°Brix)		Titratable acidity (mg citric acid. 100g <sup>-1</sup> FW)		Ascorbic acid (mg. g <sup>-1</sup> FW)		Anthocyanin (mg. 100g <sup>-1</sup> FW)	
	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival	Fortuna	Festival
2017-2018								
Fulvic acid	6.24±0.21 b-e	5.89±0.23 e	0.45±0.027 ef	0.48±0.022 de	0.65±0.047 ab	0.66±0.011 ab	56.77±0.37 c-e	52.21±0.83 ef
Humic acid	6.70±0.17 ab	6.11±0.36 b-e	0.56±0.012 ab	0.57±0.024 ab	0.68±0.045 ab	0.63±0.024 ab	57.19±0.46 b-e	53.49±3.17 d-f
Salicylic acid	5.78±0.31 e	5.95±0.24 de	0.50±0.005 cd	0.55±0.007 ab	0.67±0.020 ab	0.64±0.017 ab	52.79±1.55 ef	50.46±1.31 f
Potassium Citrate	6.98±0.16 a	6.65±0.31 a-c	0.39±0.014 h-j	0.42±0.010 f-h	0.71±0.051 ab	0.67±0.048 ab	60.08±0.95 a-c	58.19±2.72 b-d
Potassium Silicate	6.74±0.08 ab	7.01±0.18 a	0.45±0.005 e-g	0.59±0.013 a	0.65±0.056 ab	0.72±0.061 ab	58.99±0.70 a-c	61.98±1.49 ab
Calcium Carbonate	7.23±0.35 a	6.63±0.17 a-d	0.35±0.016 j	0.40±0.016 g-i	0.71±0.072 ab	0.74±0.019 a	63.88±2.98 a	57.97±1.47 b-d
Water (Control)	5.97±0.23 c-e	5.94±0.16 de	0.36±0.007 ij	0.53±0.005 bc	0.62±0.033 b	0.65±0.014 ab	52.21±2.04 ef	51.04±1.07 f
Mean	6.52 a	6.31 a	0.43 b	0.50 a	0.66 a	0.67 a	57.41 a	55.04 b
2018-2019								
Fulvic acid	6.67±0.07 ab	5.95±0.10 c	0.45±0.043 d-f	0.49±0.049 c-e	0.74±0.008 bc	0.66±0.011 e	58.14±0.62 ab	52.06±0.89 d
Humic acid	6.73±0.03 ab	5.92±0.17 c	0.55±0.018 a-c	0.63±0.018 a	0.75±0.003 a-c	0.66±0.019 e	58.92±0.25 ab	51.32±1.56 d
Salicylic acid	5.93±0.13 c	5.86±0.19 c	0.53±0.035 b-d	0.61±0.031 ab	0.66±0.015 e	0.65±0.021 e	51.49±1.21 d	51.26±1.64 d
Potassium Citrate	7.05±0.03 a	6.47±0.19 b	0.39±0.019 fg	0.42±0.033 e-g	0.79±0.04 a	0.72±0.021 cd	61.69±0.30 a	56.07±1.74 bc
Potassium Silicate	6.78±0.03 ab	6.91±0.10 a	0.44±0.009 d-f	0.59±0.018 ab	0.76±0.004 a-c	0.77±0.011 ab	59.44±0.31 ab	60.45±0.86 a
Calcium Carbonate	7.02±0.22 a	6.72±0.14 ab	0.35±0.043 g	0.40±0.023 e-g	0.78±0.025 ab	0.75±0.016 a-c	61.40±1.96 a	59.15±1.31 ab
Water (Control)	6.07±0.19 c	6.02±0.05 c	0.36±0.019 fg	0.53±0.013 b-d	0.68±0.021 de	0.67±0.005 e	53.60±1.74 cd	52.78±0.44 cd
Mean	6.60 a	6.26 b	0.43 b	0.52 a	0.73 A	0.69 b	57.81 a	54.72 b

<sup>2</sup>Mean values followed by a letter in common were not significantly different according to Duncan's multiple range test (p < 0.05). Mean value ± standard error.

Regarding anthocyanin concentration in fruits, potassium Silicate in the two cultivars, in addition to potassium citrate and calcium carbonate in Fortuna cultivar exhibited the highest anthocyanin concentration in the first season, while all bio-stimulants increased the content of anthocyanin in Fortuna, except salicylic acid in the second season, but in festival cultivar, potassium silicate, and calcium carbonate displayed the highest concentration of anthocyanin.

Cultivars displayed a significant effect on titratable acidity and anthocyanin concentration in the both seasons and TSS and ascorbic acid in the second season.

Similar findings by Al-Qadi and Alimam (2023) evaluated three concentrations of humic acid (0, 3, and 6 ml L<sup>-1</sup>) and found that 3 ml L<sup>-1</sup> increased total acidity content in strawberry fruits. Kazemi et al. (2014) assessed the effect of two concentrations of humic acid (15 and 25 mL L<sup>-1</sup>) on fruit quality and found that both levels of humic significantly increased TSS, vitamin C, and acidity %. Rozbiany and Taha (2020) reported that foliar spraying with calcium significantly increased TSS, vitamin C, and acidity %. EL-Sayed et al. (2023) tested the effect of foliar spray with potassium silicate and calcium on the chemical fruit quality traits of strawberry fruits. Calcium significantly improved anthocyanin concentration, while potassium silicate had no significant effect on this trait.

The highest concentration of vitamin C was achieved using 0.6% potassium silicate, also 1% and 2% of calcium treatment increased this content. Calcium treatment and 0.6 % potassium silicate application increased acidity %, but TSS was not influenced in both seasons. Nada (2020) tested four concentrations of potassium silicate (0, 0.2, 0.4, and 0.6 g L<sup>-1</sup>) on strawberry plants, and found that all concentrations caused significant increments in TSS, acidity%, vitamin C, and anthocyanin concentration in fruits. In contrast, the interaction effects of foliar application of humic acid and two cultivars (Camarosa and Nabila) on quality parameters. TSS, titratable acidity, and vitamin C were found non-significant (Chakraborty et al., 2023), also Ullah et al. (2017) tested four levels of humic acid (0, 1.5, 3, and 4.5 ml L<sup>-1</sup>) on strawberry plants and found that humic acid showed a non-significant effect on the titratable acidity of fruit (Ullah et al., 2017).

## CONCLUSION

The impact of different bio-stimulants i.e., humic acid, fulvic acid, salicylic acid, potassium citrate, potassium silicate, and calcium carbonate on plant vigor, yield, and quality attributes in two strawberry cultivars (Fortuna and Festival) in a plastic greenhouse were studied. Calcium carbonate and potassium silicate enhanced all most studied traits except dry matter, alkaloids in both stimulants, antioxidant activity, and titratable acidity in calcium carbonate, and their effects were different among the cultivars.

## REFERENCES

- Abd-El-Kareem, F., Elshahawy, I. E., and Abd-Elgawad, M. M. (2019). Management of strawberry leaf blight disease caused by *Phomopsis obscurans* using silicate salts under field conditions. *Bulletin of the National Research Centre*, 43(1), 1-6.
- Abdelwines, M. A., and Ahmed, M. M. (2022). The effect of some fertilizer compounds as a resistance inducer in strawberry plants on life history parameters of *Tetranychus urticae* (Acari: Tetranychidae). *Persian Journal of Acarology*, 11(2), 275-293.
- Abdusameea, M. S. I., Amira, R. M., Hassanb, I., MudasarAlia, Z. Z., Khana, A. U. and Nasira, H. (2023). A Comprehensive Review on the Health-Orientated Aspects of Strawberries. *Food Science & Applied Microbiology Reports*, 2(1), 1-7.
- Abou El Hassan, S. and Husein, M. E. (2016). Response of tomato plants to foliar application of humic, fulvic acid and chelated calcium. *Egyptian Journal of Soil Science*, 56(3), 401-141.
- Aghaeifard, F., Babalar, M., Fallahi, E., and Ahmadi, A. (2016). Influence of humic acid and salicylic acid on yield, fruit quality, and leaf mineral elements of strawberry (*Fragaria × Ananassa* duch.) cv. Camarosa. *Journal of Plant Nutrition*, 39(13), 1821-1829.

- Al-Qadi, R. A. and Alimam, N. M. (2023). Response of three varieties of strawberry to the nano-NPK fertilizer, humic acid. *International Journal of Agricultural & Statistical Sciences*, 19(1),89-99.
- Aminifard, M. H., Aroiee, H., Nemati, H., Azizi, M. and Jaafar, H. Z. (2012). Fulvic acid affects pepper antioxidant activity and fruit quality. *African Journal of Biotechnology*, 11(68), 13179-13185.
- AOAC. (1990). Association of Official Agricultural Chemists. *Official Methods of Analysis*. 15<sup>th</sup> ed, Washington. D.C., USA., 1298 p.
- Awad, E. M., Mohamed, R. A. and Asfour, H. E. (2010). Effect of compost, foliar spraying with potassium and boron on growth, yield and fruit quality of strawberry. *Journal of Plant Production*, 1(8), 1101-1112.
- Chakraborty, B., Basak, S., Sherpa, Z. W., Samanta, D., Apoorva, N. D. and Gurung, S. (2023). Effect of foliar application of humic acid and seaweed extract in strawberry (*Fragaria x ananassa* Duch.). *Plant Archives*, 23(2),68-72.
- Chandler, C.K., Folta, K., Dale, A., Whitaker, V.M., Herrington, M. (2012). Strawberry. In: Badenes, M., Byrne, D. (eds) *Fruit Breeding. Handbook of Plant Breeding*, vol 8. Springer, Boston, MA. [https://doi.org/10.1007/978-1-4419-0763-9\\_9](https://doi.org/10.1007/978-1-4419-0763-9_9)
- Chang, C. C., Yang, M. H., Wen, H. M. and Chen, J. C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of food and drug analysis*, 10(3), 178–182.
- Coskun, D., Britto, D. T. and Kronzucker, H. J. (2017). The nitrogen–potassium intersection: membranes, metabolism, and mechanism. *Plant, cell & environment*, 40(10), 2029-2041.
- Dakheel, M. G., Ullah, I., Doğan, D. E. and Demirsoy, L. (2022). Potential role of salicylic acid on drought stress tolerance of strawberry plants. In *International Symposium on Soil Science and Plant Nutrition*, 269-275.
- De Resende, J. T. V., Matos, R., Zeffa, D. M., Constantino, L. V., Alves, S. M., Ventura, M. U. and Youssef, K. (2021). Relationship between salicylic acid and resistance to mite in strawberry. *Folia Horticulturae*, 33(1), 107-119.
- De Vasconcelos, A. C. F. and Chaves, L. H. G. (2019). Biostimulants and their role in improving plant growth under abiotic stresses. *Biostimulants in plant science*, 3-16.
- Dehghanipoozeh, S., Ghobadi, C., Baninasab, B., Gheysari, M. and Shiranibidabadi, S. (2018). Effect of silicon on growth and development of strawberry under water deficit conditions. *Horticultural Plant Journal*,4(6),226-232.
- DuBois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. T. and Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analyticalchemistry*, 28(3),350-356.
- El-Sayed, H.A., Eata, A. E. M. and Khater, M. A. K. (2019). Response of early sowing pea crop to some natural and chemical substances for improving yield and quality and components. *Journal of Plant Production*, 10 (10), 849 – 858.
- El-Sayed, S. F., Mohamed, M. I. and Hanafy, A. M. (2023). Comparison among different amelioratives on improvement of tomato growth, yield and quality under low temperature condition. *Eur. Chem. Bull*, 12(Special Issue 1), 2703-2730.
- EL-Sayed, S. S. F., EL-Mogy, M. M. and Mahmoud, G. A. (2023). Potassium silicate and calcium chloride production and shelf-life of strawberries. *Fayoum Journal of Agricultural Research and Development*, 37(4), 736-759.
- Emam, S. S., Marzouk, A. S. and Abdelghany, R. A. (2023). Effect of pre-harvest application of salicylic acid on strawberry growth and resistance against gray mold. *International Journal of Scientific Research and Sustainable Development*, 6 (1): 37-36
- FAOSTAT, (2024). *Food and Agricultural Commodities Production*; online: <http://faostat.fao.org> (Accessed 8 march 2024).
- Garza-Alonso, C. A., Olivares-Sáenz, E., González-Morales, S., Cabrera-De la Fuente, M., Juárez-Maldonado, A., González-Fuentes, J. A., Tortella, G.; Valdés-Caballero, M.V. and Benavides-Mendoza, A. (2022). Strawberry biostimulation: From mechanisms of action to plant growth and fruit quality. *Plants*, 11(24), 3463.
- Ghoname, A. A., Dawood, M. G., Riad, G. S. and El-Tohamy, W. A. (2009). Effect of nitrogen forms and biostimulants foliar application on the growth, yield and chemical composition of hot pepper grown under sandy soil conditions. *Research Journal of Agriculture and Biological Sciences*, 5(5), 840-852.
- Gomez, A. K. and Gomez, A. A. (1984). *Statistical Procedures for Agricultural Research*. 2<sup>nd</sup> ed. John Wiley & Sons Pub., pp.139-153.
- Hassoon, A. S. and Abduljabbar, I. A. (2020). Review on the role of salicylic acid in plants. *Sustainable crop production*, 61-64.
- Hussain, Z. Z., Al-Dahwi, S. S. J. and Lelo, H. M. (2023). Effect of foliar fertilization with potassium silicate on some morphological characters of strawberry leaves reducing sucking insects. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1225, No. 1, p. 012085). IOP Publishing.
- Kazemi, M. (2014). The impact of foliar humic acid sprays on reproductive biology and fruit quality of strawberry. *Thai Journal of Agricultural Science*, 47(4), 221-225.
- Kedare, S.B. and Singh, R. (2011). Genesis and development of DPPH method of antioxidant assay. *Journal of food science and technology*, 48(4), 412-422.
- Mansour F. Y.O. and El- Metwaly, H.M.B. (2019). Influence of addition methods of fulvic acid and foliar spray with potassium sources on productivity and quality of potato crop. *Fayoum J. Agric. Res. & Dev.*,33(1):130-147.
- Meda, A., Lamien, C. E., Romito, M., Millogo, J. and Nacoulma, O. G. (2005). Determination of the total phenolic, flavonoid and proline contents in Burkina Fasan honey, as well as their radical scavenging activity. *Food chemistry*, 91(3), 571-577.
- Mitic, V., Jovanovic, V. S., Dimitrijevic, M., Cvetkovic, J. and Stojanovic, G. (2013). Effect of food preparation technique on antioxidant activity and plant pigment content in some vegetables species. *Journal of Food and Nutrition Research*, 1(6), 121-127.
- Mohamed, R.A., Al-Kharpotly, A., and Abd-Elkader, D.Y. (2017). Salicylic acid effects on growth, yield, and fruit quality of strawberry cultivars. *Journal of Medicinally Active Plants*, 6 (2),1-11.

- Mousa, T. A., Bardisi, S. A., Esmail, H. E. M. and Zayd, G. A. (2023). Plant growth, yield, and tuber quality of some potato cultivars as affected by potassium sources as foliar application under sandy soil conditions. *Zagazig Journal of Agricultural Research*, 50(6), 863-880.
- Nada, M. M. (2020). Effect of foliar spray with potassium silicate and glycine betaine on growth and early yield quality of strawberry plants. *Journal of Plant Production*, 11 (12), 1295 – 1302.
- Navazio, L., Formentin, E., Cendron, L. and Szabò, I. (2020). Chloroplast calcium signaling in the spotlight. *Frontiers in plant science*, 11, 186. <https://doi.org/10.3389/fpls.2020.00186>.
- Ore, O. T., Adeola, A. O., Fapohunda, O., Adedipe, D. T., Bayode, A. A., and Adebisi, F. M. (2023). Humic substances derived from unconventional resources: extraction, properties, environmental impacts, and prospects. *Environmental Science and Pollution Research*, 30(21), 59106-59127.
- Pandey, S. K. and Singh, H. (2011). A simple, cost-effective method for leaf area estimation. *Journal of botany*, Volume 2011, Article ID 658240, 1-6. doi:10.1155/2011/658240.
- Pertot, I., Zasso, R., Amsalem, L., Baldessari, M., Angeli, G. and Elad, Y. (2008). Integrating biocontrol agents in strawberry powdery mildew control strategies in high tunnel growing systems. *Crop protection*, 27(3-5), 622-631.
- Pirie, A. and Mullins, M. G. (1976). Changes in anthocyanin and phenolics content of grapevine leaf and fruit tissues treated with sucrose, nitrate, and abscisic acid. *Plant physiology*, 58(4), 468-472.
- Roshdy, A. E. D., Alebidi, A., Almutairi, K., Al-Obeed, R. and Elsabagh, A. (2021). The effect of salicylic acid on the performances of salt stressed strawberry plants, enzymes activity, and salt tolerance index. *Agronomy*, 11(4), 775.
- Rozbiany, P. M. K. and Taha, S. M. (2020). Influence of calcium foliar application to increase growth and yield of two strawberry cultivars (*Fragaria* × *ananassa* Duch.). *Zanco Journal of Pure and Applied Sciences*, 2218-0230.
- Sardans, J. and Peñuelas, J. (2021). Potassium control of plant functions: Ecological and agricultural implications. *Plants*, 10, (2), 419.
- Shukla, A. K. (2011). Effect of foliar application of calcium and boron on growth, productivity and quality of Indian gooseberry (*Emblica officinalis*). *Indian Journal of Agricultural Sciences*, 81(7), 628.
- Soppelsa, S., Kelderer, M., Casera, C., Bassi, M., Robatscher, P., Matteazzi, A. and Andreotti, C. (2019). Foliar applications of biostimulants promote growth, yield and fruit quality of strawberry plants grown under nutrient limitation. *Agronomy*, 9 (9), 483.
- Suh, H. Y., Yoo, K. S. and Suh, S. G. (2014). Effect of foliar application of fulvic acid on plant growth and fruit quality of tomato (*Lycopersicon esculentum* L.). *Horticulture, Environment, and Biotechnology*, 55, 455-461.
- Thor, K. (2019). Calcium—nutrient and messenger. *Front. Plant Sci.* 10:440.
- Ullah, I., Sajid, M., Shah, S. T., Khan, S., Iqbal, Z., Wahid, F., Hassan, E., Shah, S.H.A. and Khan, R. (2017). Influence of humic acid on growth and yield of strawberry cv. Chandler. *Pure and Applied Biology (PAB)*, 6(4), 1171-1176.
- Yaghubi, K., Vafae, Y., Ghaderi, N. and Javadi, T. (2019). Potassium silicate improves salinity resistant and affects fruit quality in two strawberry cultivars grown under salt stress. *Communications in Soil Science and Plant Analysis*, 50(12), 1439-1451.
- Youssef, S. M. S., Abu El-Azm, N. A. I. and Abd Elhady, S. A. E. (2017). Frequent foliar sprayings of salicylic acid with elevated concentrations enhance growth, yield and fruit quality of strawberry (*Fragaria x ananassa* Duch. cv. Festival) plants. *Egyptian Journal of Horticulture*, 44(1), 61-74.
- Zakher, A. G. and Elashry, M. A. M. (2016). Physiological studies on growth and productivity of tomato grown in sandy soil under low temperature conditions. *Journal of Plant Production, Mansoura Univ.*, 7(11): 1147 – 1154.
- Zydlík, Z. and Zydlík, P. (2023). The effect of a preparation containing humic acids on the growth, yield, and quality of strawberry fruits (*Fragaria* × *ananassa* (Duchesne ex Weston) Duchesne ex Rozier). *Agronomy*, 13(7), 1872.

## زيادة النمو والمحصول والمركبات النشطة بيولوجيا في الفراولة بواسطة المنشطات الحيوية

محمد إسماعيل عبد الله محمد ومحمد كامل فتح الله الطواشي

قسم الخضار، كلية الزراعة، جامعة القاهرة، الجيزة، مصر

### الملخص

حظيت المنشطات الحيوية باهتمام عالمي نظراً لتأثيراتها المفيدة على نمو وتطور العديد من المحاصيل. أجريت التجربة خلال موسم الشتاء لتقييم ومقارنة تأثير منشطات حيوية مختلفة على قوة النمو والمحصول وصفات الجودة في صنفين من الفراولة (فورتونا وفيسنتفال). وكانت المعاملات: 1) حامض الهيوميك 15% (10 مللي/لتر)، و2) حامض الفولفيك 15% (10 مللي/لتر)، و3) سترات بوتاسيوم (2 جرام/لتر)، و4) سيليكات البوتاسيوم (2 جرام/لتر)، و5) حامض السلسيليك (1 جرام/لتر)، و6) كربونات الكالسيوم (2 جرام/لتر)، و7) كترول (ماء). كل المعاملات كانت رش على الأوراق ماعدا الهيوميك كان إضافة أرضية. وتمت معاملة نباتات الفراولة اسبوعيا حيث تمت اول معاملة بعد 15 يوم من الشتل وتم تكرارها 10 مرات. أفضل أداء كان لسيليكات البوتاسيوم وكربونات الكالسيوم حيث أن كربونات الكالسيوم حسنت كل الصفات المدروسة عدا المادة الجافة والقلويدات الكلية ونشاط مضادات الأكسدة في الأوراق بينما حسنت سيليكات البوتاسيوم كل الصفات ماعدا المادة الجافة والقلويدات الكلية في الأوراق. أظهرت سترات البوتاسيوم أعلى قيمة للفلافونويدات في الأوراق وصلابة الثمار والمادة الصلبة الذاتية الكلية. الهيوميك أظهر أعلى نشاط لمضادات الأكسدة في الأوراق والحموضة المعاكسة. الفولفيك أظهر أعلى محتوى من الكربوهيدرات في الأوراق. فيما يتعلق بالتأثير التفاعلي بين الأصناف ومنشطات النمو اختلفت في كل الصفات. فيما يتعلق بتأثير الأصناف كان هناك اختلافات جوهريه بين الأصناف في كل الصفات فيما عدا الفينولات والقلويدات الكلية ونشاط مضادات الأكسدة في الأوراق والمحصول المبكر والكلبي ومتوسط وزن الثمرة والمواد الصلبة الذاتية الكلية وفيتامين ج في الموسم الأول وماعدا عدد الأوراق/النبات والقلويدات الكلية ونشاط مضادات الأكسدة والكربوهيدرات في الأوراق والمحصول الكلي في الموسم الثاني. توفر النتائج الحالية بيانات مهمة لمنتجي الفراولة لزيادة الإنتاجية وتحسين صفات جودة الثمار بواسطة المنشطات الحيوية، وخاصة كربونات الكالسيوم وسيليكات البوتاسيوم.

الكلمات المفتاحية: الفراولة، المنشطات الحيوية، الإنتاجية، الجودة