

# Journal of Plant Production

Journal homepage & Available online at: [www.jpp.journals.ekb.eg](http://www.jpp.journals.ekb.eg)

## Reusing Agricultural Wastewater for Tomato Production using Different Sources of Organic Fertilizer under the Conditions of the Delta Region

Abd El-Basir, A. E.<sup>1</sup>; Walaa M. E. Swelam<sup>2</sup>; Nada A. Mostafa<sup>3</sup> and Amira A. Abo Elmakarm<sup>4</sup>



<sup>1</sup>Self-Pollination Vegetables Crops Research Dept., Hort. Res. Inst., Agric. Res. Center, Giza, Egypt.

<sup>2</sup>Veget. and Flori. Dept. Fac. Agric., Mans. Univ., Egypt.


<sup>3</sup>Sgric. Botany Dept. Fac. Agric., Mans. Univ., Egypt.

<sup>4</sup>Post-harvest and Handling Vegetables Dept., Hort. Res. Inst., Agric. Res. Center, Giza, Egypt.

### ABSTRACT

The availability of fresh water (both in quantity and quality) is decreasing and becoming a major concern worldwide. The agricultural sector is the largest consumer of fresh water, so there is an urgent need to improve the agricultural practices to fill the gap between both water supply and demand, which is a fundamental aspect of the main basics of sustainable agriculture. Hence, one of the possible strategies to achieve this is the reuse of agricultural drainage water, especially with the current climate change. This research focuses on reusing agricultural wastewater in tomato production by applying organic fertilizers. The impacts of applied bioshar, compost, cattle manure and vermicompost on vegetative growth, leaf chemical composition, and productivity of tomato plants irrigated with agricultural drainage water were evaluated. As a result of the experiments, all organic soil amendments significantly enhanced vegetative growth parameters including plant height, fresh, dry weights, and leaf area compared to the control plants. These treatments also significantly increased photosynthetic pigments, yield and fruit quality of tomato plants. Conversely, the activity of antioxidant enzymes and heavy metal content were significantly reduced in response to organic fertilizers. Biochar exhibited the highest values for the studied vegetative growth parameters, leaf pigments, nutrient content as well as productivity, but the lowest levels of enzymatic activity and heavy metal content. Overall, the research suggests that agricultural wastewater can be reused for tomato production, especially in area with limited fresh water availability by using organic fertilizers such as biochar, vermicompost, compost and cattle manure.

**Keywords:** Tomato – Agricultural drainage water – Biochar – Vermicompost



### INTRODUCTION

Tomato (*Lycopersicon esculentum* Mill) is popular vegetable crop rich in vitamins and has appropriate sugar content, so it is considered the most consumed fruit on the planet, whether fresh, dried, or manufactured. It is produced worldwide, grown under different soils and climate conditions and covers nearly over 6.5 million hectares (FAOSTAT, 2022). Tomato is a high water-demand crop, so it is considered as a sensitive crop to water stress (Hou *et al.* 2020). Global water demand is increasing due to the rapid industrialization, climate change, and economic activities worldwide. This has eventually led to a scarcity of fresh water supply and increase wastewater generation, which was 359 billion m<sup>3</sup> in 2020. It is predicted to reach nearly 470 billion m<sup>3</sup> by 2030 and may increase to about 574 billion m<sup>3</sup> by 2050 (Qadir *et al.* 2020). Water scarcity, exacerbated by the current climate changes seriously affects the sustainable development of agriculture, productivity, and economic growth leading to potential risks for food security (Paudel *et al.* 2018).

There is an urgent need to improve the agriculture practices and methods to achieve the balance between both water supply and demand, also at the same time improve the resilience to climate changes (Fischer *et al.* 2019) specially with agriculture expansion and water scarcity. A promising instrument to address this need is reusing agriculture wastewater. Reusing wastewater has already been used in many regions, Egypt reuses about six

Billion m<sup>3</sup> of wastewater in the Delta Nile for crops production (Hawary and Shaban, 2018). Besides reducing stress on freshwater and pollution, it is rich in nutrients which enhances soil fertility and reduces chemical fertilizer requirement, so provides farmers with economic benefits (Seyoum *et al.*, 2022). On tomato production, Kolekar *et al.* 2023 indicated that irrigation with treated wastewater increased not only soil fertility, but also improved crop yield.

Additionally, organic manure can be applied to improve wastewater quality. Organic manure, derived from animals and plants waste and can improve soil fertility by increasing organic matter content, available nutrients, and improving soil moisture availability, which in turn improves tomato quality (Gao *et al.* 2023a). Appropriate management of agricultural waste as organic fertilizer is necessary for sustainable agricultural development, which is why biochar, compost, cattle manure as well as vermicompost have been studied. Biochar, an organic fertilizer derived from the thermal decomposition of organic matter, contributes to sustainable development and mitigates pollution caused by waste incineration. Consequently, it has many environmental, economic and agricultural benefits (Obadi *et al.* 2024) and enhancing crop production, soil properties, and water use efficiency by reducing the adverse impacts of drought and salinity stresses (Faloye *et al.* 2019). Biochar addition enhanced vegetative growth traits, increase yield, and enhanced tomato fruit quality (Obadi *et al.*, 2024).

\* Corresponding author.

E-mail address: [walaamohameed@mans.edu.eg](mailto:walaamohameed@mans.edu.eg)

DOI: 10.21608/jpp.2024.336010.1415

Compost, a suitable way of waste management that converts large volumes of waste into materials, contributes to improving soil minerals and organic matter. It also has positive impacts on both the microbial and physicochemical properties of the soil (Toledo *et al.*, 2018). The combination of compost and wastewater enhanced tomato growth and productivity (Tzortzakis *et al.*, 2020). Cattle manure is produced in large quantities due to the rapid development of animal breeding on livestock farms. Un-processed cattle manure is directly applied to the soil as organic manure to improve crop yield, as it contains large amounts of nutrients (Lee *et al.* 2021). Gao *et al.*, 2023b indicated that animal manure significantly enhanced tomato growth, yield and quality.

Vermicompost, an organic fertilizer derived from earthworms, increase soil fertility, supplies essential nutrients, and improves soil structure, and enhancing beneficial soil microorganisms (Mohammed and Alkobaisy, 2024). The application of vermicompost significantly improved not only vegetative growth, but also tomato yield and quality tomato (Turan *et al.*, 2023).

The research aims to reuse agricultural drainage water and reduce its harmful effects by using organic fertilizers (biochar, compost, cattle manure, and vermicompost) on tomato production under the conditions of Delta.

## MATERIALS AND METHODS

### Experimental Layout and Treatments:

To investigate the effects of biochar, compost, cattle manure, and vermicompost manure on tomato plants performance and productivity, the study was conducted at a private vegetable farm in Sahragt Elsoghra village, Aga district, Dakahlia governorate, Egypt, during the early summer seasons of 2022 and 2023. The tomato cultivar Ellisa was transplanted on February 10<sup>th</sup> in both seasons using drip irrigation system. The experiment plot area was 72 m<sup>2</sup> (2 drip lines × 30 m long × 1.2 m wide). Tomato seedlings were transplanted on one side of the drip line, spaced 60 cm apart. Some physical and chemical properties of the experimental site are shown in Table 1. Soil preparation, pest control, fertigation were conducted according to the guideline of the Ministry of Agriculture and land reclamation throughout the growing season,. The experiment design was a randomized complete blocks design with four replicates.

**Table 1. Some physical and chemical properties of soil profile from the experimental cultivated area.**

Soil texture	Organic matter	pH	EC dsm <sup>-1</sup>	Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>++</sup>	Mg <sup>++</sup>	CO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>
Clay loam	1.22 %	7.78	0.899	3.97	0.15	12.49	11.53	1.68	4.86	0.99

### Soil Organic Amendments

The four organic soil amendment treatments were: the individual application of biochar (5 m<sup>3</sup>/fed), compost (8 ton/fed), cattle manure (8 ton/fed), vermicompost (2 ton/fed), and the control. The different applied organic manure were added to the soil and incorporated before cultivation, in conjunction with plowing, as recommended by the Egyptian Ministry of Agriculture and land reclamation. Table 2 presents some properties of the applied organic manure. All plants were irrigated with agricultural drainage water under drip irrigation system. The characteristics of agricultural drainage water used in this study presented in Table 3.

**Table 2. Some properties of biochar, compost, cattle manure, and vermicompost applied to the soil.**

Properties	Units	Biochar	Compost	Cattle manure	Vermicompost
pH	---	8.82	7.5	-	6.0
EC	(dSm <sup>-1</sup> )	1.71	3.5	-	6.0
O.M		30.33	75	20.3	34.5
C	(%)	60	33.11	12.2	18
N		0.24	1.82	1.22	1.5
C:N ratio	---	250:1	18:1	10:1	12:1
P		0.22	1.29	0.3	0.75
K	(%)	0.88	1.25	2.9	5.05
Moisture		3.53	25	17.6	-
Mn	ppm	-	1110	446	18
Zn	ppm	-	280	180	20

**Table 3. The characteristics of agricultural drainage water used in this study**

Parameter	Unit	Drainage water
pH	---	7.85
EC	dS/m	2.43
Na <sup>+</sup>		154.1
K <sup>+</sup>		9.89
Ca <sup>++</sup>		22.5
Mg <sup>++</sup>	mg L <sup>-1</sup>	48.9
CO <sub>3</sub> <sup>-</sup>		0.04
HCO <sub>3</sub> <sup>-</sup>		6.79
Cl <sup>-</sup>		2.9
Fe <sup>++</sup>		0.451
Zn <sup>++</sup>		0.388
Mn <sup>++</sup>		0.389
Cd <sup>++</sup>	µg L <sup>-1</sup>	0.049
Cu <sup>++</sup>		0.71
Ni <sup>++</sup>		0.589
Pb <sup>++</sup>		0.434

### Parameters:

Morphological, chemical and productivity parameters were tested to assess the impacts of drainage water on plant performance in response to the different applied organic sources. Morphological and chemical data were recorded two months after transplanting. Five plants from each plot were randomly chosen to determine the following:

**Vegetative growth Characteristics included:** plant height, total fresh and dry weight and leaf area as described by Koller (1972).

**photosynthetic pigments content of leaves:** Chlorophyll a, b, Total chlorophylls (a+b) as well as carotenoids according to the methods described by Wettstein (1957).

**Mineral content of Leaves:** Samples of 100 grams were taken from the leaves two months after transplanting, dried in an oven at 70 °C for three days, and weighed to determine the concentrations N, P, K and Ca as described by Kalra (1998).

**Enzymes activity in leaves:** Peroxidase, Superoxide dismutase, Catalase activity according to Cao *et al.* (2005)

**Yield parameters: (early yield, total yield, and marketable yield):** Fruits were harvested starting three months after transplanting. Early yield calculated from fruit fresh weight of the first two pickings and calculated as ton/fed. Concerning marketable yield, the weight of usually healthy fruits (which are not infected by any disease and free from physiological defects).

**Fruit Quality:** Samples of 10 uniform, fully red-ripe fruits were sampled at the 3<sup>rd</sup> picking from each plot to estimate the total soluble solids, vitamin C, and acidity as described as indicated in AOAC (2012) and lycopene according to the method of Nagata and Yamashita (1992).

**Heavy Metals Content in Fruits:** Samples of the 3<sup>rd</sup> picking throughout the season were collected to determine Fe, Zn,

Mn, Cu and Pb. were analyzed in tomato fruits according to Rangana (1977).

**Statistical analysis**

The collected data underwent statistical analysis through Analysis of Variance (ANOVA), and the least significant differences (LSD) method, following the procedure detailed by Duncan (1955), were used to distinguish means at a significance level of 0.05. This analytical approach aligned with the methodology prescribed by Gomez and Gomez (1984).

**RESULTS AND DISCUSSION**

**Results**

**Vegetative growth and leaf pigments:**

Tables 4 and 5 show that there is a positive impact of the different applied organic manure (biochar, compost,

cattle manure and vermicompost) on vegetative growth parameters including plant height, fresh and dry weights, leaf area and leaf pigments such as chlorophylls a, b, total chlorophyll as well as carotenoids of plants irrigated with agricultural drainage water in both seasons. The superior performance of plants (vegetative growth and leaf pigments) was noticed when plants were supplied with biochar. Plants supplied with compost came second followed by plants supplied with vermicompost in most cases in both growing season. In contrast, the lowest values of all mentioned studied parameters were recorded by tomato plants irrigated with agriculture wastewater in the absence of organic manure, this may depend on irrigation water quality.

**Table 4. Vegetative growth characteristics of tomato plants as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Plant height (cm)		Total fresh weight (gm/plant)		Total dry weight (gm/plant)		Leaf area (cm <sup>2</sup> /plant)	
	2022	2023	2022	2023	2022	2023	2022	2023
Biochar	91.1 a	80.0 a	1048.6 a	981.3 a	117.9 a	124.7 a	85.24 a	94.79 a
Compost	83.7 b	76.2 b	932.3 b	885.8 b	102.0 b	107.2 b	68.08 b	79.57 b
Cattel manure	79.3 c	71.3 c	871.4 c	821.1 c	101.4 b	106.5 b	63.08 c	74.08 c
Vermicompost	77.2 c	69.3 c	855.2 c	815.9 c	92.7 c	89.9 c	59.94 c	73.27 c
Control	62.1 d	55.5 d	712.8 d	672.6 d	72.5 d	77.5 d	36.27 d	44.89 d

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Table 5. Photo-synthetic pigments content of tomato leaves as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Chl. (a)(mg/gm F. Wt.)		Chl. (b) (mg/gm F. Wt.)		Total chl. (a + b) (mg/gm F.Wt.)		Carotenoids (mg/gm F. Wt.)	
	2022	2023	2022	2023	2022	2023	2022	2023
Biochar	1.433 a	1.577 a	0.955 a	0.889 a	2.388 a	2.466 a	0.848 a	0.837 a
Compost	1.166 b	1.251 b	0.940 a	0.880 a	2.106 b	2.131 b	0.773 b	0.786 b
Cattel manure	1.163 b	1.248 b	0.827 b	0.750 b	1.990 b	1.998 c	0.687 c	0.709 c
Vermicompost	1.060 c	1.144 c	0.755 b	0.707 bc	1.815 c	1.851 d	0.620 d	0.630 d
Control	1.048 c	1.132 c	0.671 c	0.654 c	1.720 c	1.786 d	0.571 e	0.559 e

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Leaves mineral content:**

Concerning the mineral content in the leaves, it can be noted that N, P, K and Ca elements under agriculture wastewater irrigation significantly increased in tomato leaves in response to all applied organic fertilizers in both growing seasons (Table 6).

Comparing the effects of biochar, compost, cattle manure and vermicompost, plants supplied with biochar are superior in all parameters compared to the control. In contrast, the lowest contents in tomato leaves were observed in plants irrigated with agricultural drainage water in both seasons.

**Table 6. Macro-minerals content of tomato leaves as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	N (%)		P (%)		K (%)		Ca (%)	
	2022	2023	2022	2023	2022	2023	2022	2023
Biochar	2.67 a	3.07 a	0.96 a	1.00 a	3.85 a	3.34 a	0.28 a	0.27 a
Compost	0.94 b	1.32 b	0.69 b	0.74 b	3.14 b	2.91 b	0.22 b	0.21 b
Cattel manure	0.76 bc	1.22 b	0.53 c	0.58 c	3.37 b	2.86 b	0.22 b	0.21 b
Vermicompost	0.45 cd	0.65 c	0.48 c	0.52 c	3.29 b	2.78 b	0.21 b	0.20 b
Control	0.22 d	0.40 c	0.30 d	0.34 d	2.07 c	1.56 c	0.17 c	0.16 c

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Enzymes activity:**

Data in Table 7 show significant differences in peroxidase, superoxide dismutase as well as catalase activity in tomato leaves. In general, the maximum activity of the mentioned antioxidative enzymes was recorded in the

untreated tomato leaves. No significant differences were recorded between the control treatment, compost and cattle manure in catalase and superoxide dismutase activity respectively in the two seasons. Conversely, the lowest values were observed in the leaves of plants supplied with biochar.

**Table 7. Enzymes activity of tomato leaves as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Peroxidase activity (/g F. Wt./h)		Superoxide dismutase activity (/g F. Wt./h)		Catalase activity (/g F. Wt./h)	
	2022	2023	2022	2023	2022	2023
Biochar	194.02 d	193.12 d	102.6 b	102.5 b	179.01 b	187.93 b
Compost	198.9 c	197.8 c	102.6 b	102.5 b	233.72 a	2.44 a
Cattel manure	224.38 b	223.4 b	109.6 a	109.5 a	202.35 b	212.73 b
Vermicompost	201.7 c	200.8 c	104.6 b	104.5 b	204.95 ab	215.23 ab
Control	286.5 a	285.6 a	111.6 a	111.5 a	233.72 a	244.16 a

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Yield and fruit quality:**

Soil application with biochar, compost, cattle manure and vermicompost (Table 8) displayed a significant enhancement in early, total and marketable yield of tomato compared to the untreated plants. Comparing all added organic manure, plants grown in biochar-treated soil showed maximum values of early, total as well as marketable yield in both seasons. Concerning fruit quality, similar significance was noticed in response to soil application with organic

fertilizers in most cases (Table 9). The highest values of total soluble solids, vitamin C, lycopene as well as acidity were observed in response to biochar application. No significant differences were recorded among all organic fertilizers in vitamin C content. Also, no significant differences were found between biochar and compost in total soluble solids in the first season and acidity in both seasons. The lowest values of the mentioned traits were registered in the absence of organic fertilizers.

**Table 8. Yield parameters of tomato plants as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Early yield (ton/fed.)		Total yield (ton/fed.)		Marketable yield (ton/fed.)	
	2022	2023	2022	2023	2022	2023
Biochar	10.428 a	9.469 a	41.278 a	41.711 a	37.629 a	38.582 a
Compost	9.302 b	8.462 b	35.773 b	36.206 b	32.870 b	34.052 b
Cattel manure	9.051 b	8.354 bc	36.775 b	37.208 b	33.652 b	34.370 b
Vermicompost	8.511 c	8.012 bc	33.611 c	34.044 c	30.532 c	31.615 c
Control	7.718 d	7.654 c	30.441 d	30.874 d	27.352 d	28.464 d

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Table 9. Fruits quality parameters of tomato plants as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Total soluble solids (%)		Vitamin C (mg/100g)		Lycopene (mg/100g)		Acidity (%)	
	2022	2023	2022	2023	2022	2023	2022	2023
Biochar	4.53 a	4.56 a	12.119 a	12.714 a	12.62 a	12.47 a	0.39 a	0.51 a
Compost	4.33 a	4.36 b	12.090 a	12.651 a	11.27 c	11.14 b	0.37 a	0.49 a
Cattel manure	4.44 a	4.47 ab	12.091 a	12.685 a	12.14 b	12.00 a	0.33 b	0.45 b
Vermicompost	3.53 b	3.55 c	11.629 a	12.323 a	11.52 c	11.37 b	0.32 b	0.44 b
Control	3.43 b	3.42 c	8.795 b	9.590 b	10.46 d	10.30 c	0.28 c	0.39 c

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Heavy metals:**

Table 10 indicates that there is a reduction in the content of heavy metals (Fe, Zn, Mn, Cu and Pb) in tomato fruits as a result of applying different organic manures (biochar, compost, cattle manure, and vermicompost) in both seasons. In

contrast, plants in the absence of organic manure recorded the best values of the previous mentioned parameters followed by plants supplied with compost in the two seasons. Plants grown in soil supplied with biochar recorded the lowest values of Fe, Zn, Mn, Cu and Pb in tomato fruits.

**Table 10. Heavy metals content of tomato fruits as affected by irrigation with agriculture drainage water and some organic fertilizer treatments during the summer seasons of 2022 and 2023.**

Treatments	Fe (mg/kg dry weight)		Zn (mg/kg dry weight)		Mn (mg/kg dry weight)		Cu (mg/kg dry weight)		Pb (mg/kg dry weight)	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
Biochar	78.4 c	68.4 c	9.86 d	11.20 d	146.2 e	136.1 e	74.2 b	16.1b	21.8 b	28.3 b
Compost	167.7 b	192.4 b	14.27 b	15.27 b	231.6 b	221.5 b	88.7 b	22.0b	25.7 b	37.4 b
Cattel manure	208.7 ab	190.8 b	11.49 c	12.83 c	214.6 c	204.5 c	100.3 b	33.6b	23.9 b	29.6 b
Vermicompost	210.8 ab	203.1 b	10.72 cd	12.06 cd	194.2 d	184.1 d	82.7 b	40.9b	33.3 b	37.4 b
Control	225.7 a	241.3 a	17.19 a	18.52 a	244.2 a	234.1 a	298.6 a	232.3 a	159.7 a	157.5 a

Means followed by the same letter in the same column are significantly different at the 5% level according to Duncan's multiple range test.

**Discussion**

Agriculture is considered the main consumer of fresh water (70 % of fresh water resources) (Kolekar *et al.* 2021), hence water is one of the main limiting factors in agriculture production. Reusing agriculture drainage water is considered the need of the hour because water quantity consumed for agriculture purposes is increasing due to water scarcity and the expansion of agriculture (Dhawan, 2017). Agricultural wastewater provides crops with nutrients due to its nutrient content (Table 3). On the other hand, reusing wastewater have negative impacts such as contamination of heavy metals, microbial pathogenic agents, other hazardous compounds and salinization (Perulli *et al.*, 2021). So, one of the sustainable approaches for managing agricultural drainage water is converting it into useful forms. Also, it is necessary to choose crops that suits irrigation with treated wastewater.


In the current study, the effects of biochar, vermicompost, compost as well as cattle manure as contaminants removal on tomato performance were examined under irrigation with agricultural drainage water. The recorded results confirmed that all the aforementioned treatments significantly improved vegetative growth traits as plant height, fresh and dry weights, leaf area, leaf pigments (chl a, b, total as well as carotenoids), chemical contents including N, P, K and Ca, yield parameters as early, total and marketable yield of tomato as well as fruits quality including total soluble solids, vitamin C, lycopene as well as acidity and fruits heavy metal contents of tomato plants irrigated with agricultural drainage water.

The observed positive effects of all applied organic manure in most cases may be due to the fact that they are considered an enriched source of the most essential nutrients which are necessary for growth and development as well as

microorganisms that greatly affects soil chemical and physical properties and fertility, resulting higher yield (Table 2 and Ganeshnauth *et al.*, 2018). Additionally, wastewater combined with organic treatments stimulates soil metabolism through increasing organic and mineral nutrients, nitrification indices ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ ) as well as metabolic (dehydrogenase activity/water-soluble carbon) (Masciandaro *et al.* 2014). Concerning the observed differences between the applied organic manure, Awogbemi and Oyewumi 2022 illustrated that the efficiency of contaminants removal depends on the type, source, the adsorbent's chemical and physical properties, anion or cation exchange capacity, and the adsorbent's dose.

Regarding the applied biochar, Awogbemi and Kallon, 2023 concluded that, biochar is easy to use, effective, and efficient wastewater contaminant adsorbent. The

application of biochar as a modification technique to enhance wastewater efficiency may be due to its polarity, porosity, high surface area, a high removal of heavy metals, nutrients, organic matter nitrate and other pollutants from wastewater, so it can offer both economic and environmental advantages (Gao *et al.* 2024 and as shown in fig 1). Besides being high contaminant removal from wastewater, it is noted in the above-mentioned results that biochar is the superior as it is considered as a long-term effective soil carbon chelator, improves soil structure and nutrient supply and improves soil microbial diversity (Ren *et al.*, 2020) as well as enhancing root activity (Xiang *et al.* (2017). Similar positive effects of biochar on vegetative growth were studied by Carter *et al.*, (2013) on cabbage and lettuce and Hannachi *et al.*, (2023) on eggplant.



**Fig. 1. The effect of biochar on soil and plant. Feedstock in the form of manure, wood and farm residues can be used for biochar manufacture. Biochar positively influences plant properties like height, number of leaves, leaf area, total biomass, yield, pigments, flavonoids, nutrient uptake and soil properties like pH buffering capacity, WHC, CEC and soil microbiota (Upadhyay *et al.*, 2024).**

The increase in chlorophylls a, b and total as well as carotenoids (Table 5) is the outcome of increasing nutrients availability such as Mg (the component of chlorophyll), N and S nutrients that positively correlate with photosynthesis rate (Gao *et al.* 2021). Similar findings were obtained by Zeeshan *et al.* (2020) on tomato. Biochar can also improve nutrients uptake by enhancing the availability of N, P, K and micronutrients in soil that are vital for biochemical processes of plants (Wong *et al.* 2021) soil microbes (Bolan *et al.* 2023) including bacteria and mycorrhizae. Pavlíková *et al.* (2017) and Durukan *et al.* (2020) observed that N, P, K, and Ca concentrations in lettuce, spinach and sugar beet plants increased with biochar amendment in agreement with the obtained results in Table 6.

The negative response of peroxidase activity, superoxide dismutase as well as catalase activity in response to biochar supplement (Table 7) may be because biochar soil application modulates reactive oxygen species (ROS) levels by affecting the antioxidants defense system (Ghassemi-Golezani and Abdoli 2023). Additionally, improving water and nutrient availability contributes to a balanced soil environment, leading to better plant health, reducing ROS production and enhancing resistance to abiotic and biotic stresses (Mansoor *et al.*, 2021). In the same manner, biochar application resulted in the highest early, total and marketable tomato yield when irrigated with agricultural wastewater

(Table 8). The increased tomato yield in response to biochar application may be attributed the enhanced vegetative growth, chlorophyll content, carotenoids, and leaf mineral content (Tables 4, 5 and 6 respectively) which positively reflected on crop yield. Additionally, the increase in yield may be due to biochar's ability to enhance microbial activity, nutrients availability and improve soil structure (Agbede and Oyewumi, 2022). Gao *et al.* 2021 indicated that increased N and P uptake by tomato plants supplied with biochar resulted in higher tomato yield. Similar findings were obtained by Almaroai and Eissa, 2020 and Obadi *et al.* 2024 on tomato.

The positive effects of biochar application on total soluble solids, vitamin C, lycopene as well as acidity (Table 9) can be attributed to the improved growth (Table 4), increased photosynthetic rate (Table 5), increased nutrient content in leaves (Table 6), improved soil fertility as well as the protection against plant pathogens (Sharma *et al.* 2022). Similar results were observed by several investigators who stated that biochar increased the vitamin C content lycopene and titratable acidity of tomatoes (Almaroai and Eissa, 2020, and Obadi *et al.* 2024 on tomato). Concerning the negative effect of biochar on heavy metal content in fruits (Table 10), this may be because biochar application of can remove nearly 80–98 % of the contaminants (heavy metals, acids, ions, pesticides, N, P, K, Ca, Zn, Mg, Cu, Ni, Na and others) in wastewater, making water suitable for

irrigation and other agricultural or industrial purposes (Awogbemi and Kallon, 2023, Srikaow et al., 2023 and Xu et al., 2023). Additionally, the structure of biochar enhances beneficial microorganisms, adsorbing pollutants and preventing the uptake of organic contaminants and heavy metals by plants, hence contributing to the availability of clean water (Qiu et al., 2021).

The positive impacts of compost on tomato performance may be due to the continuous and slow release of nitrogen from compost into the soil, which increases soil fertility and reduces nitrogen leaching, positively affecting both growth and yield (Weber et al., 2014). Concerning the animal-derived organic manure, it improves soil physical properties as well as soil microbial community, which has a positive effect on plant

performance (Thomas et al., 2019). These findings are in agreement with Gao et al., (2023b) on tomato plants.

Regarding the effects of vermicompost on tomato plants performance, the combination of vermicompost and wastewater is effective in enhancing soil fertility by increasing nutrient availability, microbial activity, and productivity as well as improving quality (Masciandaro et al. 2014 and fig 2). Vermicompost also enriches plant growth and improves microbial activity in the soil, which supports plant growth and enhances photosynthesis (Raza et al. 2024). Similar results were obtained by Alam et al., 2024 indicating that the use of vermicompost and irrigation with wastewater significantly affected tomato plants performance, particularly in terms of growth and yield.



Fig. 2. Vermicompost serves as a plant growth regulator and biocontrol agent, enhancing the physical, biochemical, and fertility properties of soil while suppressing pest attack (Rehman et al., 2023).

### CONCLUSION

The findings of this study highlight the significant potential impacts of biochar, vermicompost, compost and cattle manure on enhancing the growth, yield, and quality of tomato plants irrigated with agriculture wastewater. Furthermore, all treatments increased vegetative growth, photosynthetic pigments, leaf N, P, K and Ca content which consequently led to improving yield as well as fruits quality of tomato plants. Thus, the applied organic manure could be used for contaminant removal from agricultural wastewater, not only to overcome water scarcity and protect fresh water resources, but also to increase productivity and improve quality. Moreover, these results underscore the importance of reusing agricultural wastewater in tomato production. Moving forward, further exploration and adoption of organic manure in other crops irrigated with agricultural wastewater should also be recommended, as part of a holistic strategy towards sustainable agriculture. By harnessing the benefits of these organic materials, we can not only improve the agricultural productivity but also contribute to environmental conservation and food security goals.

### REFERENCES

Agbede, T. M. and Oyewumi, A. (2022). Soil properties, sweet potato growth and yield under biochar, poultry manure and their combination in two degraded Alfisols of humid tropics. *Sci Hortic.*, 304: 111331.

Alam, M., Khan, A., Zaman, R., Khan, S., Khan, M. A., Ahmad, I., Jalal, A. and Kim, K. I. (2024). Vermi-remediation impacts on growth and metals bioaccumulation in tomato irrigated with wastewater. *Chemosphere*, 362, 142848.

Almaroai, Y. A. and Eissa, M. A. (2020). Effect of biochar on yield and quality of tomato grown on a metal-contaminated soil. *Sci Hortic.*, 265: 109210.

AOAC International (2012). Official Methods of Analysis of AOAC International. 19<sup>th</sup> edition. 2<sup>nd</sup> revision. Gaithersburg, MD, USA, Association of Analytical Communities.

Awogbemi, O. and Kallon, D.V.V. (2022). Pretreatment techniques for agricultural waste, *Case Stud. Chem. Environ. Eng.*, 6: 100229.

Awogbemi, O., & Von Kallon, D. V. (2023). Progress in agricultural waste derived biochar as adsorbents for wastewater treatment. *Applied Surface Science Advances*, 18, 100518.

Bolan, S., Hou, D., Wang, L., Hale, L., Egamberdieva, D. and Tammeorg, P. (2023). The potential of biochar as a microbial carrier for agricultural and environmental applications. *Sci. Total Environ.*, 886: 163968.

Cao, S.; Q. Xu; Y. Cao; K. Qian; K. An; Y. Zhu; H. Binzeng; H. Zhao and B. Kuai (2005). Loss-of- function mutation in DET2 gene lead to an enhanced resistance to oxidative stress in Arabidopsis. *Physiol. Plant.*, 123:57-66.

Carter, S., Shackley, S., Sohi, S., Suy, T. B. and Haeefe, S. (2013). The impact of biochar application on soil properties and plant growth of pot grown lettuce (*Lactuca sativa*) and cabbage (*Brassica chinensis*). *Agronomy*, 3(2):404–18.

Dhawan, V. (2017). Water and agriculture in India: background paper for the South Asia expert panel during the Global Forum for Food and Agriculture (GFFA)

- Duncan, B.D. (1955). Multiple Range and Multiple F-Tests. *Biometrics*, 11: 1-42.
- Durukan, H., Demirbas, A. and Turkecul, I. (2020). Effects of biochar rates on yield and nutrient uptake of sugar beet plants grown under drought
- Faloye, O., Alatise, M., Ajayi, A. and Ewulo, B. (2019). Effects of biochar and inorganic fertiliser applications on growth, yield and water use efficiency of maize under deficit irrigation. *Agric. Water Manag.*, 217, 165–178.
- FAOSTAT, F. (2022). Food and agriculture organization of the United Nations. *Statistical database*.
- Fischer, B. M., Manzoni, S., Morillas, L., Garcia, M., Johnson, M. S. and Lyon, S. W. (2019). Improving agricultural water use efficiency with biochar—A synthesis of biochar effects on water storage and fluxes across scales. *Science of the Total Environment*, 657, 853-862.
- Ganeshnauth, V., Jaikishun, S., Ansari, A. A., & Homenauth, O. (2018). The effect of vermicompost and other fertilizers on the growth and productivity of pepper plants in Guyana. *Automation in Agriculture-Securing Food Supplies for Future Generations*.
- Gao, F., Li, H., Mu, X., Gao, H., Zhang, Y., Li, R., ... & Ye, L. (2023b). Effects of organic fertilizer application on tomato yield and quality: a meta-analysis. *Applied sciences*, 13(4), 2184.
- Gao, R., Ding, S., Liu, Z., Jiang, H., Liu, G., and Fang, J. (2024). Recent advances and perspectives of biochar for livestock wastewater: modification methods, applications, and resource recovery. *Journal of Environmental Chemical Engineering*, 113678.
- Gao, Y., Shao, G., Cui, J., Lu, J., Tian, L., Song, E. and Zeng, Z (2023a). Effects of Drought Hardening and Saline Water Irrigation on the Growth, Yield, and Quality of Tomato. *Agronomy*, 13, 2351.
- Gao, Y., Shao, G., Yang, Z., Zhang, K., Lu, J. and Wang, Z. (2021). Influences of soil and biochar properties and amount of biochar and fertilizer on the performance of biochar in improving plant photosynthetic rate: a meta-analysis. *Eur J Agron.*, 30: 126345.
- Ghassemi-Golezani, K. and Abdoli, S. (2023). Alleviation of salt stress in rapeseed (*Brassica napus* L.) plants by biochar-based rhizobacteria: new insights into the mechanisms regulating nutrient uptake, antioxidant activity, root growth and productivity. *Arch Agron. Soil Sci.*, 69(9):1548–1565.
- Gomez, K.A., and Gomez, A.A. (1984). “Statistical procedures for agricultural research”. John Wiley and Sons, Inc., New York. 680 p.
- Hannachi, S., Signore, A. and Mechi, L. (2023). Alleviation of Associated Drought and Salinity Stress’ Detrimental Impacts on an Eggplant Cultivar (‘Bonica F1’) by Adding Biochar. *Plants*, 12, 1399.
- Hawary, A. and Shaban, M. (2018). Improving drainage water quality: constructed wetlands-performance assessment using multivariate and cost analysis. *Water Sci.*, 32, pp. 301-317
- Hou, X., Zhang, W., Du, T., Kang, S. and Davies, W.J. (2020). Responses of water accumulation and solute metabolism in tomato fruit to water scarcity and implications for main fruit quality variables. *J. Exp. Bot.*, 71 (4), pp. 1249-1264
- Kalra, Y. P. (1998). *Handbook of Reference Methods for Plant Analysis*. CRC Press Inc., Boca Raton, FL, USA.
- Kolekar, S., Sankapal, P., Khare, K., & Chinnasamy, P. (2023). Assessment of recycled treated wastewater for sustainable tomato crop production: A comprehensive review. *Environmental Challenges*, 13(12)100791.
- Kolekar, S., Joshi, R., Kadam, K., Khare, K. and Chinnasamy, P. (2021). Bibliometric survey on reuse of treated wastewater for agriculture. *Libr. Philos. Pract.*, 1, pp. 1-26.
- Koller, H.R. (1972). Leaf area-Leafm weight relationship in soybean canopy. *Crop Sci.*, 12:180-183.
- Lee, N., Hong, S. H., Lee, C. G., Park, S. J., & Lee, J. (2021). Conversion of cattle manure into functional material to remove selenate from wastewater. *Chemosphere*, 278, 130398.
- Mansoor, S., Kour, N., Manhas, S., Zahid, S., Wani, O.A., Sharma, V. (2021). Biochar as a tool for effective management of drought and heavy metal toxicity. *Chemosphere*, 271: 129458.
- Masciandro, G., Peruzzi, E., Doni, S. and Macci, C. (2014). Fertigation with wastewater and vermicompost: soil biochemical and agronomic implications. *Pedosphere*, 24(5), 625-634.
- Mohammed, I. Q., and Alkobaisy, J. S. (2024). Effect of bio, vermicompost and mineral fertilizers on some soil properties and soybean (*Glycine Max*
- Nagata, M. and I. Yamashita. (1992). Simple method for simultaneous determination of chlorophyll and carotenoids in tomato fruit. *J. Japan Soc. Food Sci., Technol.* 39: 928–932.
- Obadi, A., Alharbi, A., Alomran, A., Alghamdi, A. G., Louki, I., Alkhasha, A. and Alqardaei, T. (2024). Enhancement in Tomato Yield and Quality Using Biochar Amendments in Greenhouse under Salinity and Drought Stress. *Plants*, 13(12), 1634.
- Paudel, A., Bar-Tal, G.J., Levy, N., Rotbart, J. and Ephrath, S., (2018). Treated wastewater irrigation: soil variables and grapefruit tree performance. *Agric. Water Manag.*, 1, pp. 126-137.
- Pavlíková D, Zemanova V, Břendová K, Kubátová P, Tlustoš P (2017). Effect of biochar application on the content of nutrients (Ca, Fe, K, Mg, Na, P) and amino acids in subsequently growing spinach and mustard. *Plant Soil Environ.*, 63(7):322–7
- Perulli, G.D., Gaggia, F., Sorrenti, G., Donati, I., Boini, A., Bresilla, K., Manfrini, L., Baffoni, L., Di Gioia, D., Grappadelli, L.C., Spinelli, F., Morandi, B. (2021). Treated wastewater as irrigation source: a microbiological and chemical evaluation in apple and nectarine trees. *Agric. Water Manag.*, 244 (3), 106403
- Qadir, M., Drechsel, P., Jiméñez Cisneros, Kim, B., Pramanik, Y.A., Mehta, P. and Olaniyan, O. (2024). Global and regional potential of wastewater as a water, nutrient and energy source, *Nat. Resour. Forum* 44: 40–51.
- Qiu, M., Hu, B., Chen, Z., Yang, H., Zhuang, L. and Wang, X. (2021). Challenges of organic pollutant photocatalysis by biochar-based catalysts. *Biochar*. 3(2):117–23
- Rangana, S. (1979). *Manual Analysis of Fruit and Vegetable Products*. Tata McGraw Hill Pub. Co. Ltd. New Delhi, pp 363.
- Rangana, S. (1977). *Manuals of Analysis of Fruit and Vegetable Products*. Tata Mc Graw Hill Publishing Co Ltd., New Delhi.

- Raza, S. T., Rong, L., Rene, E. R., Ali, Z., Iqbal, H., Sahito, Z. A., and Chen, Z. (2024). Effects of vermicompost preparation and application on waste recycling, NH<sub>3</sub>, and N<sub>2</sub>O emissions: A systematic review on vermicomposting. *Environmental Technology & Innovation*, 103722.
- Rehman, S.u.; De Castro, F.; Aprile, A.; Benedetti, M.; Fanizzi, F.P. (2023). Vermicompost: Enhancing Plant Growth and Combating Abiotic and Biotic Stress. *Agronomy*, 13, 1134. <https://doi.org/10.3390/agronomy13041134>
- Ren, H., Huang, B., Fernandez-Garcia, V., Miesel, J., Yan, L. and Lv, C. (2020). Biochar and rhizobacteria amendments improve several soil properties and bacterial diversity. *Microorganisms*, 8, p. 502.
- Seyoum, M.M., Lichtenberg, R., Orlofsky, E., Bernstein, N., Gillor, O., (2022). Antibiotic resistance in soil and tomato crop irrigated with freshwater and two types of treated wastewater. *Environ. Res.* 211, 1–6.
- Sharma, S., Rana, V.S., Rana, N., Prasad, H., Sharma, U. and Patiyal, V. (2022). Biochar from fruit crops waste and its potential impact on fruit crops. *Sci Hort.*, 299: 111052
- Srikhaow, A., Win, E.E., Amornsakchai, T., Kiatsiriroat, T. Kajitvichyanukul, P. S. and Smith, M. (2023). Biochar derived from pineapple leaf non-fibrous materials and its adsorption capability for pesticides, *ACS Omega* 8: 26147–26157.
- Thomas, C.L.; Acquah, G.E.; Whitmore, A.P.; McGrath, S.P.; Haefele, S.M. (2019). The Effect of Different Organic Fertilizers on Yield and Soil and Crop Nutrient Concentrations. *Agronomy*, 9, 776.
- Toledo, M., Siles, J., Gutiérrez, M. and Martín, M. (2018). Monitoring of the composting process of different agroindustrial waste: Influence of the operational variables on the odorous impact. *Waste Manag.*, 76, 266–274.
- Turan, M., Kocaman, A., Tüfenkçi, Ş., Katırcıoğlu, H., Güneş, A., Kıtır, N., and Yıldırım, E. (2023). Development of organic phosphorus vermicompost from raw phosphate rock using microorganisms and enzymes and its effect on tomato yield. *Scientia Horticulturae*, 321, 112323.
- Tzortzakakis, N., Saridakis, C., & Chrysargyris, A. (2020). Treated wastewater and fertigation applied for greenhouse tomato cultivation grown in municipal solid waste compost and soil mixtures. *Sustainability*, 12(10), 4287.
- Upadhyay, V., Choudhary, K.K. & Agrawal, S.B. (2024). Use of biochar as a sustainable agronomic tool, its limitations and impact on environment: a review. *Discov Agric* 2, 20.
- Weber, J., Kocowicz, A., Bekier, J., Jamroz, E., Tyszcza, R., Debicka, M., ... and Kordas, L. (2014). The effect of a sandy soil amendment with municipal solid waste (MSW) compost on nitrogen uptake efficiency by plants. *European journal of Agronomy*, 54, 54-60.
- Wettstein, D. (1957). "Chlorophyll". *Lethale under Submikroskopische Formwechsel derplastiden*. *Exptl. Cell Res.*, 12: 427-506.
- Wong, J. W. and Ogbonnaya, U. O. (2021). Biochar porosity: a nature-based dependent parameter to deliver microorganisms to soils for land restoration. *Environ Sci. Pollut. Res.*, 28:46894–909.
- Xiang, Y., Deng, Q., Duan, H. and Guo, Y. (2017). Effects of biochar application on root traits: a meta-analysis. *GCB Bioenergy*, 9(10):1563–72. 55.
- Xu, C., Liu, R. and Chen, L. (2023). Removal of phosphorus from domestic sewage in rural areas using oyster shell-modified agricultural waste–rice husk biochar, *Processes*, 11 (9): 2577.
- Zeeshan, M., Ahmad, W., Hussain, F., Ahamd, W., Numan, M., Shah, M. and Ahmad, I. (2020). Phytostabilization of the heavy metals in the soil with biochar applications, the impact on chlorophyll, carotene, soil fertility and tomato crop yield. *J. Clean. Prod.*, 255, 120318.

## إعادة استخدام مياه الصرف الزراعي لإنتاج الطماطم باستخدام مصادر مختلفة من الأسمدة العضوية تحت ظروف منطقة الدلتا

عبد البصير السيد عبدالصير<sup>1</sup>، ولاء محمد السعيد سويلم<sup>2</sup>، ندى علاء الدين مصطفى السيد<sup>3</sup>، وأميرة عبدالبدیع أبو المكارم<sup>4</sup>

<sup>1</sup>قسم بحوث الخضار ذاتية التلقيح – معهد بحوث البساتين – مركز البحوث الزراعية – الجيزة - مصر

<sup>2</sup>قسم الخضار والزينة – كلية الزراعة – جامعة المنصورة - مصر

<sup>3</sup>قسم النبات الزراعي – كلية الزراعة – جامعة المنصورة - مصر

<sup>4</sup>قسم تداول الخضار – معهد بحوث البساتين – مركز البحوث الزراعية – الجيزة - مصر

### المخلص

ينخفض توافر المياه العذبة (كما ونوعاً) في جميع أنحاء العالم و يعد القطاع الزراعي هو الأكثر استهلاكاً للمياه العذبة لذلك هناك حاجة ملحة لتطوير الممارسات الزراعية لسد الفجوة بين الإحتياجات والمتوافر (العرض والطلب) على المياه باعتبارها واحدة من الركائز الأساسية للزراعة المستدامة. ومن هنا فإن إحدى الاستراتيجيات الممكنة لتحقيق ذلك هي إعادة استخدام مياه الصرف الزراعي خاصة في ظل تحديات التغيرات المناخية. لذلك يهدف البحث إلى إعادة استخدام مياه الصرف الزراعي في إنتاج الطماطم من خلال إضافة بعض الأسمدة العضوية. ومن أجل تحقيق ذلك تم دراسة تأثير كل من البيوشار والكمبوست والسماد البودي والسماد البلدي على النمو الخضري والمحتوى الكيميائي للأوراق وكذلك إنتاجية نباتات الطماطم ومحتوى الثمار من العناصر الثقيلة. أظهرت النتائج أن جميع الأسمدة العضوية المضافة إلى التربة أدت إلى تحسين مؤشرات النمو الخضري المتمثلة في ارتفاع النبات و الوزن الطازج والجاف و المساحة الورقية وذلك مقارنة بالنباتات التي تم الري بماء الصرف الزراعي فقط. كذلك أدت هذه المعاملات إلى زيادة معنوية في محتوى الأوراق من صبغات البناء الضوئي والعناصر الغذائية وكذلك محصول وجودة ثمار الطماطم. وعلى العكس سجل نشاط الإنزيمات المضادة للأكسدة وكذلك محتوى الثمار من العناصر الثقيلة أقل القيم. سجلت النباتات التي تم تسقيها بالبيوشار أفضل القيم للقياسات الخضريّة وصبغات البناء الضوئي ومحتوى الأوراق من العناصر الغذائية والمحصول وجودة الثمار ولكن أقلها للنشاط الإنزيمي والمعادن الثقيلة. وبصفة عامة يشير البحث إلى إمكانية إعادة استخدام مياه الصرف الزراعي لإنتاج محصول الطماطم وتقليل الأثر المترتبة على ذلك عن طريق إضافة الأسمدة العضوية مثل البيوشار أو السماد البودي أو الكمبوست أو السماد البلدي تحت ظروف المنطقة محل الدراسة.

**الكلمات الدالة:** الطماطم – مياه الصرف الزراعي – البيوشار – السماد البودي