BEHAVIOR OF EGYPTIAN HYBRID RICE CULTIVAR (EHRI) UNDER DIFFERENT IRRIGATION REGIMES AND HARVESTING TIMES

Omar, A. M.¹; I. S. El-Refaee² and Hasnaa A. Ghazy²

¹ Agron. Dept., Fac. Agric., Kafr El-Sheikh University, Egypt

² Agric. Res. Center, Field Crops Res. Inst., Rice Res. & Train. Center,

33717 Sakha - Kafr El-Sheikh, Egypt

ABSTRACT

These studies were carried out at the Experimental Farm of Rice Research and Training Center (RRTC), Sakha, Kafr El-Sheikh, Egypt, during 2007 and 2008 summer seasons. The purpose was to investigate the effect of irrigation intervals, cutoff irrigation and harvesting times on grain yield and its attributes as well as grain quality characters and water relations of Egyptian hybrid rice cultivar (EHR1). A split split–plot design, with four replicates, was used. The main plots were devoted to three irrigation intervals namely: continuous flooding (CF), irrigation every 6 (6D) and 12 days (12D). The sub-plots were assigned to five times for cut-off irrigation, i.e. cut-off irrigation after complete heading (ACH), 1-, 2-, 3- and 4- weeks after complete heading (WACH). However, the sub-sub plots were consisted of three harvesting times, i.e. harvest rice plants at 10-, 15- and 20- days after cut-off irrigation (DACI).

The main results showed that grain yield and its attributes, as well as milling recovery and protein content, were significantly decreased as irrigation intervals increased from CF up to twelve days, but, unfilled grains (%) and amylose content were increased. Irrigation every six days came in between with insignificant difference with CF in most of previously mentions traits. Delayed cut-off irrigation up to 3- and 4-WACH significantly increased grain yield and most of its attributes, as well as grain quality. However, it reduced amylose content in 2007 and unfilled grains (%) in both seasons, as compared with cut-off irrigation ACH. Harvesting rice plants 10 and 15 DACI recorded the highest number of grains/panicle, panicle weight, 1000 grain weight and grain yield. However, the lowest values of milling recovery and protein content were recorded when plants were harvested at 10 DACI.

Continuous flooding consumed the highest amount of irrigation water, while increasing irrigation intervals up to six and twelve days tended to decrease the amount of water used. Furthermore, 6D treatment recorded the highest water productivity (0.796 and 0.798 kg/m³) and minimum grain yield reduction (4.54 and 3.64 %) with water saved about 8.04 and 7.08 %, compared to CF in both seasons, respectively. Delay cut-off irrigation recorded higher water productivity than early cut-off irrigation after complete heading.

Generally, in case of water shortage, it was concluded to use irrigation every six days and cut-off irrigation 3-WACH with harvest plants 10 DACI for the highest water productivity and grain yield, as well as acceptable grain quality characters, of Egyptian hybrid rice cultivar (EHR1).

Keywords: Hybrid rice, irrigation regimes and harvesting times.

INTRODUCTION

Hybrid rice technology is one of the innovative breakthroughs that can further increase rice production and food security in Egypt. Hybrid rice varieties can out-yield conventional modern ones by 19 % even at the same

input levels (Lin, 1994). In general, rice hybrid have yield advantage of 1.0 to 1.5 t/ha over inbred high yielding rice varieties, yielding 5 to 6 t/ha in India (Virmani, 2002).

Rice (Oryza sativa L.) is a simi-aquatic plant and does not need standing water for a successful rice crop. In Egypt, the dominant practice in rice production is flooded irrigation, which consumes large amounts of water as being approximately 18 % of the total water resources (Badawi et al., 2002). Rice, however, it is a heavy consumer of freshwater and approximately 50 % of the freshwater used in Asian agriculture was used for rice production (Guerra et al., 1998). With limited water resources, a future increases in rice production requires the development of water saving technologies. In Egypt, water resources are not sufficient for both reclaiming and irrigation purposes for the soil. So, saving some of irrigation water is a necessity demand to face this problem, in the future. Several water saving irrigation techniques for rice have been reported previously. The most widely adopted water saving practice in China was alternate wetting and drying (Zhi, 1993). Other way to save water was by increasing irrigation intervals with minimum grain yield reduction. In Egypt, increasing intervals between irrigations, wherever, allowed the rice fields to dry for a few days between irrigations for six to eight days. Awad (2001) found that grain yields were not affected by irrigation intervals, ranging from four to eight days. Belder et al., (2005) reported that water productivity was higher, in the alternately submerged and nonsubmerged regimes, than in the continuous submerged regime.

Scheduling last irrigation (cut-off irrigation) at the correct stage of maturity may influence moisture content in grain at harvest and, then, affect grain yield and quality. Whereas, moisture content in the grain at harvest affects the head rice recovery (Govindaswamy and Ghosh, 1970). Cut-off irrigation on rice fields early may cause moisture stress in grains before they are physiologically mature. Early cut-off irrigation may lead to lower harvest moisture contents associated with lower head rice yield. This practice causes a significant reduction in grain yield and increasing the unfilled, immature and broken, grains. Current recommendations for cutting off irrigation suggest maintaining the flood until 25 to 28 days after 50 % heading to prevent reduction in grain yield and milling quality (Slaton, 2001). This corresponds to physiological maturity for most of the kernels on the panicle. El-Refaee (2007) reported that withholding of irrigation 21 days after heading is considered as the optimum timing of last irrigation to rice field to get high grain yield and quality.

Cut-off irrigation and harvesting times are important factors to be considered for an optimum rice harvest and they are keys to high head rice yields. Harvest management preserves rice quality and grain yield that contribute directly to profit. Early harvest may lead to increases the unfilled and immature grains. This prematurely grains may result in partially chalky kernels and milk white kernels and increasing the thickness of the bran and aileron layers (Dewedar, 2004). Harvest, generally, commences in most countries when the average moisture content of the grains on the panicles is in the 20 - 27 % (Elaine *et al.*, 2005). Hossain *et al.*, (2009) concluded that the rice harvesting at 30 to 35 days after flowering (DAF) was found to be

suitable for higher grain quality, in respect of head rice outturn, and amylose content. However, the highest protein content was obtained from early harvesting (25 DAF).

The present study aimed to find out the best irrigation intervals, as well as irrigation cut-off after complete heading and harvest times, for higher grain yield and acceptable grain quality characteristics of EHR1 hybrid rice cultivar.

MATERIALS AND METHODS

Two field experiments were conducted at the Experimental Farm of Rice Research and Training Center (RRTC), Sakha Kafr El-Sheikh, Egypt, during 2007 and 2008 summer seasons. The purpose was to investigate the effect of irrigation intervals, cut-off irrigation after complete and harvesting times on grain yield and its attributes, as well as grain quality characters and water relations, of Egyptian hybrid rice cultivar (EHR1). The meteorological data of the experiment sites are presented in Table 1 according to Sakha Meteorological Station.

A split-split plot design, with four replicates, was used in this study. The main plots were devoted to three irrigation intervals namely: continuous flooding (CF), irrigation every 6 (6D) and 12 days (12D) with 5-6 cm water head at the time of water addition. The sub plots were assigned to five timing of cut-off irrigation, i.e. cut-off irrigation after complete heading (ACH), 1, 2, 3 and 4 weeks after complete heading (WACH). However, the sub-sub plots were consisted of three harvesting times namely: 10, 15 and 20 days after cut-off irrigation (DACI).

Ч	enou										
			2007		2008						
Month	A tempe (º(ir Frature C)	RH %		E (mm/	A tempe (º(ir rature C)	RI	Η%	E (mm/	
	Max.	Mini.	7:30	13:30	day)	Max.	Mini.	7:30	13:30	day)	
May	30.0	12.0	76.3	45.0	6.48	29.0	10.0	70.5	42.5	6.91	
June	33.0	16.5	82.4	56.0	7.61	33.0	15.0	82.5	50.0	7.33	
July	32.6	17.3	83.0	54.0	6.88	32.0	15.7	80.0	55.7	6.79	
August	32.5	17.2	83.0	56.5	6.40	33.0	16.3	83.2	56.0	6.53	
September	32.0	13.0	74.5	52.0	5.56	33.5	15.0	77.3	47.7	6.08	
Mean	32.02	15.2	79.84	58.7	6.59	32.1	14.4	78.7	50.38	6.73	

Table 1: Monthly temperature means, relative humidity (RH) and pan evaporation (E) at the study area during the experimental period

The experiments were preceded by barley (*Hordeum vulgare L.*) in both seasons. The soil was clay with pH 8.1 and 8.2 and an organic matter content of 1.7 and 1.6 %. The total N was 500 and 515 ppm in both seasons, respectively. Phosphorus fertilizer, at the rate of 35.5 kg P₂O₅/ha, was applied to the soil during land preparation. Potassium fertilizer was added, at rate of 57 kg K₂O/ha, as basal dose and incorporated into dry soil. The recommended dose of Nitrogen fertilizer (165 kg N/ha), in the form of Urea (46 % N), was applied in three equal splits (as basal, top dressing at panicle

initiation and late booting). Seeds of EHR1 hybrid rice cultivar, at a rate of 24 kg/ha, were soaked in sufficient water for 24 hours and incubated for another 48 hours to enhance germination. Zinc (Zn So₄), as well as all other cultural practices, was applied as recommended. The experiments were sown on 3rd and 5th of May in the two successive seasons. Two to three, thirty days old, seedlings were transplanted at 20 cm distance between hills and rows in plot size of 30 m² (5 x 6 m) each.

To avoid lateral movement and more water control, each main plot was separated by two meter wide ditches. Water pump provided with a calibrated water meter was used for all water measurements. Water productivity (WP) was calculated as weight of grains per unit of water used (kg grain/m³ water).

At harvest, plant height was estimated and total number of panicles were counted and, then, conformed to numbers/m². Ten random panicles were collected from each sub sub-plot to estimate panicle length, total grains/panicle, unfilled grain percentage, panicle weight and 1000-grain weight. Grain and straw yields was measured from 12 m² (3 x 4 m) and adjusted to 14 % moisture content and, then, conformed to t/ha. Harvest index was determined according to Yoshida (1981) by subdividing grain yield (economic yield) on the total dry weight of grains and straw (biological yield).

Milling recovery (hulling, milling and head rice %) were measured according to the method described by Juliano (1971). Improved Kjeldahl methods of A.O.A.C. (1970) was used to determine the N content, then, multiplied by the factor of 5.95 to estimate the crude protein in rough grains. Amylose content was estimated according to Williams *et al.* (1958).

The analysis of variance was carried out according to Gomez and Gomez (1984) using GENSTAT 5th Edition Computer Program. Means were compared, using the least significant differences (LSD) at 5% probability level.

RESULTS AND DISCUSSION

Grain yield and its attributes:

Prolonged irrigation intervals caused a remarkable reduction in grain yield and its attributes (Fig. 1 and Table 2). In both seasons, continuous flooding (CF) followed by irrigation every 6 days (6D) gave the highest values of plant height, number of panicles, number of grains/panicle, panicle weight, 1000 grain weight, harvest index, grain and straw yields. However, increased irrigation intervals up to 12 days (12D) significantly reduced all previously mentions traits. While, unfilled grains (%) reached the maximum values. Grain yield was 11.25, 10.74 and 8.08 t/ha, in the first season, and 11.76, 11.33 and 8.73 t/ha, in the second season, for CF, 6D and 12D, respectively. These results indicated that exposed rice plant to drought stress (12D) caused significant reduction in grain yield, this held true since all yield components were affected by such conditions. It is a fact that the unavailability of water inhibits the production of dry matter content in the different plant organs, because of the water imbalance inside the pant, beside number of panicles/m², number of filled grain/panicle and grain yield. Similar finding were reported by Awad (2001) and El-Gewaily (2006).



Fig. 1: Plant height, No. of panicles and panicle length of Egyptian hybrid rice cultivar (EHR1) as influenced by irrigation intervals

The effect of cut-off irrigation times on grain yield and its attributes were significant, except straw yield in both seasons (Table 2). Delayed irrigation termination up to 3- and 4-WACH significantly increased grain yield and most of its attributes. However, it reduced unfilled grains (%) as compared with cut-off irrigation after complete heading. The increased grain yield attributes were the reason for increasing grain yield with cut-off irrigation at 21 days (3 weeks) after complete heading. Continuing irrigation up to 3-WACH might be improved the translocation of photosynthates and thereby increased the grain yield. However, under early cut-off irrigation the carbohydrates tended to decrease due to the shortage of water required to photosynthetic processes and the restriction of translocation of insufficient metabolite form source to sink owing to plant-water deficit, which, led to decrease the grain yields. These results are in conformity with the findings of Uppal and Bali (1994) and El-Refaee (2007).

Grain yield and some of its attributes were significantly affected by harvesting times (Table 2). Harvesting plants 10 DACI recorded the highest number of grains/panicle, panicle weight and 1000 grain weight, followed by 15 DACI, in both seasons. However, the lowest values of all previous characters were obtained when plants were harvested 20 DACI as compared with the other harvest time. The highest grain yields (10.23 and 10.70 t/ha) were obtained when plants harvested 10 DACI and (10.02 and 10.77 t/ha) 15 DACI, with no significant difference between each other. While, the lowest values (9.82 and 10.35 t/ha) were recorded when plants harvested 20 DACI in the two respective seasons. It is means that harvested rice plants after 10 or 15 DACI were the optimum, because of the complete filling, resulted in produces the heaviest panicles and grains. These results are in agreement with those of Asano *et al.* (2000) and Dewedar (2004).

Omar, A. M. et al.

Data in Table 3 showed that the highest number of grains/panicle, in 2007, was obtained when cut-off irrigation 1-WACH with harvest plants 10-DACI were combined. Continuous flooding with 2-WACH recorded that the highest number of grains/panicle in 2008. The lowest value of unfilled grains (%) was obtained by either cut-off irrigation 3- and 4-WACH or harvesting plants 20 DACI under CF treatment in the first season, however, the lowest values (9.1 and 7 %) of unfilled grains (%) were obtained when cut-off irrigation 3- and 4-WACH with harvest plants 20-DACI in both seasons, respectively. The highest values of panicle weight (3.89 g) was recorded when irrigation terminated 4-WACH and harvest plants 10-DACI in 2007. Continuous flooding with cut-off irrigation 3-WACH recorded the highest grain yield (12.43 and 12.60 t/ha) in both seasons, respectively (Table 3). The highest values of straw yield were obtained when irrigation terminated after complete heading with harvest plants 10-DACI in 2007. Irrigation every 6 days with cut-off irrigation 4- and 3- WACH recorded the highest values of harvest index in both seasons, respectively (Table 3).

Grain quality:

Data in Table 4 revealed that milling recovery (hulling, milling and head rice %) and both amylose and protein contents were significantly affected by irrigation intervals in both seasons, except hulling (%) in 2008. Continuous flooding caused an increase in milling recovery and protein content. However, 12D decreases all previous traits, under study, but, gave the highest amylose content. Irrigation every 6 days comes in between with insignificant difference with CF in the most of the studied characters. These results could be attributed to the decrease in moisture content of the grains leading to the reduction in milling recovery percentage. The same trend was found by Nour *et al.* (1994) and El-Refaee (1997).

Early cut-off irrigation after heading significantly decreased milling recovery and protein content as compared with other treatments in both seasons (Table 4). However, cut-off irrigation early after complete heading, in 2007, and 4-WACH, in 2008, significantly decreased amylose content. The milling recovery in the last two treatments (3- and 4-WACH) was high due to the corresponding increase in the moisture content of the grain. The lowest milling recovery in the treatment received last irrigation ACH and 1-WACH might be due to early drying of the grains before the completion of ripening. Besides, delay in harvesting along with the other treatments may result in low moisture content in grain during day and in night some amount of moisture is re-absorbed by the deposition of dew. Such alternate drying and wetting cycles cause mechanical stress which may result in the development of fissures and ultimately higher percentage of broken rice. The results are in agreement with those obtained by Uppal and Bali (1994) and El–Refaee (2007).

Table 3:	Grain yield and	some of its attrib	outes of Egyptia	n hybrid rice
	cultivar (EHR1)	as influenced by	the interaction	between the
	studied factors.			

				-									
<u> </u>	Cut-off		Ha	arvest date	2007			Irriga	tion treatmen	t 2008			
ō,	irrigation	10-D/	ACI	15-DACI		20-	DACI	CF	6 days	12 days			
5	ACH	166	7	177.3		1(67.5	165.3	185.8	185.5			
in	1-WACH	188	.4	167.1		10	69.6	157.6	182.4	194.9			
å	2-WACH	175	.5	176.1		10	69.1	172.0	182.7	201.7			
us	3-WACH	165	.6	173.6		10	66.0	174.4	186.3	193.6			
o.ia	4- WACH	158	.7	164.0		14	48.7	144.4	180.5	186.7			
Zō	L.S.D 5%			9.8				12.3					
			Cut-	off irrigatio	n 200	07		н	arvest date 20	07			
	Irrigation		1. 2.		3-	3- 4-							
	treatment	ACH	WACI	H WĀCH	WĂ	VACH WACH		10-DACI	15-DACI	20-DACI			
	CE	13.0	11 1	86	63	6.2 6.2		93	9.8	8.0			
~	6 days	16.3	13.5	12.7	12	2	11.5	14.6	12.4	12.8			
%	12 days	21.5	13.9	12.8	12.	.9	11.5	14.3	17.0	13.0			
S	L.S.D 5%			2.1					1.7	•			
air	2.0.2 070						Harvest	date					
Ъ	Cut-off			2007			nai vest	uuto	2008				
led	irrigation	10 D		15 DACI		20			15 DACI				
Ę	1011	10-07		13-DACI		20-		IU-DACI	15-DACI	20-DACI			
5		15.	8	20.6		4	2.0	24.3	24.6	17.0			
		13.	/ 8	13.1		1	1.0	22.0	12.0	0.0			
	3-WACH	11.	4	10.9		1	0.9	85	77	83			
	4- WACH	11.	1	9.9		ġ	9.9	9.0	8.0	7.0			
	LSD5%			21					3.0				
	Cut-off		2.1				Harvest date 2007						
Ħ	irrigation		10-D/			110	15-DA		20-D	ACI			
igl	ACH		3.2	2			2.88	01	200	6			
×			3.21	5			2.00		2.3	2			
e e	2-WACH		3.6	5			3.62		3.3	6			
-ic	3-WACH		3.50	0			3.67		3.0	3			
Ра	4- WACH		3.89	9		3.75			3.2	.1			
	L.S.D 5%				0.32								
	0	Irrigation treatment											
a)	irrigation			2007					2008				
Ę	inigation	CF		6 days		12 days		CF	6 days	12 days			
р	ACH	9.6	8	9.55		7	.69	10.16	9.81	7.89			
/ie	1-WACH	10.5	53	9.92		7	.87	11.46	11.18	8.83			
ŝ	2-WACH	11.6	62	10.89		8.42		12.20	11.65	8.68			
rai	3-WACH	12.4	13	11.66		8.39		12.60	11.96	9.78			
G	4- WACH	12.0	JU	11.68		8	3.03	12.39	12.06	8.45			
	L.S.D 5%			0.74					0.57				
	Cut-off				In	rigat	ion treat	ment 2007					
σ	irrigation		CF				6 day	s	12 d	ays			
jel (ACH		15.6	51			12.92	2	11.3	35			
2 er	1-WACH		14.8	37			13.90)	12.1	18			
t a			14.5	01			14.22		12.4	49			
S	4-WACH		15.4	12 17			14.20		12.0	30			
			10.1				1 0.22		10.	50			
	2.0.0 070					Irri	nation tr	atment					
	Cut-off			2007			gation ite	aunent	2008				
еX	irrigation	CE	. 1	6 days		12	dave	CE	6 days	12 days			
pu		0.20	22	0 424		- 12	402	0.416	0.416	0 202			
st	1-WACH	0.30	6	0.424		0.	392	0.410	0.410	0.393			
Xe.	2-WACH	0.4	14	0.434		0	402	0.458	0.461	0.405			
lar	3-WACH	0.43	37	0.449		0.	.399	0.446	0.473	0.422			
Т	4- WACH	0.43	38	0.468		0	.376	0.464	0.468	0.404			
				0.005					0.016				

CF=continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation.

Milling recovery and protein content were significantly responded to different harvest times in both seasons (Table 4). The lowest values were recorded when plants were harvested 10-DACI as compared with harvesting times at 15- and 20-DACI with insignificant differences between each other for most of traits in both seasons. The relatively lower milling recovery at first harvest times is probably due to the presence of unripe, light and deformed grains. The results are in agreement with those reported by Ntanos *et al.* (1998) and Surek *et al.* (1998).

Data in Table 5 indicated that CF with irrigation termination 3-WACH recorded the highest percentage of hulling, milling and head rice in 2007.

Table 4:	Gra	in quality ch	nara	cters of Eg	yptian hyb	rid rice	cultivar (El	HR1)
	as har	influenced vest times.	by	irrigation	intervals,	cut-off	irrigation	and

Treatment	Hulling (%)		Mil (?	Milling (%)		d rice %)	Amy con (؟	/lose itent %)	Protein content (%)		
	2007	2008	2007	2008	2007	2008	2007	2008	2007	2008	
Irrigation intervals											
(1):											
CF	76.61	77.61	70.84	71.17	62.12	61.47	18.67	19.47	7.23	7.31	
6 days	75.85	77.30	70.50	70.35	61.91	61.04	19.60	19.78	7.19	7.20	
12 days	74.99	76.78	68.32	69.77	59.66	59.31	19.81	20.15	6.76	6.91	
L.S.D 5%	0.55	NS	1.03	0.62	1.44	1.25	0.63	0.60	0.08	0.26	
Cut-off irrigation (C):											
ACH	74.06	76.21	66.92	69.38	56.26	59.10	18.74	20.03	6.87	6.87	
1-WACH	75.93	75.98	70.08	70.37	61.91	59.49	19.99	20.31	7.04	6.90	
2-WACH	75.98	77.44	70.81	71.14	62.84	61.68	19.25	19.86	7.02	7.08	
3-WACH	76.58	78.26	70.53	70.48	62.77	61.94	19.66	19.20	7.16	7.44	
4- WACH	76.52	78.26	71.09	70.78	61.31	60.82	19.06	19.61	7.22	7.42	
L.S.D 5%	0.86	0.97	0.99	1.09	1.73	1.13	0.80	0.71	0.19	0.23	
Harvest times (H):											
10-DACI	75.07	76.64	68.45	69.75	59.67	59.08	19.37	19.62	6.97	7.08	
15 DACI	75.86	77.51	70.21	70.93	60.65	61.37	19.48	19.89	7.05	7.18	
20-DACI	76.52	77.54	70.99	70.61	62.72	61.37	19.22	19.90	7.17	7.17	
L.S.D 5%	0.65	0.76	0.87	0.76	1.62	0.80	NS	NS	0.13	NS	
Interaction											
IxC	*	NS	*	NS	**	**	*	**	**	**	
IxH	NS	NS	NS	NS	**	NS	NS	NS	**	*	
СхН	**	NS	*	**	**	**	NS	NS	**	**	
IxCxH	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	

CF=continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation. NS = Not significant, * and ** = Significant at 5 and 1 % levels, respectively.

	Cut-off	l li	Irrigation treatment (2007) Harvest time (2007)												
_	irrigation	С	F	6	days	12 d	lays		10-E	DACI	15-D	DACÌ	20)-D	ACI
%)	ACH	75	.58	7	4.41	72	.20		71	.56	74.	.82	-	75.	82
6	1-WACH	75	.95	7	5.84	76	.02		75	75.31		.01		76.49	
llin	2-WACH	76	.92	7	5.90	75	.12		75.43		75.57			76.94	
P	3-WACH	77	77.94		6.64	75.17			76.32		76.56			76.87	
-	4- WACH	76	.66	7	76.45		.43		76	.73	76.35			76.47	
	L.S.D 5%				1.40	-					1.4	45			
		Irri	igatior	n inte	erval					Harves	st time	;			
	Cut-off		20	07				20	07			2	800		
-	irrigation	CF	6 d	avs	12 davs	10-		1	5-	20-	10-		15-	_	20-
%)		•••	• •	uyo		DAC		DA		DACI	DAC	I D	ACI	D	DACI
βι	ACH	68.85 67.26		.26	64.64	63.47	7 6	<u>58</u> .	.04 6	59.23	68.62	2 6	3.84	7	0.68
illi	1-WACH	70.13	3 70	.40	69.70	69.05	5 7	70.	.18	71.00	69.01	1 7	1.50	7	0.58
Μ	2-WACH	/1.5	1 /1	.63	69.27	69.68	3 1	/1.	.36	(1.37	/1./9	9 7	1.58		0.05
	3-WACH	72.3	1 /1	/1.43 67.8		69.04	1 /	70. 70	.88	(1.68	69.05	$\frac{1}{2}$	1.72	1	0.66
	4- WACH	/1.3	((1)	.79	70.11	71.07	1 1	10.	.57	/1.69	70.30) /	1.00	1	0.05
	L.S.D 5%		1.	/5		<u> </u>		1.8	86			1	.74		
			val					Harves	st tim	e					
	Cut-off	2007				2008			10- 15-			40	200	8	
	irrigation	CF	6D	12 [CF	6D	12	D		15- DACI	20-	10-	15	-	20-
		60.65	57 21	50.0	1 61 22	59.22	577	75	54 62	54 16	50.00	58 G	1 50 (7	50 50
		50.00	62 35	50.9 64 0	8 60 37	60.22	57.7	73	54.02 61 75	61 83	59.99 62 14	58.0	+ 59.0	20	50 61
%)	2-WACH	60 17	64.06	64.2	9 62 14	61 55	61 3	25	61 02	64 24	63.26	60.80	161	76	62.38
e	3-WACH	66.34	62 73	59.2	3 61 68	62 46	61 F	38	59.99	63.00	65.31	59.9	1 62 8	33	63 10
ri	4- WACH	64.13	59.98	59.8	2 61.84	62.57	58.0)7	60.98	60.02	62.93	57.0	0 63.3	30	62.18
ad	L.S.D 5%		2.93			3.02				2.71			1.8	2	
Ψ	Irrigation					Har	vest	: ti	me (2	007)					
	treatment		1 0- D	DACI			15	5-D	ACI			20-	DAC		
	CF		61	.53			6	62.	.77		62.05				
	6 days		59	.96			5	59.	.76		64.07				
	12 days		57	.52		59.42					62.05				
	L.S.D 5%							2.0	05						

 Table (5): Milling recovery of Egyptian hybrid rice cultivar (EHR1) as influenced by the interaction among the studied factors.

However, 6D treatment gave the highest head rice percentage with harvest time 20-DACI in 2007 and with terminates irrigation 4-WACH in 2008. Cut-off irrigation 2-WACH recorded the highest hulling and milling (%) when plants were harvested 20- and 10-DACI in 2008, respectively. On the other hand, cut-off irrigation 3-WACH with harvesting plants 20-DACI recorded the highest milling (%) in 2007 and head rice (%) in both seasons.

Results in Table 6 revealed that the highest values of amylose content were given by 6D with cut-off irrigation 1- and 2-WACH in 2007 and with cut-off irrigation after complete heading in 2008. However, the highest protein content was given by CF with cut-off irrigation 2- and 4-WACH in both seasons, respectively. Continuous flooding with harvest plants 15- and 20-DACI recorded the highest protein content in 2007 and 2008, respectively. The highest protein content was obtained when cut-off irrigation 3- and 4-WACH with harvest plants 20-DACI in both seasons.

CF=continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation.

				Irrigation	n interval								
tent	Cut-off irrigation		2007			2008							
nte		CF	6 days	12 days	CF	6 days	12 days						
8	ACH	16.98	19.45	19.78	19.73	21.04	19.33						
e	1-WACH	19.09	20.58	20.31	20.02	20.51	20.41						
ő	2-WACH	18.83	20.58	20.34	19.19	19.98	20.41						
л Х	3-WACH	19.05	19.61	20.30	19.13	18.25	20.23						
Ar	4- WACH	19.38	19.46	18.31	19.30	19.14	20.39						
	L.S.D 5%		1.34			1.20							
				Irrigation	n interval								
	Cut-off irrigation		2007			2008							
		CF	6 days	12 days	CF	6 days	12 days						
	ACH	7.15	6.86	6.60	6.69	7.15	6.78						
	1-WACH	6.98	7.46	6.67	7.44	6.41	6.85						
	2-WACH	7.64	6.70	6.71	7.06	7.21	6.99						
	3-WACH	7.36	7.47	6.66	7.60	7.70	7.01						
	4- WACH	7.01	7.48	7.16	7.79	7.54	6.93						
ц	L.S.D 5%	0.30 0.42											
tei	Irrigation	Harvest time											
lo	treatment		2007		2008								
С С	liealinein	20-DACI	15-DACI	10-DACI	20-DACI	15-DACI	10-DACI						
iei	CF	7.01	7.35	7.33	7.23	7.24	7.47						
ē	6 days	7.27	6.99	7.33	7.14	7.18	7.29						
Ъ.	12 days	6.63	6.80	6.84	6.87	7.13	6.73						
	L.S.D 5%		0.19			0.33							
	Cut-off irrigation	20-DACI	15-DACI	10-DACI	20-DACI	15-DACI	10-DACI						
	ACH	6.81	6.80	7.00	7.21	6.45	6.95						
	1-WACH	7.28	6.94	6.89	6.61	7.24	6.85						
	2-WACH	6.86	7.32	6.86	6.87	7.16	7.22						
	3-WACH	6.83	7.15	7.52	7.60	7.31	7.40						
	4- WACH	7.07	7.02	7.52	7.11	7.75	7.41						
	L.S.D 5%		0.30			0.39							

Table 6: Amylose and protein contents of Egyptian hybrid rice cultivar (EHR1) as influenced by the interaction between the studied factors.

CF=continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation.

Water relations:

The amounts of water input, before starting irrigation treatments, were 4270 and 4410 m³/ha, in 2007 and 2008 seasons, respectively, (Table 9). Nour and Mahrours (1994) indicated that the amount of water used in land preparation for transplanting rice was 4525.5 m³/ha. Comparing the different treatments of irrigation (Table 9), it was observed that CF received the highest amounts of water throughout the season, while, the lowest amounts were received by 12D treatment. Data showed, also, that increasing irrigation intervals from CF up to six and twelve days tended to decrease the amount of water used from 14656 and 15324 m³/ha to 13476 and 14183 m³/ha and 13476 and 14183 m³/ha in both seasons, respectively. There were no large variations in the amounts of irrigation water input due to the stable conditions (temperature, relative humidity and evaporation rates) in both studied seasons as previously shown in Table (1). El-Refaee *et al.* (2007) found that under continuous flooding rice field received the highest amounts of water

throughout the season while, the lowest amount was received by irrigation every 8 days.

Under CF, delayed the time for cut-off irrigation from complete heading to 4-WACH increased water used from 13380 and 14032 m³/ha to 16011 and 16477 m³/ha, however, it increased from 12382 and 13077 m³/ha to 14750 and 15004 m³/ha with 6D and from 9658 and 10332 m³/ha to 11083 and 11856 m³/ha with 12D in the both seasons, respectively. These mainly due to continuous applied of water for more than four weeks with delayed the time of cut-off irrigation.

Data in Table 9 demonstrated that, over cut-off irrigation treatments, water saved due to increasing irrigation intervals compared to CF were (8.04 and 7.08 %) and (29.27 and 27.30 %) with corresponding grain yield reduction of (4.54 and 3.64 %) and (28.2 and 25.8 %) for 6 and 12D in the two successive seasons. Ebaid and El-Refaee (2007) pointed out that water saved due to increasing irrigation interval 8 and 12 days, compared to continuous flooding, were 11.2 and 20.5 % with corresponding grain yield reduction of 11.0 and 24.9 %, respectively. Results showed, also, that cut-off irrigation saved some of irrigation water with different values of reduction in grain yield under both of irrigation every 6 and 12 days as compared with the same treatments under CF in both seasons. Delay in cut-off irrigation significantly increased the grain yield % (Table 7). Compared to cut-off irrigation after complete heading, cut-off irrigation 3-WAH gave the highest percentage of increasing grain yield of 28.47 and 24.00 % under CF, however, cut-off irrigation 4-WAH increased grain yield by 22.33 and 22.99 % under 6D in both season, respectively. Under 12D the highest value of increasing grain yield was recorded with cut-off irrigation 2-WACH (9.46 5) in the first season and 3-WACH (23.98 %) in the second season.

Regarding water productivity (WP), 6D was considered the best water productivity (0.796 and 0.798 kg/m³) followed by 12D (0.784 and 0.784 kg/m³). However, CF recorded the lowest values (0.767 and 0.767 kg/m³) in the first and second seasons, respectively (Table 9). Bouman and Tuong (2001) reported that WP was higher in alternate submerged and non-submerged regimes than in the continuous submerged regime. The WP values were relatively high in irrigation every 6 days compared with the other irrigation intervals. The extremely high values of WP, in this treatment, were caused by the extremely high grain yield and low water inputs in this treatment

Under all irrigation intervals delayed cut-off irrigation recorded higher WP than cut-off irrigation ACH. High WP, with delayed cut-off irrigation, was associated with high grain yield. The productivity of irrigation water could be increased to reach it maximum value of 0.798 and 0.881 kg/m³ in first season and 0.794 and 0.795 kg/m³ in second season under CF, and about 0.820 and 0.826 kg/m³ in first season and 0.817 and 0.810 kg/m³ in second season when irrigation intervals increased up to six days with cut-off irrigation 2- and 3- WACH, respectively. However, the highest WP under 12D was obtained when irrigation terminated 2-WACH (0.822 kg/m³) in first season and 3-WACH (0.851 kg/m³) in second season.

			20	07			
Irrigation	Cut-off	Grain	Grain yield	Grain yield	Total water	Water	WP
treatment	irrigation	(t/ha)	reduction (%)	increased (%)	used (m³/ha)	saved (%)	(kg/m³)
	ACH	9.677	-	-	13380	-	0.723
	1-WACH	10.527	-	8.78	14013	-	0.751
CF	2-WACH	11.617	-	20.05	14555	-	0.798
	3-WACH	12.432	-	28.47	15320	-	0.811
	4-WACH	12.002	-	24.03	16011	-	0.750
Average		11.251	-	20.33	14656	-	0.767
	ACH	9.546	1.35	-	12382	7.46	0.771
c	1-WACH	9.919	5.78	3.91	12847	8.32	0.772
o dave	2-WACH	10.892	6.24	14.10	13290	8.69	0.820
uays	3-WACH	11.664	6.18	22.19	14113	7.88	0.826
	4-WACH	11.678	2.70	22.33	14750	7.87	0.792
Average		10.740	4.54	15.63		8.04	0.796
	ACH	7.688	20.55	-	9658	27.82	0.796
12 days	1-WACH	7.868	25.26	1.00	9939	26.92	0.792
	2-WACH	8.415	27.56	9.46	10241	29.64	0.822
	3-WACH	8.385	32.55	9.07	10540	31.20	0.786
	4-WACH	8.025	33.14	4.38	11083	30.78	0.724
Average	-	8.076	28.22	5.98	10292	29.27	0.784
		•	20	08			
	ACH	10.159	-	-	14032	-	0.724
	1-WACH	11.459	-	12.80	14901	-	0.769
CF	2-WACH	12.199	-	20.08	15364	-	0.794
	3-WACH	12.597	-	24.00	15846	-	0.795
	4-WACH	12.387	-	21.93	16477	-	0.752
Average		11.760	-	19.70	15324	-	0.767
- U	ACH	9.808	3.46	-	13077	6.81	0.750
<u>_</u>	1-WACH	11.183	2.41	14.02	13820	7.25	0.809
0 Have	2-WACH	11.646	4.53	18.55	14249	7.26	0.817
days	3-WACH	11.958	5.07	21.92	14763	6.24	0.810
	4-WACH	12.063	2.62	22.99	15004	7.82	0.804
Average		11.332	3.64	19.37	14183	7.08	0.798
- U	ACH	7.889	22.34	-	10332	26.37	0.764
	1-WACH	8.834	22.91	11.98	10812	27.44	0.817
12 days	2-WACH	8.680	28.85	10.03	11191	27.16	0.776
-	3-WACH	9.781	22.35	23.98	11494	27.46	0.851
	4-WACH	8.453	31.76	7.15	11856	28.05	0.713
Average	•	8.727	25.79	13.29	11137	27.30	0.784

 Table 7: Grain yield, yield reduction, grain yield increased, total water used, water saved and water productivity (WP) of EHR1rice cultivar as influenced by irrigation treatments.

CF=continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation.

Generally, in case of water shortage, it could be concluded to use irrigation every six days and cut-off irrigation 3-WACH with harvest plants 10 DACI for the highest water productivity and grain yield as well as acceptable grain quality characters of Egyptian hybrid rice cultivar (EHR1).

REFERENCES

- A.O.A.C. (1970). Official Methods of Analysis. 11th Ed. Washington, D.C., USA.
- Asano, H.; F. Hirano; K. Isobe and H. Sakurai (2000). Effect of harvest time on the protein composition (glutelin, prolamin, albumin) and amylase content in paddy rice cultivated by Aigamo duck farming system Japanese. J. Crop. Sci. (59): 320-323.
- Awad, H. A. (2001). Rice production at the North of Delta region in Egypt as affected by irrigation intervals and nitrogen fertilizer levels. J. Agric. Sci. Mansoura Univ., 26(2): 1151-1159.
- Badawi, A. T.; M. A. Maximos and I. R. Aidy (2002). Rice improvement in Egypt during 85 years (1917- 2001). In: Rice in Egypt. Ministry of Agriculture and Land Reclamation. ARC, Egypt. 1-24.
- Belder, P.; J. H. J. Spiertz; B. A. M. Bouman; G. Lu and T. P. Tuong (2005). Nitrogen economy and water productivity of lowland rice under watersaving irrigation. Field Crops Res. 93:169-185.
- Bouman, B. A. M. and T. P. Toung (2001). Field water management to save water and increase its productivity in irrigated lowland rice. Agric. Water Manage., 49: 11-30.
- Dewedar, G. A. A. (2004). Effects of some an agricultural practice on yield, yield components and some technological characters. M. Sc. Thesis, Face. Agric., Al-Azahr Univ., Egypt.
- Ebaid, R. A. and I. S. El-Refaee (2007). Utilization of rice husk as an organic fertilizer to improve productivity and water use efficiency in rice fields. African Crop Science Conference, El-Minia Univ., October, 27-31, (8):1923-1928.
- Elaine, T. C.; K. L. Bett-Garber; J. Thompson; R. Mutters; C. C. Grimm and A. M. McClung (2005). Effects of drain and harvest dates on rice sensory and physicochemical properties. Cereal Chemistry 82(4):369-374.
- El-Gewaily, E. E. A. (2006). Behavior of some nutrient elements in rice soils under different irrigation intervals. Ph.D Thesis Dept. Fac. Agric., Mansoura Univ., Egypt.
- El-Refaee, I.S. H. (1997). Effect of some irrigation treatments on growth and yield of rice. M.Sc. Thesis, Fac. of Agric., Kafr El-Sheihk, Tanta Univ., Egypt.
- EI-Refaee, I. S. H. (2007). Effect of cut–off irrigation date on grain yield and quality of some Egyptian rice cultivars. J. Agric. Res. Kafr El-Sheikh Univ., 33(1):123-139.
- El-Refaee, I. S.; A. E. Abd El-Wahab; F. N. Mahrous and S. A. Ghanem (2007). Irrigation management and splitting of nitrogen application as affecting on grain yield and water productivity of hybrid and inbred rice. African crop science conference, El-Minia Univ., October, 27-31, (8):45-52.

- Gomez, K. A. and A. A. Gomez (1984). Statistical Procedure for Agricultural Research. 2nd Edition. A Wiley Interscience Pub., John Wiley and Sons, New York. USA. Pp.628.
- Govindaswamy, S. and A. K. Ghosh (1970). Time of harvest for high head rice recovery. Oryza 7:95-96.
- Guerra, L. C.; S. I. Bhuiyan; T. P. Tuong and R. Barker (1998). Producing more rice with less water from irrigated systems. SWIM Paper 5. IWMI/IRRI, Colombo, Sri Lanka, 24 p.
- Hossain, M. F.; M. S. U. Bhuiya; M. Ahmed and M. H. Mian (2009) Effect of harvesting time on the milling and physicochemical properties of aromatic rice. Thai J. of Agric. Sci., 42(2): 91-96
- Juliano, B. O. (1971). A simplified assay for milled rice amylose. Cereal Sci. Today (16): 334-338,340-360.
- Lin, J. Y. (1994). Impact of hybrid rice on input demand and productivity. Agricultural Economics. 10(2):153-164.
- Nour, M. A. M. and F. N. Mahrours (1994). Effect of varying irrigation intervals during tillering, reproductive and ripening stages on rice yield and its components. Egypt J. Appl. Sci., 9(7):869–879.
- Nour, M. A. M.; S. A. Ghanem and A. M. El–Serafy (1994). Effect of different water regime on the productivity of some transplanted rice. Proc. 6th Conf. Al–Azhar Univ., Cairo, Egypt, (1):247–259.
- Ntanos, D.; N. Pillippou and S. H. Zinoviad (1998). Effect of rice harvest on milling yield and grain breakage. Cahiers Optios Mediterraneennes, 15(1):23-27.
- Slaton, N. A. (2001). Rice production handbook. University of Arkansas Cooperative Extension Service MP192. Little Rock, Ark., USA.
- Surek, H.; H. Aydin; M. Negis and H. Kushu (1998). Determining the suitable rice harvesting time in Erdine. Cahiers Options Mediterraneenes. 15(1): 29-31.
- Uppal, H. S. and A. S. Bali (1994). Effect of scheduling and cut-off date of irrigation on yield and quality of rice (*Oryza sativa*) in a non-cracking soil. Indian J. Agron., 39 (4):628-630.
- Virmani, S. S. (2002). Development and large scale adoption of hybrid rice in India. In: India Grains. pp 15-16.
- Williams, V. R.; W. T. Wu; H. Y. Tsai and H. G. Bates (1958). Varietal differences in amylase content of rice starch. J. Agric. Foodchem. 6: 47-48.
- Yoshida, S. (1981). Fundamentals of rice crop sciences. International Rice Research Institute, Los Banas, Philippines. pp. 269.
- Zhi, M. (1993). Environmental impact of water-saving irrigation for rice. Proceeding of Environmental Sound Water Resource Utilization, AIT Bangkok, Thailand, 8-11 Nov., p 143-148.

سلوك صنف الأرز الهجين "هجين مصرى واحد" تحت نظم رى و مواعيد حصاد مختلفة

- عبد الحميد محمد عمر ' ، اسماعيل سعد الرفاعي ' و حسناء عبد الحميد غازي ' ' قسم المحاصيل – كلية الزراعة – جامعة كفر الشيخ
- ٢ مركز البحوث و التدريب في الأرز معهد بحوث المحاصيل الحقلية مركز البحوث الزراعية-
 - سخا- كفر الشيخ مصر

أجريت هذه الدراسة فى المزرعة البحثية لمركز البحوث و التدريب فى الأرز – سخا – كفر الشيخ – مصر - خلال موسمي صيف ٢٠٠٧ و ٢٠٠٨ م بهدف دراسة تأثير زيادة فترات الرى و مواعيد توقف الرى و الحصاد على محصول الحبوب و مكوناتة و على جودة الحبوب و بعض العلاقات المائية لصنف الأرز الهجين "هجين مصرى واحد". إستخدم تصميم القطع المنشقة مرتين ذو أربع مكررات بحيث إحتوت القطع الرئيسية على ثلاث معاملات للري هي الغمر المستمر و الري كل ستة أيام و اثنى عشر يوماً و إحتوت القطع الشقية الأولى على خمسة مواعيد لتوقف الرى و هى توقف الرى بعد تمام الطرد و ١ و ٢ و ٢ و ٢ تمام الطرد. فى حين إحتوت القطع الثانية على ثلاثة مواعيد للحصاد وهى: الحصاد بعد ١ و ٢ و ٢ و ٢ و ٢ و ٢ و ٢ و ٢ يوماً من ميعاد توقف الرى.

و توضح أهم النتائج أن محصول الحبوب و مكوناتة بالإضافة الى صفات التبيض و محتوى البروتين قد نقصت معنويا بزيادة فترات الرى من الغمر المستمر حتى الرى كل ١٢ يوماً، بينما أعطت معاملة الرى كل ١٢ يوماً أعلى القيم لنسبة الحبوب الفارغة و محتوى الأميلوز. فى حين سجلت معاملة الرى كل ستة أيام قيما وسطاً مع عدم وجود فروق معنوية بينها و بين معملة الغمر المستمر فى معظم الصفات السابقة, فى كلا الموسمين.

أدى التأخير في توقف الري حتى ٣ و ٤ أسابيع من تمام الطرد إلى الزيادة المعنوية فى محصول الحبوب و معظم مكوناتة و كذلك صفات الجودة، في حين أدى تأخير توقف الري إلى النقص المعنوي في محتوى الاميلوز فى موسم ٢٠٠٧ و نسبة الحبوب الفارغة فى الموسمين و ذلك بالمقارنة بتوقف الري مبكرا بعد تمام الطرد. كما سجل ميعادي الحصاد بعد ١٠ و ١٥ من توقف الري زيادة معنوية فى عدد السنيبلات/دالية، وزن الدالية ، وزن ١٠٠٠ حبة، السعة المحصولية، كثافة الدالية و محصول الحبوب في بالموسمين. بينما سُجلت أقل القيم لصفات التبيض و محتوى البروتين عند الحصاد بعد ١٠ و بالمقارنة بياقى مواعيد الحصاد الأخرى.

استهلكت معاملة الغمر المستمر طوال الموسم أكبر كميه مياه رى فى حين قلت كميات مياه الرى المستهلكة بزيادة فترات الرى إلى ٦ و ١٢ يوماً. علاوة على ذلك سجلت معاملة الرى كل ٦ أيام أعلى إنتاجية لوحدة المياه (٢٩٦، و ٢،٧٩٨ كجم/م٢) و أقل نقص فى محصول الحبوب (٢،٤٤ و ٣،٦٤ %) و وفرت كمية من مياه الرى تعادل ٢،٠٤ و ٢،٠٧ % و ذلك بالمقارنة بمعاملة الغمر المستمر فى كلا الموسمين، على التوالى. كما أدى التأخير فى توقف الرى الى زيادة إنتاجية وحدة المياه و ذلك بالمقارنة بتوقف الرى بعد تمام الطرد.

و بصفة عامة، بناء على النقص في مياه الري, يمكن الإستنتاج بأن إستخدام معاملة الرى كل ستة أيام مع توقف الرى بعد ثلاثة أسابيع من الطرد و الحصاد بعد عشرة أيام من توقف الري, أدى الى الحصول على أعلى إنتاجية لمحصول الحبوب و وحدة المياه بالإضافة إلى صفات جودة مقبولة للحبوب لصنف الأرز الهجين "هجين مصرى واحد".

قام بتحكيم البحث

أ.د / عبد الرحيم عبد الرحيم ليله أ.د / احمد عزت عبد الوهاب

كلية الزراعة – جامعة المنصورة مركز البحوث الزراعية

J. Plant Production, Mansoura Univ., Vol. 3 (3), March, 2012

Treatment	eatment No. of grains/ panicle		Unfilled grains (%)		Panicle weight (g)		1000 grain weigh (g)		Grain yield (t/ha)		Straw yield (t/ha)		Harvest index	
	2007	2008	2007	2008	2007	2008	2007	2008			2007	2008	2007	2008
rrigation intervals (I):														
CF	191.9	192.5	9.1	10.8	4.00	3.91	24.52	24.18	11.25	11.76	15.18	14.51	0.425	0.447
5 days	175.3	183.5	13.3	12.6	3.62	3.60	24.68	23.80	10.74	11.33	13.71	13.62	0.439	0.454
12 days	139.6	162.5	14.8	15.3	2.52	3.27	22.58	22.78	8.08	8.73	12.39	12.62	0.394	0.408
S.D 5%	6.8	6.8	1.3	1.5	0.20	0.40	0.54	0.73	0.44	0.38	0.59	0.65	0.016	0.008
Cut-off irrigation (C):														
АСН	170.5	178.9	17.0	22.0	3.02	2.88	23.12	22.71	8.97	9.29	13.29	13.36	0.404	0.408
1-WACH	175.0	178.3	12.8	15.6	3.24	3.19	23.77	23.19	9.44	10.49	13.65	13.46	0.409	0.437
2-WACH	173.6	185.7	11.4	10.6	3.55	4.00	24.13	23.98	10.31	10.84	13.74	13.60	0.427	0.441
3-WACH	168.4	184.7	10.5	8.2	3.49	4.14	24.40	24.40	10.83	11.45	14.10	13.96	0.431	0.450
4- WACH	157.1	170.6	10.2	8.0	3.61	3.75	24.39	23.66	10.57	10.97	14.00	13.54	0.427	0.446
S.D 5%	5.3	7.5	1.2	1.7	0.21	0.25	0.49	0.74	0.42	0.31	NS	NS	0.014	0.009
<u>Harvest date (H):</u>														
10-DACI	171.0	184.9	12.8	15.1	3.52	3.69	24.24	23.88	10.23	10.70	13.98	13.72	0.420	0.437
15 DACI	171.6	181.0	13.1	12.7	3.41	3.62	23.90	23.50	10.02	10.77	13.79	13.58	0.419	0.440
20-DACI	164.2	173.0	11.3	10.9	3.21	3.47	23.74	23.38	9.82	10.35	13.51	13.46	0.419	0.432
S.D 5%	0.46	6.2	0.9	1.4	0.14	0.18	0.39	0.45	0.28	0.31	NS	NS	NS	NS
nteraction														
xC	NS	**	**	NS	NS	NS	NS	NS	**	**	**	NS	**	**
хН	NS	NS	**	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
CXH	**	NS	**	**	*	NS	NS	NS	NS	NS	NS	NS	NS	NS
хСхН	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 2: Grain yield and some of its attributes of Egyptian hybrid rice cultivar (EHR1) as influenced by irrigation intervals, cut-off irrigation and harvest times.

CF= Continuous flooding, ACH= After complete heading, WACH= Week after complete heading and DACI = Days after cut-off irrigation. NS = Not significant, * and ** = Significant at 5 and 1 % levels, respectively.